

[12670-5]

ACCEPTED

APR 13 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 12670-5
SUBJECT TO THE FOLLOWING COMMENTS

DIRECTIONS FOR USE

For most effective results use in a good sprayer adjusted to deliver a fine mist.

MOSQUITOES, FLIES, GNATS: Close all doors and windows. Direct the spray to all parts of the room. Fill the room with mist, then leave the treated area. Keep room closed for at least 15-30 minutes. Ventilate the room when treatment is completed.

WASPS: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary.

FOR CRAWLING INSECTS: Roaches, earwigs, grain mites, sowbugs, spiders and centipedes: Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insect where possible.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

E.P.A. REG. NO. 12670-5

Quality is Our Most Important Product



PEST-CIDE

FOR INDOOR USE ONLY

FOR: Bakeries, Food Plants, Restaurants, Schools, Hotels, Motels, Office Buildings, Hospitals and Food Handling Plants.

NET CONTENTS 1 GALLON

CAUTION: KEEP OUT OF REACH OF CHILDREN.

SEE OTHER CAUTIONS ON BACK PANEL

SOLD EXCLUSIVELY BY

GLOBE CHEMICAL CO., INC.

DECATUR, GEORGIA

ACTIVE INGREDIENTS:

*†(5-Benzyl-3-furyl)methyl, 2,2-dimethyl-3-(2-methylpropenyl)cyclopropanecarboxylate	0.250% w/w
Related compounds	0.034%
Aromatic petroleum hydrocarbons	0.331%
Petroleum distillate	99.375%
INERT INGREDIENTS:	0.010%
	<u>100.000%</u>

*Cis, trans isomers ratio: max. 30% (±) cis and min. 70% (±) trans.

†U.S. Patent No. 3,465,007

CAUTION:

KEEP OUT OF REACH OF CHILDREN.
Harmful if swallowed. Do not store near heat or open flame. Avoid contamination of feed and foodstuffs. Use in food processing plants only when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. In hospitals, patients should be removed from room prior to treatment. Room should be ventilated for 2 hours after spraying. Do not return patients to room until after ventilation.

Remove pets, birds, and cover fish aquariums before spraying.