IKE-CHLOR POOL SANITIZER ' (Chlorinating Solution)

ACTIVE INGREDIENT: SODIUM HYPOCHLORITE......9%

KEEP OUT OF REACH OF CHILDREN

DANGER

FIRST AID: If on Skin, wash with plenty of snap and water, if in eyes, thish with water for at least 15 minutes. Get medical attention, if swallowed, drink large quantities of milk, or gelatin solution or, if these are not available, drink large quantities of water. Do not give venegal or other acids. Do not induce vomiting. Get prompt medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive, may cause severe skin irritation or chemical burns to broken skin. Causes eye damage. May be fatat if swallowed, Avoid breathing vapors. Do not get in eyes, on skin or un clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash hands after handling, Vacate poorly ventilated areas as soon as possible. Do not return until odors have gissipated.

ENVIRONMENTAL HAZARDS. This pesticide is toxic to fish, keep out of lakes, streams or ponds. Treated effluent cannot be discharged into lakes, streams, ponds or public waters unless a discharge permit is obtained. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS: STRONG OXIDIZING AGENT: Mix only with water according to label directions. Mixing this product with gross filth such as feces, urine, etc., or with ammonia acids, detergents or other chemicals will release hazardous gases infitating to eyes, lungs and mucous membranes.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

it is a violation of federal law to use this product in a manner inconsistant with its labeling.

DIRECTIONS FOR SWIMMING POOL CHLORINATION

For a new pool or for spring start-up, superchlorinate with one (1) pint (16 flu. ozs.) of sodium hypochlorite solution for each 2500 gallons of water. This dosage is equivalent to 5 ppm available chlorine by weight.

by weight.

FOR POOL MAINTENANCE

A) Adjust pool water pH to 7 6 7.2 range and maintain.

B) Add manually or by a feeder device, sodium hypochlorite solution at a rate to maintain an available chlorine residual of 0.6 to 1.0 ppm. One half pint (8 flu, nzs.) of sodium hypochlorite solution for each 6000 gallons of water will provide 1.0 ppm available chlorine by weight. Frequency of additions to maintain 0.6 to 1.0 ppm available chlorine will depend on temperature and number of swimmers. Use test kit to make certain the pH and chlorine residual are in the proper range.

NOTAL: This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available

STORAGE AND DISPOSAL: Store in a cool, dry area away from direct sunlight. In case of spill, flood area with large quantities of water. Rinse empty container thoroughly with water and either return to manufacturer or discard by placing in trash collection or burying in an approved land fill. Product or rinsate that cannot be usad, should be diluted with water and disposed of in a sanitary sewer. Do not contaminate food or feed by storage, disposal or cleaning or equipment.



1290 E. Cummercial Blvd.: Fort Lauderdale, Florida 33306 / 771-5700 *Detray Beach 421-1711

EPA REG NO 17-11-47 EPA EST. NO. 12711 FL 01 HET CONTEN S 25 GALLONS

DATE PACKAGED

NOT FUR USE AFTER

ACCEPTED PAY 2 1980

UNDER THE FELERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER-ID UNDER NO.

30