10 POUNDS

1/2% CHLORPYRIFOS GRANULES

Controls chinch bugs, sod webworms (lawn moths), ants, brown dog ticks, crickets, cutworms, earwigs, sowbugs (pillbugs), certain white grubs1, and hyperodes (turf grass) weevils2 infecting lawns and other ornamental turf areas.

EPA Reg. No. 13283-8 EPA Est. No. 13283-AL-1

ACTIVE INGREDIENT

Chlorpyrifos (0.0-diethyl 0-[3,5,6-trichloro-2-pyridyl]

INERT INGREDIENTS 99.6%

TOTAL ...

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately. IF INHALED: Remove victim to fresh air and apply respiration if indicated. IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly. IF IN EYES: Rinse eyes for at least 15 minutes with water. Call-a physicia CAUTION CEPTED immediately.

KEEP OUT OF REACH OF CHILDREN
See back panel for additional preceduring statements

Unant the Federal Impedicide, Fungities, and Redenticide Act as unsaich, for the posticide tegislored under EPA Reg. No.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its

FOR CONTROL OF CHINCH BUGS, SOD WEBWORMS (LAWN MOTHS), ANTS, BROWN DOG TICKS, CRICKETS, CUTWORMS, EARWIGS, AND SOWBUGS (PILLBUGS):

Apply Do-it-Yourself Insect Control granules uniformly to home lawns and other ornamental and recreational turf grass areas, using any hand or power or election spreader (gravity or rotary type) for applying granular materials. Treat when insects

rea to be eated	Amount of granules	Amount of water	first appear. Apply Insect Control at rate of 10 pounds of granules per 2,000 sq. ft. For more uniform distribution, apply one half.
200 sq ft	1 lb	6 gal	the required amount of granules in one direction, then apply the remaining balf in one direction at a right angle, to the
.000 sq ft	5 lbs	30 gal	
.000 sq ft	10 lbs	60 gal	

remaining half it *angles*te ake direction of the first application. Immediately after, sprinkle turf with 60 gallons of water per 2,000 sq. ft., so as to wash the granules into the zone of insect intestation. Use chart to determine amounts of insect Control granules and water required for smaller areas. Re-treat as necessary

FOR CONTROL OF CERTAIN WHITE GRUBS (LARVAE OF **EUROPEAN CHAFER AND JAPANESE BEETLE):**

Use the recommended amount of granules for the specified area. Apply granules uniformly over the area treated and immediately water the area thoroughly. For best ing and surface. August.

Area to be rreated		Amount of water	results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service for proper treatment time in your area.
200 sq ft 1 000 sq fi 2 000 sq fi	2 4 lbs 10 20 lbs 20 40 lbs	30 gat	

FOR CONTROL OF HYPERODES (TURF GRASS) WEEVILS:

Use the recommended amount of granules for the specified area shown in chart

Amount to be treated	Amount of granules	Amount of water	
200 sq ft	2 lbs	6 gais	
1 000 sa ft	10 lbs	6 gals 30 gals	
2 2000 sq. ft	20 4bs	60 gals	

Treat suspected problem areas in mid-April and again in mid-July. Consult your local Agricultural Extension Service for the best treatment time in your area

FOR CONTROL OF ANTS IN BURIED CABLE CLOSURES: Apply 3 ounces of granules per square foot of enclosed surface area. This 3-ounce package is specifically intended for use in buried cable closures of power and utility companies and the telecommunications industry. Sample closures are telephone pedestals and transformer pads. Distribute granules uniformly

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Do II-HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Do-ItYourself Insect Control may be harmful if swallowed. Do not get an eyes or on slun.
Wash thoroughly after handling Wash contaminated clothing before re-use Flush
contaminated eyes with plenty of water and get prompt medical attention. Avoid
breathing vapors or dust particles. Avoid contacting food crops. Keep away from
food, feedstuffs and domestic water supplies. Keep container closed. Keep children
a of the stream of the supplies washed into the soil and grass is dry
NOTE TO PHYSICIAN. Chlorpyrifos is a cholinesterase inhibitor. Treat
symptomatically Atropine, only by injection, is an antidote.
ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and other
wildlife Birds and other wildlife in treated areas may be killed. Keep out of takes,
streams or ponds. Do not apply where runoff is likely to occur. Do not apply when
weather conditions favor drift from area treated. Do not contaminate water by
cleaning of equipment or disposal of wastes. Apply this product only as specified on
this label.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash

Manufactured by Do-It-Yourself Pest Control Utility, Inc. P. O. Box 26445, Birmingham, Alabama 35226 Telephone (205) 942-3328

16