

# AID GROOM & REPEL

A Concentrated Pyrethrin Micro Emulsion

•CLEAR •STABLE  
•LEAVES NO OILY OR OBJECTIONABLE RESIDUES

Effective, Fast Acting Insecticide

For Use in Stables, Barns, Dairies, and on Livestock

Apply with AID Horse & Stable Sprayer or Compressed Air Equipment

<b>ACTIVE INGREDIENTS:</b>	
Pyrethrins	0.2%
*Piperonyl Butoxide, Technical	2.0%
Petroleum Distillate	0.8%
<b>INERT INGREDIENTS</b>	<b>97.0%</b>
	<b>100.0%</b>

\*Equivalent to 0.8% (butylcarbityl (6-propylpiperonyl) ether and 0.2% related compounds

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See side panel for additional precautionary statements  
DO NOT TRANSPORT OR STORE UNDER 32 F

EPA Reg. No. 13130-B

EPA Est. No. 34797 IA-1 34145 NB-1,  
11715 TN-1 4371 FL-01

NET CONTENTS 1 GALLON (128 FL.OZ.) (3.785 L.)

**AID LABORATORIES, INC.**

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. Do not contaminate food and feed stuff. Do not apply where food processing takes place. After spraying in food processing plants, wash all equipment, benches, shelving, etc. where exposed food will be handled, with potable water. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water. Do not apply to aquatic life or to areas of surface water. Do not apply to areas of surface water. Do not apply to areas of surface water.

**STORAGE AND DISPOSAL**

Do not store in areas where food or feed is stored or disposed.

**PESTICIDE DISPOSAL**

Pesticide spray mixtures or residue that cannot be used according to label instructions must be disposed of in accordance with Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

**CONTAINER DISPOSAL**

Triple rinse for equipment. Then offer for recycling or reconditioning or puncture and dispose in an approved land fill or by other approved state and local procedures.

**DIRECTIONS FOR USE**

**DAIRIES:** The use of this product in food processing establishments should be confined to time periods when the premises are not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Vacate treated area and ventilate before reoccupying. For maximum effectiveness a combination of spot surface treatment and fogging is recommended.

**BARN AND STABLES:**

**SURFACE SPRAYING:** Use bagged sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored food, around the base of machinery, behind shelves and drawers. If surface application only is to be used, spray walls and other surfaces applying a rate of 1 gallon to 750 square feet of surface.

**SPACE SPRAYING:** Close doors and windows and shut off ventilating systems. Apply with a hand machine adjusted to deliver a fine mist or fog. Direct spray at an upward angle and distribute uniformly throughout the entire area at a rate of 3.6 ounces per 1000 cubic feet of space.

**FOGGING:** To kill exposed accessible stages of the stable fly and house fly, use a rate of 3.6 ounces per 1000 cubic feet of space. Control house flies directly whenever necessary.

**ON ANIMALS:** To protect cattle and horses from horn flies, house flies, mosquitoes and gnats, apply a light mist directly over the surface of the hair. To control stable flies, horse flies, and deer flies, apply a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatments at intervals necessary to give continued protection. For blood sucking lice, apply to the infested areas of the animal. A single application will get lice off the animal. Repeat every 2 or 3 weeks if required.

**AID LABORATORIES, INC.**  
Ocala, Florida U.S.A. 33472

1982  
13330-8