

SANADINE

for Sanitizing, Cleaning, Disinfecting

SANADINE... is a sanitizing, cleaning and disinfecting agent for poultry and livestock buildings and equipment, veterinary quarters and equipment; milk handling and processing equipment and machines.

SANADINE... is not adversely affected by low temperature or by water hardness.

SANADINE... Amber to brown color of use dilution is indicator of germicidal strength. When the initial color fades noticeably a fresh solution should be prepared.

See other pages for Use Directions

ACTIVE INGREDIENTS:	
...
...
INERT INGREDIENTS:	
...

DANGER: KEEP OUT OF REACH OF CHILDREN.



EPA Reg. No. 13680-17

NET CONTENTS:

manufactured by
OZARK CHEMICAL COMPANY
NORTH LITTLE ROCK, ARKANSAS 72113

ACCEPTED
13680-17
JUL 1 1975
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO.

SANADINE

Use-Directions for Poultry and Livestock Processing

For cleaning and disinfecting in one operation:

Use 4 oz. SANADINE per 5 gal. water (100 ppm titratable iodine)

1. Remove all poultry, livestock and feed from premises. Truck, coops, crates.
2. Remove all litter and droppings from floors, walls and surface of facilities occupied or traversed by poultry.
3. Apply solution to all surfaces to be sanitized.
4. Wash floors with water.

For sanitizing previously washed and sanitized, milk handling equipment and milk processing equipment. Use 1 oz. SANADINE per 5 gal. water (25 ppm titratable iodine). Allow minimum 10 minutes contact time before rinsing with potable water or allow to remain in equipment and rinse just before next use.

For disinfecting veterinary instruments, rubber gloves, tubing, etc. Use 1 oz. SANADINE per 1 gallon water (125 ppm titratable iodine) to soak for minimum of 10 minutes. Rinse thoroughly with potable water before use.

DANGER:

FIRST AID

CALL A PHYSICIAN.

RINSE EMPTY CONTAINER COMPLETELY WITH WATER BEFORE DISCARDING.