

# TRANSVAAL

## WEED-RHAP®

### LV OXY-6 D HERBICIDE

#### WHERE TO USE

WEED-RHAP LV OXY-6D is used to control broad-leaved weeds in cereal crops, corn, sorghum, weeds and brush in rangeland, pastures, similar noncrop uses, and for aquatic weed control.

#### PLANTS CONTROLLED

WEED-RHAP LV OXY-6D will kill or control the following as well as many other noxious plants susceptible to 2,4-D:

Alligatorweed	Chicory	Locoweed	Sagebrush
Arrowweed	Cocklebur	Mexicanweed	Shepherdspurse
Artichoke	Coffeebean	Morning-glory	Smartweed
Bindweed (hedge, field, and European)	Creeping Jenny	Mustard	Sowthistle
Bitter wintercress	Curly Indigo	Nutgrass	Stinkweed
Boxelder	Duckweed	Parsleyleaf	Sumac
Buckhorn	Elderberry	Pennywort	Sunflower
Bullthistle	Goldenrod	Plantain	Virginia creeper
Bulrush	Ground Ivy	Poison Ivy	Water hyacinth
Burdock	Hemp	Pokeweed	Water lily
Bur ragweed	Hoary cress	Povertyweed	Water primrose
Buttercup	Honeyuckle	Puncturevine	Wild garlic
Canada thistle	Indigo	Purslane	Wild lettuce
Catnip	Ironweed	Rush	Wild onion
Chickweed	Jimsonweed	Russian thistle	Wild radish
	Lambquarters		Willow

#### DIRECTIONS FOR USE

Add the recommended amount of product to about one-half the volume of water to be used in spraying. Mix well, then add remaining water and until spray mixture is uniform. Agitate thoroughly while spraying to a uniform emulsion. Best results are obtained when applications are made while the plants are well-leaved and actively growing.

**On Fall-Planted Wheat, Oats and Barley**—Use 1/3 to 1/2 pint of product in 5 to 10 gallons of water to cover one acre. Apply in early spring when weeds are small and before crop has reached the boot stage.

**On Spring-Planted Wheat, Oats and Barley**—Use 1/3 pint of product in 5 to 10 gallons of water to cover one acre. Apply after the fully tillered stage, except during the boot to dough stage.

**DO NOT** forage or graze treated grain fields within two weeks after treatment with 2,4-D. **DO NOT** feed treated straw to livestock.

**On Corn and Sorghum**—Use 1/3 pint of product in 5 to 10 gallons of water to cover one acre when weeds are in active growth. Local climatic conditions determine when treatment should be made. Best results are usually obtained when weeds are small and corn is 4 to 18 inches tall. **DO NOT** apply from tasseling to dough stage. Treat sorghum when 4 to 12 inches tall. Use drop nozzles when either crop is over 10 inches tall.

**For preemergence on Corn**—Use 1 1/3 to 2 2/3 pints of product in 5 to 10 gallons of water per acre. Use the higher rate on heavy soils. **DO NOT** use on sandy soils. Plant corn as deep as feasible.

**On Ornamental Turf**—Depending on type of weeds and stage of growth, use 2 1/2 to 2 pints of product and enough water to give sufficient coverage to one acre on established stands of perennial grasses. **DO NOT** use on creeping grasses such as bent except for spot spraying. Newly seeded turf should not be treated until after the second mowing, and the lower dosage rate should be used. Possible injury may occur to Dichondra and clover.

**On Rangeland, Pastures and General Weed Control**—Use 2 1/2 to 2 3/4 pints of product in sufficient water to give good coverage to one acre, depending on type of weeds and stage of growth. Use only on established stands of perennial grasses. **DO NOT** graze dairy animals on treated areas within seven days after application.

**For Aquatic Weeds in Lakes, Ponds and Drainage Ditchbanks**—Use 1 1/2 to 3 pints of product in 50 to 100 gallons of water and apply to one acre. Application should be made when leaves are fully developed above water line and plants are actively growing. Treatment of dense weed areas can result in oxygen loss from decomposition of dead weeds. This loss can cause fish suffocation. Therefore, to minimize this hazard treat 1/3 to 1/2 of the water area in a single operation and wait at least 14-14 days between treatments. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas.

Perennial and other Hard-To-Control Weeds may require a repeat application to give adequate control.

#### CAUTION

This product may cause skin irritation. Avoid inhaling spray mist. Harmful if swallowed. Avoid contact with the eyes, skin or clothing. Flush eyes with clear water and get prompt medical attention.

**DO NOT** apply this product to susceptible broad-leaved plants that are desirable such as cotton, soybeans, tomatoes, grapes, clover or ornamentals. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

Minute quantities may cause severe injury by drift. Avoid drift of spray and use a coarse spray which is less likely to drift. Although this product contains an ester of low volatility, use it with caution where the spreading of vapors can damage nearby susceptible plants.

**DO NOT** contaminate water in irrigation ditches or water used for domestic purposes.

**DO NOT** contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Users should note that herbicide treatment of public waters requires a permit from appropriate state agencies in most states. Your State Conservation Department or Game and Fish Commission will aid you in securing a permit if required in your state.

**DO NOT** store this product near fertilizers, seeds, insecticides or fungicides.

**DO NOT** contaminate feed or foodstuffs.

**DO NOT** use spray equipment contaminated with this product for any other purpose.

**DO NOT** apply insecticides, fungicides or other agricultural products unless the equipment has been cleaned very carefully with a suitable chemical cleaner.


**DO NOT** reuse containers. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

If stored below freezing, it may be necessary to warm to 40°F. and agitate before using. This does not affect the efficiency of the product.

**DO NOT** USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

#### WARRANTY

TRANSVAAL warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purpose. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.



## 2,4-D Low Volatile

6 LBS. OF 2,4-D ACID EQUIVALENT PER GALLON

<b>ACTIVE INGREDIENTS:</b>	
Butoxyethyl Ester of 2,4-Dichlorophenoxyacetic Acid*	88.08%
<b>INERT INGREDIENTS:</b>	11.92%
<b>TOTAL</b>	<b>100.0%</b>
*Equivalent to 60.61% or 6 pounds per gallon of 2,4-Dichlorophenoxyacetic Acid.	

MADE BY

### TRANSVAAL, INC.

JACKSONVILLE, ARKANSAS 72076

## ACCEPTED

NET CONTENTS: GAL.

Nov. 12, 1973

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 11687-62

EPA REGISTRATION NO. 11687-62

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**  
READ SIDE PANELS FOR ADDITIONAL HANDLING PRECAUTIONS.

# TRIFLURALIN TECHNICAL

Net Wgt. 110 lbs.

## ACTIVE INGREDIENT :

Trifluralin [alpha, alpha, alpha-trifluoro-2,6-dinitro-N,N-dipropyl-  
p-toluidine] . . . . .

97%

## INERT INGREDIENTS :

3%

Total 100%

(FOR MANUFACTURING USE ONLY)

## CAUTION: Keep out of reach of children.

Keep away from domestic animals and foodstuffs. Harmful if swallowed or absorbed through the skin. Avoid breathing of dust.

Use only with adequate ventilation. Avoid contact with eyes and skin. Do not smoke or eat while handling the product or before having washed with soap and water. In case of contact, flush with plenty of water.

Do not allow to freeze. Do not store near heat or flame. Do not reuse empty containers.

This product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate any body of water.

## ANTIDOTE:


If swallowed, give tablespoonful of salt with a glass of warm water to induce vomiting.

Toxicological class: Category 3

## DIRECTIONS FOR USE:

For use by commercial pesticide formulators.  
For reformulation or repackaging use only.

EPA Reg. No. 13801-12



**INIRACHEM SA**

Cendrier-Centre 17  
GENEVA

1201 GENÈVE  
Switzerland