

# PRAIRIE DOG, GROUND SQUIRREL, FIELD MICE, AND KANGAROO RAT BAIT

NET WEIGHT 50 LBS.

1973

Active Ingredient: . . . . .  
Inert Ingredient: . . . . .  
Total: . . . . .

132023

## WARNING KEEP OUT OF REACH OF CHILDREN

This product is toxic to wildlife, and birds. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**DIRECTIONS FOR USE:** For Prairie Dog: Scatter table spoonful quantities of bait near active mounds preferably on the side away from the pile of dirt which has been removed from the burrow. For Ground Squirrels: Scatter teaspoonful quantities of bait on surface of ground near burrow entrance. For Field Mice (Meadow Mouse and White-footed Mouse): Place one teaspoon of bait in surface trails or at the mouth of holes leading to underground burrow systems. Two to four bait placements should be made near the base of each infested tree. Cover bait by pulling surrounding grass back for access. Do not disturb the runway system. Baiting should be repeated at the rate of once per month. For Kangaroo Rat: Scatter one to two cups of bait around the perimeter of the mounds, by hand or by use of a large funnel.



# POISON



**ANTIDOTE:** CHEMICAL INDUCTION OF VOMITING. If less than ten minutes have passed since a person was taken, give a tablet of ipecac or a glass of warm water to induce vomiting. Treatment by this method after ten minutes may induce onset of convulsions. If vomiting does not occur, give 1-2 glasses of water. If there is no respiration, resuscitate by artificial respiration, mouth-to-mouth, and keep warm.

For information on the use of this product, contact the manufacturer or your local health department.

When placed out of the range of children and pets, and in areas where wildlife, not of value to you, are present, this product may be used.