

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

CAUTION

KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS AND PETS. MAY BE FATAL OR HARMFUL IF SWALLOWED. AVOID SKIN CONTACT. DO NOT BREATHE DUST. AVOID CONTAMINATION OF FEED OR FOOD STUFFS. AVOID CONTACT WITH ACIDS.

ENVIRONMENTAL HAZARDS

THIS PRODUCT IS TOXIC TO WILDLIFE AND FISH. BIRDS AND OTHER WILDLIFE FEEDING ON TREATED BAIT MAY BE KILLED. KEEP OUT OF ANY BODY OF WATER. DO NOT APPLY WHERE RUNOFF IS LIKELY TO OCCUR.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:—For control of prairie dogs [Black tailed (*Cynomys ludovicianus*), whitetailed (*C. leucurus*), and Gunnison (*C. gunnisoni*)] on rangeland in Western United States (North Dakota, South Dakota, Nebraska, Kansas, Oklahoma, Texas, New Mexico, Arizona, Colorado, Montana, Utah and Wyoming) where non-target species (especially blackfooted ferrets) will not be endangered.

Before undertaking prairie dog control with the product, consult with Local, State and Federal Game Authorities to insure that the area proposed for treatment is not inhabited by the blackfooted ferret, an endangered species and predator of the prairie dog.

PREBAITING — Prebait with 4 grams (one teaspoon) of untreated steam-rolled oats per mound, one to two days prior to baiting, to increase acceptance of treated baits by prairie dogs.

BAITING — Apply bait only after all or most of pre-bait is eaten and only to areas where untreated bait was consumed. Establish observation period during prebaiting. Apply bait by hand as a six-inch bait spot on edge of each mound or in adjacent feeding areas. Application rate should not exceed 4 grams (one teaspoon) per bait spot. Treat during late summer or fall period (July-December). Do not apply more than one bait application during this period. Dispose of spilled or unwanted bait by burial.

RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION

ZINC PHOSPHIDE ON STEAM-ROLLED OATS
FOR THE CONTROL OF PRAIRIE DOGS:
BLACK-TAILED (*CYNOMYS LUDOVICIANUS*)
WHITE-TAILED (*C. LEUCURUS*)
AND
GUNNISON (*C. GUNNISONI*)

Active Ingredient
Zinc Phosphide.....2.0%
by weight
Inert Ingredients.....98.0%
Total.....100.00%

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

If swallowed-Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger.

Avoid use of all oils. Have patient lie down and keep warm. Do not induce vomiting or give anything to an unconscious person. Get medical attention. If on skin or in eyes-Flush with plenty of water.

STORAGE AND DISPOSAL

PROHIBITIONS — Do not contaminate water, food or feed by storage or disposal. Open burning and dumping is prohibited. Do not reuse empty containers.

PESTICIDE DISPOSAL — Bait that cannot be used should be buried in a safe place away from water supplies in accordance with procedures approved by Federal, State, or Local Disposal Authorities.

CONTAINER DISPOSAL — Consult Federal, State, or Local Authorities for approved procedures.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than that indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

SD Dept. of Agric.
Rodent Control Fund
Anderson Building
Pierre, SD 57501
Est. No.1380-SD-1
EPA Registration No. 13808-6
Net Contents 25 lbs

REST DOCUMENT AVAILABLE

ACC. 12/28/60
OCT 27 1960
12/28/60

49