

ACCRA PAC Industrial Insecticide

A PYRENONE® Insecticide

Industrial Strength - Low Dosage Required

For use in Schools, Hospitals, Institutions, Hotels, Dairies, Supermarkets, Restaurants, Drive-ins, Motels, Drug Stores, Food Plants and Animal Quarters.

Provides Rapid Control of Houseflies, Mosquitoes, Small Flying Moths (Millers), Gnats, Wasps and accessible stages of Roaches, Ants, Water Bugs, Silverfish, Flour Beetles, Weevils, Grain Mites and Spiders.

ACTIVE INGREDIENTS:

Pyrethrins	0.40%
*Piperonyl Butoxide, Technical	2.00%
Petroleum Distillate	7.40%

INERT INGREDIENTS:	90.20%
	100.00%

*Equivalent to 1.6% (butylcarbityl) (6-propylpiperonyl) ether and 0.4% related compounds

PYRENONE - Registered trademark of Fairfield American Corporation

EPA Reg. No. 13891-24
EPA Est. No. 13891-IN-1

Manufactured By:

Accra Pac, Incorporated
2730 Middlebury St.
Elkhart, IN 46515

NET WT. 16 OZ.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements.

ACCEPTED

SEP 28 1983

U.S. Federal Insecticide
and Fungicide
Administration
EPA Reg. No. 13891-24

DIRECTIONS FOR USE

It is a violation of law to use this product in a manner inconsistent with the label directions.

SHAKE BEFORE AND AT INTERVALS DURING PROLONGED USE.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

Houseflies, Mosquitoes, Gnats, Wasps and Small Flying Moths: Close all doors and windows and spray room at the rate of 1 second per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat treatment as necessary.

Roaches, Water Bugs, Silverfish, Spiders, Crickets, Centipedes and Sowbugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Indian Meal Moths and Angoumois Grain Moths: Close doors and windows. Direct spray into all parts of room especially around stored food product containers, pallets and darkened areas.

Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needs.

Carpet Beetles: Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.

SEE REVERSE SIDE

8278L9

BEST DOCUMENT AVAILABLE

Commoned Flour Beetles, Saw-toothed Grain Beetles, Granary Weevils, Rice Weevils, Red Flour Beetles, Grain Mites: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary.

Lice: Spray on garments, bedding, mattresses, furniture, floors, carpeting and inanimate surfaces. First spray an inconspicuous area to test for possible staining or discoloration. Inspect after drying, then proceed to spray entire area to be treated.

Hold can upright with nozzle away from you. Press button and spray surface to be treated from a distance of 10 inches. Spray each square foot for 3 seconds, directing spray into all infested surfaces until moist.

For bedding: Make sure spray is directed into corners of fitted sheets and blankets as well as inside and outside of pillow cases. In the case of garments and headgear, close attention should be given to seams, hems, cuffs, collars, and creases. After treatment, bedding and all garments must be either laundered in hot water or dry cleaned as required.

For mattresses, furniture and carpeting: Direct spray into all folds, tucks, seams and crevices of infested areas until moist. Allow all treated surfaces to dry completely before use.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and frames. Repeat treatment as necessary.

Fleas: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

DISPOSAL

Do not reuse empty container.

Wrap container and put in trash collection.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, with potable water. All food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 120°F may cause bursting.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.