

ACTIVE INGREDIENT

2-chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate 3.00%

INERT INGREDIENT 97.00%

TOTAL 100.00%

PRECAUTIONS IN USING


CAUTION: Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and food stuffs. Keep away from heat and open flame, do not apply in dwellings.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

NOTICE OF WARRANTY

THE MANUFACTURER MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its use which extend beyond the use of the product under normal conditions in accord with the statements made on this label.



AGRI-BON

LIVESTOCK DUST

WITH RABON*

A ready-to-use Insecticide Dust to control HORN FLIES, LICE and aid in the control of FACE FLIES on BEEF and DAIRY CATTLE and LICE on SWINE.

*Shell Chemical Co. Trademark

CAUTION: Keep Out of Reach of Children.

See other cautions on side panel

NET WT. 10 LBS.

AGRICULTURAL ENTERPRISES, INC.

933 WEST 6th
FREMONT, NEB. 68025 U.S.A.

USE DIRECTIONS

The livestock dust is a ready-to-use insecticide which can be applied by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter.

SELF-TREATING DUST BAG:

FORCED USE: Put the livestock dust in a dispensing bag and hang in barn door exits or alleyways leading from animal buildings, salt or mineral blocks, or water holes. Protect cloth or burlap bags from weather.

FREE CHOICE USE: Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The free choice use aids in the control of lice.

HAND DUSTING: Apply approximately 2 oz. of dust by shake can, rotary duster or by spoon to the upper portions of the back, neck and poll, and to the face as an aid in the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary.

LICE ON SWINE: Apply 3-4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One pound of 3% dust should be applied per 150 sq. ft. of bedding.

TAKE TIME



OBSERVE LABEL DIRECTIONS

E.P.A. Reg. No. 14651-2
E.P.A. EST. 14651-NB 1

14651-2