RAT-A-TAC THRO PAC Rodenticide

ACTIVE INGREDIENT: Diphacinone 2-diphen laceteyl-1, 3-indandione	y- %
INERT INGREDIENTS 99.995	
TOTAL 100.000	

CAUTION:

KEEP OUT OF REACH OF CHILDREN ee back panel for additional cautions

NET WEIGHT 4 OZ.

Manufactured for AGRICULTURAL ENTERPRISES, INC.

933 W. 6th St. Fremont, Nebraska 68025 U.S.A.

DIRECTIONS:

RATS: Place 1 to 2 packets next to burrows in runways or where rats are accustomed to feeding. Check these bait stations daily to assure an uninterrupted supply of bait for a period of not less than 12 days. Continue baiting until all signs of feeding have stopped. One packet is effective for one or two rats. MICE: Place 1 ounce of bait in a shallow dry container next to burrows, in runways or where mice are accustomed to feeding. It is recommended that the bait be placed at intervals of 8 to 12 feet. Check bait stations daily to assure uninterrupted supply of bait for a period of not less than 15 days.

LONG TERM CONTROL: Bait continuously. Stale, damp or moldy bait should be replaced with fresh

bait promptly.

CAUTION: Keep Out Of Reach Of Children. Protect children, pets and domestic animals from the bait. Bait should be placed at locations not accessible to children, pets and domestic animals or in tamper-proof bait boxes. In case baits are eaten by humans, call physician at once. Treat as for overdose of Dicoumarol®.

CAUTION: This product is toxic to wildlife. Do not contaminate water by disposal of waste. Use only as specified on this label.

EPA Reg. No. 14651-8 EPA Est. No. 14651-NB-1