

11-19-74

THE FEDERAL BUREAU OF INVESTIGATION
U.S. DEPARTMENT OF JUSTICE
WASHINGTON, D.C. 20535

14651-1

FOR CONTROL OF HOUSE FLIES, FACE FLIES, AND STABLE FLIES IN AND AROUND LIVESTOCK BUILDINGS

Mix 2 quarts of Emulsifiable Insecticide (1.1 lb./gal) per 6 gallons of water (1% concentration). Spray inside and outside barn walls (including dairy barns), eaves, fences, garbage, manure, and other resting or breeding places at least twice monthly using approximately 1 gallon of finished spray per 1000 sq. ft. (40' x 25'). Do not use under, in or around poultry houses. Improved house fly and face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6 - 10 gallons of spray. Do not contaminate food, feed or drinking water.

WARNING

Keep out of reach of children. Do not contaminate feed or foodstuffs. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean rubber gloves and a mask or respirator of a type found suitable by the U.S. Bureau of Mines for protection against CIODRIN Insecticide. Use this material only for recommended purposes and at recommended dosages.

WARNING: HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN.

Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating and smoking, wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Keep away from heat and open flame. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Call a physician in all cases of suspected poisoning.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. DESTROY USED CONTAINERS. Do not reuse for any purpose. Decontaminate containers by rinsing thoroughly with water and alkaline detergent. Destroy containers by breaking and burying fragments in isolated area. Dispose of rinsings in a way as not to constitute a hazard, or contaminate water supplies.

NOTICE OF WARRANTY

THE MANUFACTURER MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

EPA Reg. No. 14651-9
EPA Est. No. 14651-NB-1



AGRI-CID

EMULSIFIABLE INSECTICIDE

(Contains 1.1 lb. CIODRIN®* Insecticide per Gallon)

FOR CONTROL OF STABLE FLIES, HORN FLIES, HOUSE FLIES, LICE AND LONE STAR TICKS ON DAIRY CATTLE, BEEF CATTLE, SHEEP, GOATS AND HOGS.

ACTIVE INGREDIENTS	BY WEIGHT
Dimethyl phosphate of alpha-methylbenzyl 3-hydroxy-cis-crotonate	11.60%
Xylene	75.49%
INERT INGREDIENTS	12.91%
	TOTAL 100.00%

*Shell Registered Trademark

WARNING: KEEP OUT OF REACH OF CHILDREN.

See other warning on side panel.

NET CONTENTS 1 GAL.

Agricultural Enterprises, Inc.

933 West 6th St.

Fremont, Nebraska 68025.

Emulsi
or low
water
to calv
show

FOR C
LONE :
Mix 2
tration
1 -- 2
peat a
7 days
For per
equipm
(1.1 lb
the an
of spr
tion as
not cor
Improv
the rat
to main

LOW V

Mix 4
water
the fac
week
per pir

FOR CC
AND B

Mix 8
of wat
cluding
tional
1 level
drinking

For use
1 quart
This mc

14651-9



AGRI-CID

EMULSIFIABLE INSECTICIDE

(Contains 1.1 lb. CIODRIN® Insecticide per Gallon)

FOR CONTROL OF STABLE FLIES, HORN FLIES, HOUSE FLIES, LICE AND LONE STAR TICKS ON DAIRY CATTLE, BEEF CATTLE, SHEEP, GOATS AND HOGS.

ACTIVE INGREDIENTS	BY WEIGHT
Dimethyl phosphate of alpha-methylbenzyl 3-hydroxy-cis-crotonate	11.60%
Xylene	75.49%
INERT INGREDIENTS	12.91%
	<hr/>
	TOTAL 100.00%

*Shell Registered Trademark

WARNING: KEEP OUT OF REACH OF CHILDREN.

See other warning on side panel.

NET CONTENTS 1 GAL.

Agricultural Enterprises, Inc.

933 West 6th St.

Fremont, Nebraska 68025

GENERAL DIRECTIONS

Emulsifiable Insecticide (1.1 lb/gal) is suitable for use in conventional hydraulic sprayers or low pressure knapsack sprayers. Prior to use, mix the required amount in sufficient water and apply in accordance with State and Federal Recommendations. Do not apply to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

DIRECTIONS FOR LIVESTOCK PEST CONTROL

FOR CONTROL OF STABLE FLIES, HORN FLIES, HOUSE FLIES, LICE, FACE FLIES AND LONE STAR TICKS ON DAIRY CATTLE, BEEF CATTLE, SHEEP, GOATS AND HOGS.

Mix 2 quarts of Emulsifiable Insecticide (1.1 lb/gal) per 6 gallons of water (1% concentration). Spray to thoroughly cover all parts of the animal, including the legs, using 1 - 2 pints of spray per large animal and proportionately less for smaller animals. Repeat applications as necessary to maintain control, but not more often than once every 7 days. Do not contaminate food, feed or drinking water.

For pen or corral operations involving spray applications to livestock with high pressure equipment, do not exceed a 0.5% concentration, mix 2 quarts of Emulsifiable Insecticide (1.1 lb/gal) per 12 gallons of water (0.5% concentration). Spray to thoroughly cover the animal's body and legs (avoid excessive spraying of the head) using 1 - 2 quarts of spray per large animal and proportionately less for smaller animals. Repeat application as necessary to maintain control, but not more often than once every 7 days. Do not contaminate food, feed or drinking water.

Improved face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6 - 10 gallons of spray. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

LOW VOLUME APPLICATION FOR CONTROL OF FACE FLIES

Mix 4 tablespoons (2 fluid oz.) of Emulsifiable Insecticide (1.1 lb/gal) per 1 1/2 pints of water (1% concentration). Using a calibrated sprayer, apply 5 teaspoons of spray to the face and a total of 5 - 10 teaspoons to the back and sides of the animal 3 times per week. Improved control may be achieved by the addition of 1 level teaspoon of sugar per pint of spray solutions. Do not contaminate food, feed or drinking water.

FOR CONTROL OF STABLE FLIES, HORN FLIES, HOUSE FLIES, AND FACE FLIES ON DAIRY AND BEEF CATTLE

Mix 8 tablespoons (4 fluid ounces) of Emulsifiable Insecticide (1.1 lb/gal) per 1 1/2 pints of water (2% concentration). Spray to thoroughly treat all parts of the animal, including the legs, using 1 - 2 fluid ounces of spray per animal. Apply daily with conventional hand or power sprayer. Improved control may be achieved by the addition of 1 level teaspoon of sugar per pint of spray solution. Do not contaminate food, feed or drinking water.

BACKRUBBER APPLICATIONS

For use in backrubbers or face rubbers for the control of horn flies and face flies, mix 1 quart of Emulsifiable Insecticide (1.1 lb/gal) per 4 gallons of back rubber base oil. This may be diesel fuel or fuel oil in accordance with local recommendations.