

GENERAL DIRECTIONS

This Insecticide is a specially prepared ready-to-use insecticide suitable for use in conventional hand, power or automatic spray equipment and in back rubbers. This product is approved for lactating dairy animals. No feedoff period is necessary when used on beef animals. Apply with hand power or automatic equipment to moisten the hair only and not in excess of 2 fluid ounces daily per animal. Do not wet the skin due to possible skin irritation. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

For the control of certain insects attacking animals apply in accordance with State and Federal Recommendations.

PRECAUTIONS IN USING

Harmful if swallowed. Avoid inhalation of mist and contact with skin. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention. Keep away from open flame. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. If swallowed, call physician immediately. Atrophine is antidotal.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

ACCEPTED



OCT 17

AGRI-TEAM

INSECTICIDE SPRAY

WITH CIODRIN® AND VAPONA®

READY-TO-USE

ANIMAL SPRAY SOLUTION

UNDER FEDERAL REGISTERED PATENT
 U.S. PATENT OFFICE
 PATENT NO. 2,851,710
 U.S. DEPT. OF AGRICULTURE

A Ciodrin® and Vapona® Insecticide Ready-to-Use Fly Spray For Dairy and Beef Cattle. Controls Horn Flies, Stable Flies, Face Flies, and Lice.

ACTIVE INGREDIENTS	% BY WEIGHT
Dimethyl Phosphate of Alpha-Methylbenzyl 3-Hydroxy-Cis-Crotonate*	1.00
2,2-Dichlorovinyl Dimethyl Phosphate	0.23**
Related Compound-	0.02**
INERT INGREDIENTS	98.75
	TOTAL 100.00

*CIODRIN® Insecticide

**Equivalent to 0.25% VAPONA® Insecticide

Shell Reg. Trade Mark

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on direction panel.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

NET 1 GAL

Agricultural Enterprises, Inc.

243 West 6th St.
Houston, Texas 77002

Lacta
the c
of th
the l

Lacta
secti
mal.
time

Back
beef
20 li
as n

Do n
Do n

THIS
POSE
tend
made


inven-
luct is
beef
y and
skin
cattle

and

f con-
Keep
before
l, call

iscard

ACCEPTED AGRI-TEAM



INSECTICIDE SPRAY

WITH CLODRIN* AND VAPONA*
READY-TO-USE
ANIMAL SPRAY SOLUTION

UNION PACIFIC
FURNISHES THE FOLLOWING INFORMATION
FOR THE USE OF THE CUSTOMER
EPA REG. NO. 14651-10
TO OBTAIN THIS PRODUCT

A CloDRIN® and Vapona® Insecticide Ready-To-Use Fly Spray For Dairy and Beef Cattle. Controls Horn Flies, Stable Flies, Face Flies, and Lice.

ACTIVE INGREDIENTS	% BY WEIGHT
Dimethyl Phosphate of Alpha-Methylbenzyl 3-Hydroxy-Cis-Crotonate*	1.00
2,2-Dichlorovinyl Dimethyl Phosphat	0.23**
Related Compounds	0.02**
INERT INGREDIENTS	98.75
TOTAL	100.00

*CLODRIN® Insecticide

**Equivalent to 0.25% VAPONA® Insecticide

®Shell Reg. Trade Mark

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on direction panel.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.
NET 1 GAL

Agricultural Enterprises, Inc.
933 West 6th St
Fremont, Nebraska 68025

USE AND DIRECTIONS FOR APPLICATION

Lactating Dairy and Beef Animals: Flies (Horn Fly, Stable Fly, Face Fly, House Fly). For the control of horn flies, stable flies, face flies, and house flies, make a light application of this insecticide daily as a fine mist to thoroughly cover all parts of the animal, including the legs. Do not exceed 2 fluid ounces per animal when applied as daily fly control.

Lactating Dairy and Beef Animals: Lice. Make 1 - 2 applications of 3 - 3 1/2 ounces of insecticide at 14 day intervals as a fine mist spray to thoroughly cover all parts of the animal. If material is applied by fogging or misting equipment, the maximum spraying time per animal is 20 seconds.

Back and Face Rubbers: For the control of horn flies and face flies on lactating dairy and beef animals, use insecticide in back rubbers and face rubbers. Use 1 gallon for each 20 linear feet of burlap back rubber. For automatic back rubbers recharge the reservoir as needed. Animals must rub face on rubbers for best fly control.

Do not treat animals within one day of slaughter

Do not apply dermally in combination with Trichlorofon.

CAUTION: KEEP OUT OF REACH OF CHILDREN

NOTICE OF WARRANTY

THIS COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

EPA Reg. No. 14651-10
EPA Est. No. 14651-NB-1