

AGRI-BON AQUA 50

A 50% WETTABLE POWDER

with Rabon®*

FOR USE AS A LIVESTOCK, POULTRY & PREMIS SPRAY

GENERAL DIRECTIONS

RABON® WP is suitable for use in conventional power or low pressure knapsack sprayers. Occasional agitation is recommended to prevent undue settling of the suspension. Follow the USE DIRECTIONS for the proper percent solution needed for a specific insect and areas or types of wall surfaces. Refer to DILUTION TABLE for the quantity of insecticide needed to make the percent solution recommended.

LIVESTOCK USE DIRECTIONS

Animal	Insect	Solution	Remarks
Beef Cattle	Horn Flies, Lice	0.35%	Apply as coarse spray. Use between 1/2 and 1 gallon of spray per animal depending on size and hair coat. There is no withholding period from last application to slaughter.
	Lone Star Ticks	0.5%	Apply as a coarse spray using 1 to 2 quarts per head to thoroughly wet the animal. Repeat in 2 weeks if necessary.
Swine	Lice	0.5%	Apply as a coarse spray using 1 to 2 quarts per head to thoroughly wet the animal. Repeat in 2 weeks if necessary.

POULTRY USE DIRECTIONS

Insect	Type of Housing	Solution	Remarks
Lice & Mites	Wire Cages	0.5%	Apply directly to birds. (1 gal 100 birds) spray vent and fluff areas from below. Repeat when necessary. Do not repeat more often than every 14 days. For individual bird treatment apply 1/2 ounce of the mixture per bird.

NOTE: For maximum lasting control of the northern fowl mite, penetration of the feathers around the vent area is absolutely essential. Use power sprayer at 100-125 psi at NO LESS THAN RECOMMENDED RATE. More attention must be given each individual bird when using low-pressure equipment. Treat roosters carefully and thoroughly to avoid reinfestation in breeding flocks.

Floor Litter	Management	0.5%	Apply 1/2 Gal 1000 sq. ft. evenly for penetration to litter surfaces. Also apply thoroughly to walls, roosts, cracks, crevices and interiors. Spray birds lightly.
Floor Litter	Management	50% WP	Treat evenly and thoroughly using 2.5 oz 100 sq. ft.
Floor Dust Box	Management	50% WP	Mix evenly throughout top layer of box contents using 2.5 oz 50 birds.
Floor Roost Point	Management	1.0%	Treat with brush or spray thoroughly particularly cracks and crevices using 1 pt 100 ft.
Fowl Tick (Blue Bug)	All Types	1.0%	Apply 1 gal 100-150 sq. ft. thoroughly to walls, ceiling, floor, cracks and crevices with a power sprayer.

ACTIVE INGREDIENTS

2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate

INERT INGREDIENTS

TOTAL

*A Shell Registered Trademark

BY WEIGHT

50.0%
50.0%
100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON SIDE PANEL.

TAKE TIME



OBSERVE LABEL DIRECTIONS

DILUTION TABLE


AMOUNT OF RABON® WP WETTABLE POWDER POUNDS	AMOUNT OF WATER FOR			
	0.35% SOLUTION GALLONS	0.5% SOLUTION GALLONS	1.0% SOLUTION GALLONS	2.0% SOLUTION GALLONS
1/2	9	6	3	1 1/2
2	36	25	12 1/2	6
4	75	50	25	12 1/2
8	150	100	50	25

NET CONTENTS 4 LBS

AGRICULTURAL ENTERPRISES, INC.

BOX O
FREMONT, NE, U.S.A. 68025

Insect	Problem Areas	Solution	Remarks
Flies	Dairy barns, poultry houses, swine barns, other animal buildings	2.0%	For dry whitewashed wood or concrete block surfaces 1 gal. of solution 500 sq. ft.
		1.0%	For unpainted wood or painted concrete block surfaces 1 gal. of solution 500 sq. ft.
		1.0%	For masonite or galvanized sheet metal surfaces 1/2 gal. of solution 500 sq. ft.
		1.0%	Apply 1 gal. of mixed solution 100 sq. ft. Penetrate problem area first time repeat every 7-10 days thereafter.
		0.5%	Spray infested areas thoroughly approximately 25 gals. of mixed solution acre. Spray along foot paths and roadsides leading to such areas.



Under the Federal Insecticide, Fungicide, and Rodenticide Act registered under EPA Reg. No. 401-11

Precautions in Using

CAUTION: Harmful if swallowed. Avoid breathing spray mist and dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Burn or discard in a safe place.

NOTICE OF WARRANTY

THE MANUFACTURER MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses, which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

EPA Reg. No. 14651-11

EPA Est. No. 1465-NB-1