


RECEIVED
 JUN 2 1984
 14651-14



FLY DIE

Sugar Bait

with Vapona*

Use Wet or Dry for Fast, Effective Fly Control

ACTIVE INGREDIENTS:

2,2-Dichlorovinyl dimethyl phosphate 0.46%
 Related Compounds 0.04%

INERT INGREDIENTS 99.50%
TOTAL 100.00

* A Shell Registered Trademark

CAUTION:

KEEP OUT OF REACH OF CHILDREN

See other cautions on instruction sheet.

NET WT.: 1 LB.

AGRICULTURAL ENTERPRISES, INC.

933 West 6th St.
 Fremont, Nebraska 68025

USE AND DIRECTIONS FOR APPLICATION

This Insecticide is a specially prepared insecticidal bait for use in the control of flies and can be used wet or dry.

LIQUID BAIT: Dissolve 1 lb. of bait in 1 gallon of water and apply as a coarse spray, sprinkle or spray.

DRY BAIT: Scatter the bait where flies congregate.

FOR REDUCING POPULATIONS OF RESISTANT AND NON-RESISTANT HOUSEFLIES IN AND AROUND OUTDOOR LIVING AREAS, PICNIC GROUNDS, BACKYARD AREAS, PATIOS, LOADING DOCKS, OUTDOOR LATRINES, PARKING AREAS, REFUSE AREAS AROUND SERVICE STATIONS, OPEN-AIR DRIVE-INS, ICE CREAM STANDS, GARBAGE COLLECTION AND DISPOSAL AREAS: Scatter the dry bait or apply the liquid bait to areas where flies congregate. For open-air drive-ins, ice cream stands and restaurants, apply on apron between serving counter and parking area, paying particular attention to trash barrels and garbage cans where flies congregate. Apply 1/2 lb. dry bait or 2 qt. liquid bait per 1,000 sq. ft.

DAIRY BARN, MILK SHEDS, STABLES, LIVESTOCK BARN, LOAFING SHEDS, PIG PARLORS AND OUTDOOR AREAS: Scatter the bait or apply the liquid bait to floors or ground areas where flies congregate. Ideal baiting sites are around doorways, feed storage rooms, alleyways and window sills. On freshly limed floors, place treatment on waste cloth or on paper. Repeat as necessary for practical fly control. Apply 1/4 lb. dry bait or 1 qt. liquid bait per 1,000 sq. ft.

CAGED POULTRY AND TURKEYS ON WIRE: Scatter the dry bait or apply the liquid bait on droppings beneath the cages or perches. Repeat at weekly intervals. Apply 1/4 lb. dry bait or 1 qt. liquid bait per 1,000 sq. ft.

POULTRY HOUSES: Scatter the dry bait or apply the liquid bait to areas where flies congregate but keep out of reach of birds. Apply to manure on window sill, outside of penned enclosures and on floor of feed storage room. Repeat as necessary. Apply 1/4 lb. dry bait or 1 qt. liquid bait per 1,000 sq. ft.

DOG KENNELS: Scatter the dry bait or apply the liquid bait outside dog runways on window sills, ledges and other surfaces out of reach of animals. For caged units apply to dropping area. Repeat as necessary. Apply 1/4 lb. dry bait or 1 qt. liquid bait per 1,000 sq. ft.

GARBAGE CANS: Apply the dry bait around and inside garbage cans. Repeat as necessary.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be used in conjunction with atropine.

NOTICE OF WARRANTY

The Manufacturer makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its uses, which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

E.P.A. Reg. No. 14651-14

E.P.A. EST. No. 14651-NB-1