

BEEF & DAIRY CATTLE

Horn Flies, Lice, Face Flies Place contents of this package in any commercially available dust bag. Suspend bags in areas frequented by cattle or in gateways or lanes through which the animals pass daily for water, feed or minerals. Bags may also be suspended in loafing sheds or in front of protected mineral feeders. For lactating dairy cows, bags may be suspended in the exit through which cows leave the milking barn. In all cases the bags should be adjusted so that the bottom of the bag will hang four to six inches below the topline of the cattle. For reduction of face flies, the bags must be located so the animals will be forced to use them daily and hung at a height to insure that the faces of the cattle will be dusted.

Horn Flies, Lice—Direct Application Apply 2.0 ounces of dust per animal by shaker can over the head, neck, shoulders, back and tailhead. Repeat as necessary.

SWINE

Lice on Swine—Direct Application Apply not more than 1 oz (3 level tablespoonfuls) per head as a uniform coat to the head, shoulders and back by use of a shaker can or suitable mechanical dust applicator. Repeat as necessary but not more often than once every ten days.

Bedding Treatment Appl. 2 oz (6 level tablespoonfuls) uniformly over each 30 square feet of fresh, dry bedding by use of a shaker can or a suitable mechanical dust applicator. Repeat as necessary but not more often than once every ten days.

In severe infestations both individual animals and the bedding may be treated as directed above.

No interval is required between treatment and slaughter or between treatment and use of milk as food.

*NOTE: 1 oz of this material equals approximately three level tablespoonfuls.

oda:

BEST DOCUMENT AVAILABLE



AGRI-DUST

ANIMAL INSECTICIDE

with Co-Ral*

For Control of Horn Flies & Lice on Beef & Dairy Cattle. As an Aid in Reducing Face Fly Population on Beef and Dairy Cattle. Control of Lice on Swine.

Use in Dust Bag, Shaker Can or Mechanical Dust Applicator.

ACTIVE INGREDIENTS

O, O Diethyl O 3-chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl phosphorothioate 10%

INERT INGREDIENTS 99.0%

100.0%

*A Reg. TM of the parent company of Farbwerke Bayer, GmH, Leverkusen.

CAUTION
KEEP OUT OF REACH OF CHILDREN
See side panel for warning statements.

NET WT. 10 LBS.

Agricultural Enterprises, Inc.

PO Box 0
Fremont, Nebraska 68025

RESTRICTIONS

For external insecticidal use on specified animals only. Avoid contamination of feed troughs, water and water utensils. Provide thorough ventilation while dusting. Persons treating animals should avoid repeated direct contact with skin or inhalation of dust.

CAUTION

Harmful if inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling.

Do not contaminate feed or food. Keep out of reach of children.

If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.


To Physicians—Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

WARRANTY

THE MANUFACTURER MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE OR OTHERWISE EXPRESS OR IMPLIED concerning this product or its use which extend beyond the use of the product under normal conditions in accord



Reg. No. 14651-15
EPA Est. No. 14651-98-1

BEST DOCUMENT AVAILABLE

BEEF & DAIRY CATTLE

Horn Flies, Lice, Face Flies Place contents of this package in any commercially available dust bag. Suspend bags in areas frequented by cattle or in gateways or lanes through which the animals pass daily for water, feed or minerals. Bags may also be suspended in loafing sheds or in front of protected mineral feeders. For lactating dairy cows, bags may be suspended in the exit through which cows leave the milking barn. In all cases the bags should be adjusted so that the bottom of the bag will hang four to six inches below the topline of the cattle. For reduction of face flies, the bags must be located so the animals will be forced to use them daily and hung at a height to insure that the faces of the cattle will be dusted.

Horn Flies, Lice—Direct Application Apply 2.0 ounces of dust per animal by shaker can over the head, neck, shoulders, back and tailhead. Repeat as necessary.

SWINE

Lice on Swine—Direct Application Apply not more than 1 oz. * (3 level tablespoonfuls) per head as a uniform coat to the head, shoulders, and back by use of a shaker can or suitable mechanical dust applicator. Repeat as necessary but not more often than once every ten days.


Bedding Treatment Apply 2 oz. * (6 level tablespoonfuls) uniformly over each 30 square feet of fresh, dry bedding by use of a shaker can or a suitable mechanical dust applicator. Repeat as necessary but not more often than once every ten days.

In severe infestations both individual animals and the bedding may be treated as directed above.

No interval is required between treatment and slaughter or between treatment and use of milk as food.

*NOTE: 1 oz. of this material equals approximately three level tablespoonfuls.

ada:



For Control of Horn Flies
Cattle—As an Aid in Reducing
Beef and Dairy Cattle—Control

Use in Dust Bag, Shaker
Applicator

ACTIVE INGREDIENTS:

O, O-Diethyl O-(3-chloro-4-
-oxo-(2H)-1-benzopyran-7-
phosphorothioate.

INERT INGREDIENTS:

®A Reg. TM of the
Bayerwerke Bayer GmbH,

CAUTION
KEEP OUT OF REACH OF CHILDREN
See side panel for

NET WT

Agricultural Formula

PO Box
Fremont, Nebraska

RESTRICTIONS

For external, insecticidal use on specified animals only. Avoid contamination of feed, troughs, water and water utensils. Provide thorough ventilation while dusting. Persons treating animals should avoid repeated direct contact with skin or inhalation of dust.

CAUTION

Harmful if inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling.

Do not contaminate feed or food. Keep out of reach of children.

If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

To Physicians—Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

WARRANTY

THE MANUFACTURER MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its use which extend beyond the use of the product under normal conditions in accord

ACCEPTED

MAY 8 1975

Reg. No. 14681-18
EPA Reg. No. 14681-18B-1

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the purpose of

10