

SEVIN DUST

EPA REG. NO. 14782-2

NET WT. 50 LBS.


DIRECTIONS FOR USE

① **APPLES, PEARS:** Codling Moth, Green Apple Aphid, Red-banded Leaf Roller, Pear Psylla, Apple Maggot, Periodical Cicada — To avoid possible fruit thinning do not apply until 30 days after full bloom. Repeat at 10 day intervals if needed. Do not apply within 1 day of harvest. Use 70-120 pounds per acre.

② **CORN:** Corn, Earworm, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetles — Use 40 pounds per acre. For larvae in whorl and foliage feeders, apply to entire plant. Apply to silks as they appear. Make 3 - 4 applications at 2 - 3 day intervals as infestation warrants. Re-apply immediately after rain. For European corn borer—consult your state agricultural experiment station for proper timing of treatments.

Application to the tassel region of sweet corn during the pollen shed period will seriously reduce bee population.

③ **COTTON:** Boll Weevil, Bollworm, Cotton Fleahopper, Lygus Bugs, Thrips, Pink Bollworm — Use 30 to 50 pounds of dust per acre. Apply when insects first threaten damage to crop, and repeat at 5 to 7 day intervals as required to maintain control.



Manufactured by
CARUSO FARM SUPPLY CO., INC.
120 N. WHITE HORSE PIKE
HAMMONTON, NEW JERSEY 08037

ACTIVE INGREDIENT:

*Carbaryl (1-Naphthyl N-methylcarbamate) 5.0%

INERT INGREDIENTS: 95.0%

100.0%

*Sevin, U. S. Pat. Nos. 2,903,478; 3,009,855

CAUTION

Keep Out of Reach of Children

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes. Do not burn. Do not reuse container.

DIRECTIONS FOR USE (CONT.)

④ **SORGHUM (GRAIN AND FORAGE):** Armyworms, Webworms, Corn Earworm, Stink Bugs, Grasshoppers, Sorghum Midge - Use 30 to 40 pounds per acre. For optimum control on grain sorghum direct dust into forming heads. For sorghum midge control, make applications 3 to 4 days after heads have emerged from boot. Do not apply within 21 days of harvest of grain. May be applied to day of harvest for sorghum forage.

⑤ **TOBACCO:** Hornworms, Budworms, Tobacco Flea Beetles, Grasshoppers, Japanese Beetles - Use at the rate of 30 to 50 pounds of dust per acre in broadcast treatment. Full coverage is essential. Begin treatments when worms are small. Repeat as needed. Use the highest dosage on large plants. If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered immediately after dust has settled. Do not apply within 3 days of priming or cutting.

*Trade-mark of Union Carbide Corporation for the active ingredient carbaryl.