

ACCEPTED

AS

AS



WARNING

KEEP AWAY from FEED or FOOD PRODUCTS

POISON

CAUTION - DO NOT DROP

IF LEAKING DON'T

**BREATHE FUMES
TOUCH CONTENTS
SWALLOW**

This is to certify that the contents of this package are properly described by name and are packed and marked and are in proper condition for transportation according to the Regulations prescribed by the Department of Transportation

**Apopka Growers Supply, Inc.
Apopka, Florida**

Shipper's name required hereon for shipments by EXPRESS

ONE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
ECONOMIC POISON REGISTERED
NO. 15091-3

5-13-76

A Gas
Special
gri.

MBS-68

FORMULATED and PACKAGED

by

APOPKA GROWERS SUPPLY, INC.

P.O. Drawer 1147
Apopka, Florida 32703

US Highway 1 South
Stuart, Florida 33494

ACTIVE INGREDIENTS	70%
Methyl Bromide	68.6%
Chloropicrin	1.4%
INERT INGREDIENTS	30%

DANGER

Keep Out of Reach
of Children



POISON



See other warnings
on side panel

DO NOT INHALE VAPORS

ANTIDOTE: Remove victim to fresh air immediately. Keep victim lying down and warm. Give artificial respiration if breathing has stopped. **CALL A PHYSICIAN IMMEDIATELY!**

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.

NET WEIGHT _____ LBS.

MADE IN U.S.A.

EPA Reg. No. 15091-3

PRINTED IN U.S.A.

MBS 68

DIRECTIONS: Use MBS 68 to control as cyst formers, meadow (lesion), root and spiral. Also for the control of white worms. Apply when the soil is in a good condition.

For preplant treatment of:

- Seed or plant beds for growing transplants of cauliflower, eggplants, muskmelons, apples, strawberries and tomatoes—2.9 lbs. of MBS 68 per 100 sq. feet. Expose 48 hours, then aerate 4 to 10 days before planting.
- Field soil treatment where tomatoes are to be grown—dosage 350 lbs. of MBS 68 per acre. Expose 48 hours. Aerate 2 weeks before transplanting in treated areas.
- For preplant treatment of soils for such as ornamental nursery beds, lawns, etc.

Use MBS 68 at the rate of 250 to 500 lbs. per acre. Apply by means of chisels spaced no more than 18 inches apart set at a depth of 4 to 6 inches. Use a higher rate in heavy soils, soils with high organic matter or when soil temperatures are low. Cover with ethylene or other gastight sheeting as soon as possible after chisel application. Leave cover on for 48 hours. If soil temperature is 60° F. or higher. Expose soil temperature is between 50° F. and 60° F.

PRECAUTIONS: Aerate treated soil before planting seed or 7 to 10 days before planting vegetative growth such as cuttings, slips, etc. Under some conditions, growth difficulties have been experienced on carnations, holly, snapdragons, roses, and certain other plants. If there is a problem on a small spot to determine safety.

Apply fertilizers containing nitrate or sulfate after the crop is established and soil temperatures are above 65° F. to avoid ammonia nitrogen starvation.

To prevent injury to desirable living plants, cover with at least 18 inches from drip line areas containing living roots of these plants.

Fumigated areas may be protected against recontamination by wooden frames or trenches to be placed around or washed. Recontamination by foot traffic should be kept to a minimum.

ACCEPTED

5-13-76

ONE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
UNION REGISTERED
NO. 15091-3



MBS-68

FORMULATED and PACKAGED
by

APOPKA GROWERS SUPPLY, INC.

P.O. Drawer 1147
Apopka, Florida 32703

US Highway 1 South
Stuart, Florida 33494

ACTIVE INGREDIENTS	70%
Methyl Bromide	68.6%
Chloropicrin	1.4%
INERT INGREDIENTS	30%

DANGER

Keep Out of Reach
of Children



POISON



See other warnings
on side panel

DO NOT INHALE VAPORS

ANTIDOTE: Remove victim to fresh air immediately. Keep victim lying down and warm. Give artificial respiration if breathing has stopped. **CALL A PHYSICIAN IMMEDIATELY!**

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.

NET WEIGHT _____ LBS.

MADE IN U.S.A.

EPA Reg. No. 15091-3

PRINTED IN U.S.A.

MBS 68

DIRECTIONS: Use MBS 68 to control nematodes such as cyst formers, meadow (lesion), root knot, lance sting and spiral. Also for the control of white grubs and wireworms. Apply when the soil is in a good friable, moist condition.

For preplant treatment of:

- Seed or plant beds for growing transplants of broccoli, cauliflower, eggplants, muskmelons, peppers, pineapples, strawberries and tomatoes—doses of 1.45 to 2.9 lbs. of MBS 68 per 100 sq. feet. Expose 24 to 48 hours, then aerate 4 to 10 days before seeding or planting.
- Field soil treatment where tomatoes or strawberries are to be grown—dosage 350 lbs. of MBS 68 per acre. Expose 48 hours. Aerate 2 weeks before setting transplants in treated areas.
- For preplant treatment of soils for non-food crops such as ornamental nursery beds, floral crop seedbeds, etc.

Use MBS 68 at the rate of 250 to 500 lbs. per acre. Apply by means of chisels spaced not more than 12 inches apart set at a depth of 4 to 6 inches. Use at the higher rate in heavy soils, soils with high organic content, or when soil temperatures are low. Cover with polyethylene or other gastight sheeting as soon as possible after chisel application. Leave cover on for 24 hours if soil temperature is 60° F. or higher. Expose for 48 hours if soil temperature is between 50° F. and 60° F.

PRECAUTIONS: Aerate treated soil for 48 to 72 hours before planting seed or 7 to 10 days before putting out vegetative growth such as cuttings, slips, sets or spuds. Under some conditions, growth difficulty has been experienced on carnations, holly, snapdragons, multiflora roses, and certain other plants. If there is a question, use on a small spot to determine safety.

Apply fertilizers containing nitrates or ammonium sulfate after the crop is established and the soil temperatures are above 65° F. to avoid ammonia injury or nitrogen starvation.

To prevent injury to desirable living plants, keep edge of cover at least 18 inches from drip line. Do not apply to areas containing living roots of these plants.

Fumigated areas may be protected against recontamination by wooden frames or trenches to prevent flooding or washing. Recontamination by footwear or equipment should be kept to a minimum.



POISON



ANTIDOTE

- Remove victim to fresh air immediately.
- Keep victim lying down and warm.
- Give artificial respiration if breathing has stopped.
- Call a physician immediately.

DANGER

WARNING: Poisonous liquid and vapor

- Vapors can be fatal if inhaled.
- Contact can cause burns.
- Do not get in eyes, on skin, or on clothing.
- In case of contact, wash skin thoroughly with soap and water. Flush eyes with water for at least 15 minutes. Get medical attention at once. Remove contaminated clothes, including shoes, immediately, and thoroughly aerate before re use.
- For skin irritation use boracic lotion or skin ointment.
- Get medical attention.
- Be sure container is closed completely.
- Store in a cool, well ventilated place.
- This product is toxic to fish. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

PERSONAL PROTECTION: When necessary to enter an atmosphere of MBS 68 use full face gas mask with BLACK Canister designed for acid gases and organic vapors.

- Empty containers should be returned to Apopka Growers Supply, Inc., Stuart, Florida via freight, collect.