

USE DIRECTIONS

Use of 2 ounces per gallon of this product will be effective against Pseudomonas aeruginosa (PRD) at this level. This has been substantiated by the ACGI Use Duration Confirmation Test. At this level, the product is reported to age best pathogen flora.

For best results, use a detergent with a detergent applicator to walls, floors and other hard surfaces.

If used for sanitization of food equipment, food contact items, and linens, use at the level indicated on the product label.

NO RINSE REQUIRED

For heavy soiling, a longer contact time is recommended.

PHENOL FREE FORMULA

GUARANTEED	Minimum
Alkalinity	42
Surfactant	27

For complete information, see the label on the product.

SANIDET

**CLEANER
DEODORIZER**

**DISINFECTANT
FUNGICIDE**

DESIGNED FOR HOSPITAL AND INSTITUTIONAL USE

ACTIVE INGREDIENTS

- Alkyl 160 C-30 1.00%
 - dimethyl benzyl ammonium chloride 1.00%
 - Alkyl 16R C-32 1.00%
 - ethylbenzyl ammonium chloride 1.00%
 - Sodium Carbonate 1.00%
 - Tetrasodium ethylene diamine tetraacetate 1.00%
- INERT INGREDIENTS** 91.50%


NET CONTENTS: 1 GALLON (128 FL. OZ.)

WARNING

Keep Out of Reach of Children. See right side panel for additional precautionary and first aid statements.

Wausau Chemical Corporation
Wausau, Wisconsin 54401

ACCEPTED
SEP 29 1978
UNDER THE FEDERAL FOOD, DRUG & COSMETIC ACT
FUNGICIDE AND ROSENGLIDE ACT
FOR EXERCISE OF PUBLIC HEALTH
ED UNDER NO. 15269



DANGER

Keep Out of Reach of Children. Avoid contact with eyes, nose, mouth, and skin. If contact occurs, wash thoroughly with water. Avoid contact with food.

FIRST AID

In case of contact, see label for first aid instructions. Wash with plenty of water for at least 15 minutes. If in eyes, call a physician. Remove and wash clothing before reuse. If swallowed, induce vomiting. If white, gelatinous material is present, do not induce vomiting. Drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN

Probable bacterial infection. Use of gastric lavage. Measure respiratory rate. Shock respiratory depression. First aid may be needed.