

ROLAR BRAND

40

Liquid Sodium Hypochlorite

DANGER

Keep Out of Reach of Children

DANGER: Harmful if swallowed. Skin and eye contact may cause severe irritation. Avoid contact with skin and eyes. Do not mix with other products or chemicals (such as household ammonia, detergents, or acids). To do so will release hazardous gases.

FIRST AID: EXTERNAL: If on skin, flush skin with plenty of soap and water. If in eyes, flush with water for at least 15 min. Get prompt medical attention.

INTERNAL: If swallowed, drink large quantities of milk, egg whites, or a starch solution or, if these are not available, drink large quantities of water. Do not give vinegar or other acids. Do not induce vomiting. Get prompt medical attention.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL: Triple rinse container (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill or other safe place.

ACCEPTED
15265-11
DEC 15 1977
UNDER THE FEDERAL INSECTICIDE ACT
EPA REG. NO. 15265-11
EPA EST. NO. 15265-WI-1
REGISTERED
10

ACTIVE INGREDIENT
Sodium Hypochlorite 10.50%
INERT INGREDIENT 89.50%
NET CONTENTS 100.00%

WAUSAU CHEMICAL CORP.
Wausau, Wisconsin

NET CONTENTS

DAIRY USE DIRECTIONS

DAIRY FARMS:

(1) Clean all milking utensils and equipment in the normal manner. Just before use, sanitize all surfaces with a 200 ppm available chlorine dilution of Rolar 40. Allow 2 min. exposure time. Always follow with a thorough rinsing operation.
(2) Pipelines: Flush clean and rinse immediately after use. Just before next milking sanitize with a dilution of Rolar 40 providing 200 ppm available chlorine. Allow 2 min. exposure time.

(3) Bulk Tanks: Flush clean and rinse when empty. Sanitize with a dilution of Rolar 40 providing 200 ppm available chlorine just before next use. Allow 2 min. exposure time.

DAIRY AND FOOD PLANTS:

(1) Clean equipment in the normal manner. Just before use, sanitize all surfaces with a dilution of Rolar 40 providing 200 ppm available chlorine. Allow at least 2 min. exposure time, and allow to drain. Do not rinse with water after this treatment. Do not soak overnight.

(2) Prepare a 200 ppm available chlorine solution (add 1.0 Rolar 40 to 5 gallons of water).

(3) The product provided with this label is for use in dairy and food plant applications. It is not intended for use in swimming pools.

TABLE OF PROPORTIONS

100 ppm use 1.0 to 2.0 gal. water
200 ppm use 1.0 to 5 gal. water
500 ppm use 1.0 to 1 gal. water