

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

NOV 10 ZDIO

Mr. Joe Weix Product Registration Wausau Chemical Corporation 2001 N. River Drive 54403/ PO Box 953 Wausau, WI 54402-0953

SUBJECT:

Application for Pesticide Notification (PRN 98-10)

Request General Label Change / Comply with State of California Regulations

EPA Reg. No.15265-19

Application Dated October 26, 2010

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 10/26/10 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-5335 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely.

Paul J. Mastradone, Ph.D., Acting

Notifications & Minor Formulations Team Leader

Registration Division (7505P) Office of Pesticide Programs

Please read instructions on n	everse before completing form.			Form Appro	oved.	UMB No. 2070-	0060	Print Form
SEPA	United States Environmental Protection Agency Washington, DC 20460				×	Registration Amendment Other		OPP Identifier Number
Application for Pesticide - Section I								
Company/Product Number Wausau Chemical Corpor		. [2		roduct Mana			3. Pro	posed Classification
4. Company/Product (Name) Hay Green		F	PM# 20. None Restricted					
5. Name and Address of App Wausau Chemical Corpo 2001 N River Drive 54403 Wausau, WI 54402-0953	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name							
			on - I					
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below. X Final printed labels in response to 04/19/2010 Agency letter dated "Me Too" Application. Other - Explain below. Other - Explain below.					FICATION			
Explanation: Use additional page(s) if necessary. (For section I and Section II.) Need to revise and update label per EPA letter (04/19/2010) and for state submission of our label to the state of California.								
		Section	on - II	1		, <u>, , , , , , , , , , , , , , , , , , </u>		
1. Material This Product Will	Be Packaged In:							
Child-Resistant Packaging Yes* X No * Cartification must be submitted	Yes Yes X No No Part No Par		Water Soluble Packaging Yes X No If "Yes" No. per Package wgt container		 .	2. Type of Container Metal X Plastic Glass Paper Other (Specify)		pacify)
3. Location of Net Contents	Information 4. Size(s) Reportainer Bulk 2400#,		r		5. Lo	cation of Label D On Label On Labeling		ns panying product
6. Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled								
		Section						000000
	items directly below for identificati	Υ	uél to be	contacted,	if nec			
Name Joe Weix	Title Analytical Chemist Telepho				e No. (Include Area Code)			
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and completion Received Backnowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment crees both under applicable law. (Stamped)								
2. Signature	3. Title Analytical	nalytical Chemist				() () () () () () () () () ()		
4. Typed Name		5. Date						, ι ε ι
Joe Weix		Oct 26, 20	010					

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Hay Green is a buffered propionic acid for use as a preservative in processing high moisture hay, corn, grain silage, (barley, corn, oat, sorghum, and wheat), haylage, and feed blends. It can be used post-harvest on alfalfa, barley grain, Bermuda grass, bluegrass, brome grass, clover, corn grain, cowpea hay, fescue, lespedeza, lupines, oat grain, orchard grass, peanut hay, peavine hay, rye grass, sorghum grain, soybean hay, sudan grass, timothy, vetch and wheat grain.

This product may only be used during the baling process if appropriate baler equipment is used which prevents exposure of non-target organisms to this pesticide product. Do not apply to pre-harvest hay and straw, cut hay and straw in the field prior to baling, or to hay and straw after it has been baled. Apply product to the hay and straw as it is fed into the baler, using the rates on this label. The baler application equipment must prevent drippage and drift, and must function properly and be used/maintained in accordance with the manufacturer's written operation instructions.

FOR USE ON HAY: Spray Hay Green on the hay as it is fed into the bailer using the application rate indicated by the chart below:

Hay	Small Square/	Large Round	Bales Large So	quare Bales
Moisture	Stem Moisture	Dew Moisture	Stem Moisture	Dew Moisture
16%-22%	4 lb/ton	2 lb/ton	6 lb/ton	3 lb/ton
22%-26%	8 lb/ton	6 lb/ton	10 lb/ton	8 lb/ton
27%-30%	16 lb/ton	12 lb/ton	Do not bale	16 lb/ton
Over 30%		Not recomme	nded	

- 1. Avoid baling if the windrow moisture level exceeds 30 percent.
- Always check the hay moisture level before baling high moisture hay.
 For rained-on hay or poorly ventilated storage, increase the application
- rate by 4-5 lb/ton.

 4. Make sure that the spray nozzle's openings and placement are such that
- Make sure that the spray nozzle's openings and placement are such tha good coverage is obtained.
- 5. Make sure that the storage area for the baled hay is well ventilated.
- 6. Do not store untreated dry hay on top of treated wet hay.

*NOT FOR USE ON SILAGE, HAYLAGE, FEEDBLEND AND CORN IN CALIFORNIA. FOR USE ON SILAGE, HAYLAGE, and FEED BLENDS: Spray Hay Green on crop at chopper intake or blower intake using the application rate indicated by the chart below:

CROP	MOISTURE	RATE
Corn silage	60-70%	2-3 lb/ton
Small grain silage	60-70%	2.5-3.5 lb/ton
Haylages	50-70%	2.5-3.5 lb/ton
Feed Blend (short term storage)		2-4 lb/ton -increase
•		for greater stability

FOR USE ON CORN: Spray Hay Green on ground or cracked corn using the application rate indicated by the chart below. Must double or triple for whole corn.

GRAIN	MOISTURE	RATE
Ear Corn	28ન્3ે4% ્રેિ	1്5 lb/ton
Shelled Corn	24-39% [ം പ്രദേ lb/ton

- 1. Make sure the silo does not have any cracks or holes that will allow air leakage
- 2. Harvest at full dent and at the proper moisture level:

 Ground ear-coin 28-24% moisture

 Cracked or rolled corn; 24-36% moisture
- 3. Ensile rapidly with iniform distribution in the silo and all kernels cracked.
- 4. Apply at the correct rate based on the moisture content.
- Pack well for good fermentation. Seal off with plastic sheeting. It is good practice to feed off the top four feet before sealing as good compaction is not possible near the top.



NGV 1 0 2010

- / 6	
ACTIVE INGREDIENT:	BY WT
Propionic Acid	68.00%
OTHER INGREDIENTS:	<u>32.00%</u>
Ammonium Hydroxide, Water, Citric Acid	
Nonylphenol ethoxylate, Dye	
TOTAL	100.00%
1 Gallon – 8.9 lbs	
FR. D. M. 150(5.10 FR. D. M. 150	C

EPA Reg. No. 15265-19 EPA Est. No. 15265-WI-001 KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED - Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow.

Do not induce vomiting unless told to do so by The poison control center or doctor. Do not give contribing the mouth to an

The poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED - Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING - Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center for doctor treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN - Probable mucosal damage may contraindicate the use of gastric lavage.

See side panel for additional precautionary statements.

Net	Wt:	2400	480	□ 260	□130	\Box	50
1101	**	L_14700	TOV		130	ш	J,

Manufactured by

Code:

WAUSAU CHEMICAL CORPORATION

P O Box 953, Wausau, WI 54402-0953

800-950-6656 715-842-2285

Fax #: 715-842-9059

E-mail: info@wausauchemical.com

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals DANGER/PELIGRO

Follow the manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentration. Do not reuse them. Corrosive. Causes irreversible eye damage. Do not get in eyes:

Wear protective eyewear (goggles, face shield, or safety glasses). Harmful if swallowed or absorbed through skin. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse. Wear chemical -resistant gloves, chemical-resistant aprons, chemical - resistant footwear, and goggles or face shield when loading application equipment unless a closed loading system is used. Avoid working near high concentrations of spray mist/vapor. Use with adequate ventilation. Wash thoroughly after handling.

USER SAFETY RECOMMENDATIONS

User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Users should wash hands before eating, drinking, chewing gum, using tobacco or using the noiler.

Environmental Hazards

For terrestrial uses. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Seal tightly. Do not store in direct sunlight. Store above zero degrees Fahrenheit

PESTICIDE DISPOSAL: Pesticides wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide/Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Offer rinsed container for recycling or reconditioning if available, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMERS AND NOTICE

It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of registrant, the manufacturer or seller. In no case shall the registrant, the manufacturer, or seller be liable for the consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer or user. Except as expressly provided herein, the Registrant, the Manufacturer, or Seller make no warranties, guarantees, or representations of any kind, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to, merchantability fitness for a particular purpose, use or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, buyer's or user's exclusive remedy, and the registrants, the manufacturer's or seller's total liability, shall be for damages not exceeding the cost of the product. No agent or employee of the registrant or seller shall be able to amend the terms of this label.

Specific Gravity 1.065 to 1.075 @ 70°F, pH 5.6 to 5.8. Freezes at -30°F. Freezing does not deteriorate product. However, citric acid may crystallize out. After warming, mix to get the citric acid back into colution.

