

C
B
A

85L

ACCEPTED
23 JUL 1979
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 15311-2

ACTIVE INGREDIENTS:

Disodium cyanodithioimidocarbonate	90.0%
Potassium N-methyldithiocarbamate	5.8%
INERT INGREDIENTS	90.0%
TOTAL	100.0%

This product contains 0.89 lb. of active ingredients per gallon and weighs 8.90 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, give patient doses of powdered charcoal immediately or all he can swallow of raw egg white, milk, gruel, or flour and water. Then induce vomiting with salt, soap, or mustard in warm water. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Do not discharge into lakes, streams, ponds or public waters unless in accordance with an NPDES permit. For guidance contact your Regional Office of the EPA.

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

85L is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits. An initial slug addition of 6.0 to 11.9 fluid ounces of 85L per 1000 gallons of water is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 2.0 to 11.9 fluid ounces of 85L per 1000 gallons of water should be employed every 1 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

STORAGE & DISPOSAL: Keep container closed when not in use. Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture, or residue that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place. Consult Federal, State, or Local disposal authorities for approved alternative procedures.

Manufactured by:

The Bond Chemical Company, Inc.
103 Pleasant Avenue
Upper Saddle River, NJ 07458

EPA REG. NO. 15311-2

EPA EST. NO. 15311-NJ-1

NET CONTENTS 30 Gallons

BEST DOCUMENT AVAILABLE

A
B