

PM 23
U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (75-767)
WASHINGTON, DC 20460

15440-32

EPA REGISTRATION NO.

15440-32

DATE OF ISSUANCE

SEP 16 1996

9/15
9/16/96

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

TERM OF ISSUANCE

Conditional

NAME OF PESTICIDE PRODUCT

MARKS 2,4-LL DMA 2

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

A. N. MARKS & COMPANY, LTD
Wyke Bradford
West Yorkshire
England BD12 9EU

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA section 5(c)(2) provided that you:

1. Submit and/or cite all data required for registration of this product under FIFRA sec. 5(c)(2) when the agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of this product under FIFRA sec. 4.
2. Submit one (1) copy of the final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 5(c). Your release for shipment of the product constitutes acceptance of these conditions. Attached copy of the label is enclosed for your records.

Joanne L. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

Joanne L. Miller

DATE

SEP 16 1996

ACCEPTED
with COMMENTS
In EPA Letter Dated

SEP 16 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

15440-32

MARKS

RECD EPA/01 P/DPD2

2,4-DB DMA 2

BROADLEAF HERBICIDE

JUL 30 1986

2/15

ACTIVE INGREDIENT:

4-(2,4-Dichlorophenoxy) butyric acid, dimethylamine salt* 26.85%

INERT INGREDIENTS: 73.15%

TOTAL 100.00%

*4-(2,4-Dichlorophenoxy) butyric acid equivalent 22.6% by weight or 2.0 pounds per gallon.

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

Corrosive. Causes irreversible eye damage. Wear goggles or face shield and rubber gloves when handling.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for at least 15 minutes. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

See side panel and inside booklet for additional PRECAUTIONARY STATEMENTS and NOTE TO PHYSICIAN.

Manufactured by:
A.H. Marks and Company Ltd.
Wyke Bradford
West Yorkshire, England BD129EJ
Telephone: 011 44 1274 691234

EPA Reg. No. 15440-GE
EPA Est. No. 15440-EN-1
07.22.96
Front Panel

NET CONTENTS ___ GALS.

PRECAUTIONARY STATEMENTS

DANGER

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Wear goggles or face shield. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for at least 15 minutes. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN

Repeat doses of activated charcoal are contraindicated if the patient has absent bowel sounds because of the risk of obstruction.

Maintain respiration, monitor ECG and blood gases. Correct acidosis with iv sodium bicarbonate. Control convulsions with diazepam and treat hypotension with volume replacement and inotropes (eg dopamine or dobutamine) as necessary. Monitor CPK for rhabdomyolysis.

Forced alkaline diuresis in the treatment of severe phenoxy acid poisoning is no longer recommended due to the risk of electrolyte imbalance. Urinary alkalization is less hazardous. The recommended regimen for urinary alkalization is as follows: To maintain urine pH greater than 7.5, adults should be given 50 ml boluses of 8.4% sodium bicarbonate iv and/or 1 l of 1.26% sodium bicarbonate + 40 mmol potassium iv over 4 hours. Children should be given 1 ml/kg of 8.4% sodium bicarbonate + 20 mmol potassium diluted in 500 ml dextrose saline infused at 2-3 ml/kg/hour.

These recommendations are subject to review and a Poison Control Center should be contacted in each case where treatment is likely to be necessary.

Probable mucosal damage may contraindicate the use of gastric lavage.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATION

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, waterproof gloves, shoes plus socks, and protective eyewear.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage or disposal. Store in original container in a dry secured storage area. Keep container tightly closed when not in use.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL CAUTIONS AND RESTRICTIONS: Do not apply this product through any type of irrigation system.

Do not use in or near a greenhouse.

GENERAL INFORMATION

Spray tank residues of 2,4-D or MCPA mixed with 2,4-DB DMA 2 can cause serious crop or ornamental plant injury. A sprayer previously used to apply these chemicals must be thoroughly cleaned with alkali and water before applying 2,4-DB DMA 2. Be sure sprayer is clean before applying 2,4-DB DMA 2.

Local conditions may affect the use of herbicides. Consult your State Agricultural Experiment Station, Farm Advisors, or Extension Weed Specialists for advice in selecting treatment from this label to best fit local conditions.

INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT.

Crop varieties vary in response to 2,4-DB and some are easily injured. Apply 2,4-DB DMA 2 only to varieties known to be tolerant to 2,4-DB. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-DB, consult your seed company, state Agricultural Extension Service or qualified crop consultant for advice. Be sure that use of this product conforms to all applicable laws, rules and regulations. Certain states have restrictions pertaining to application distances from susceptible crops. The applicator should become familiar with these laws, rules, or regulations and follow them exactly.

PRECAUTIONS

Mixing and Loading: Most cases of ground water contamination involving phenoxy herbicides such as 2,4-DB have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-DB pesticides at such sites to prevent contamination of ground water supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent ground water contamination.

Do not apply 2,4-DB DMA 2 directly to or otherwise permit it to come in contact with cotton, okra, grapes, tomatoes, fruit trees, vegetables, flowers, or other desirable crop or ornamental plants. Do not permit spray mist to drift onto susceptible plants since very small quantities of 2,4-DB can cause severe injury during the growing or dormant periods. Use coarse sprays to minimize drift. Do not apply with nozzles that produce fine spray droplets.

Drift from ground application may be reduced by: (1) keeping the spray boom as near to the crop as possible in order to obtain complete coverage; (2) by applying 10 gallons

or more of spray per acre; (3) by using no more than 20 pounds of pressure at the nozzle tips; and (4) by not spraying when wind exceeds 5 miles per hour.

Drift from aerial application may be reduced by: (1) applying as near to the target as possible to obtain adequate coverage; (2) by applying 5 or more gallons of spray per acre; (3) by using 20 pounds pressure or less at the nozzle tips; (4) by using nozzles which produce a coarse spray pattern; and (5) by spraying when there is no possibility for a temperature inversion at time of spraying.

Applications by aircraft, ground rig and hand sprayers should be carried out only when there is no hazard from spray drift.

APPLICATION PROCEDURES

2,4-DB DMA 2 can be applied by ground or aerial application. The following provides recommended methods of application for each crop.

GROUND APPLICATION:

Use a standard herbicide sprayer that provides uniform and accurate application. Sprayer should be equipped with screens no finer than 50 mesh in the nozzle tips and in-line strainers.

Select a spray volume and delivery system that will ensure thorough and uniform spray coverage. For optimum spray distribution and thorough coverage, use of flat fan nozzles (maximum tip size 8008) with a minimum spray pressure of 20 psi at the nozzle tips are recommended. Other nozzle types that produce coarse spray droplets may not provide adequate coverage of the weeds to ensure optimum control. Raindrop nozzles are not recommended as weed control with 2,4-DB DMA 2 may be reduced. In general a minimum spray volume of 10 gallons per acre (GPA) is recommended for optimum spray coverage. When using higher speed equipment, a maximum speed of 10 mph is suggested if field conditions cause excessive boom movement during application and subsequent poor spray coverage. Ground applications made when dry, dusty field conditions exist may provide reduced weed control in wheel track areas.

AERIAL APPLICATION:

Use orifice discs, cores and nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage. In general a minimum spray volume of 5 GPA and a maximum pressure of 20 psi are recommended.

MIXING INSTRUCTIONS:

2,4-DB DMA 2 alone: Fill the spray tank 1/2 to 3/4 full with clean water. Begin agitation and add the recommended amount of 2,4-DB DMA 2. Add water to the spray tank to the desired level. Maintain sufficient agitation to ensure a uniform spray mixture during application.

TANK MIXTURES: 2,4-DB DMA 2 can be applied in tank mixture with other herbicides registered for use on approved crops. Refer to the specific crop section for rate recommendations and other restrictions. To apply 2,4-DB DMA 2 in mixture with another product, fill the spray tank 1/2 to 3/4 full with clean water and begin agitation. If tank mixing with wettable powder, soluble powder, flowable or dry flowable products, add the powder or flowable product first. After the other herbicide is thoroughly mixed with water, add the recommended amount of 2,4-DB DMA 2 and add water to the spray tank to the desired level. If tank mixing the other product types, add the 2,4-DB DMA 2 first before adding the other product. Always mix one product in water thoroughly before adding another product or compatibility problems may occur. Never mix two products together without first mixing in water.

Maintain sufficient agitation while mixing and during application to ensure a uniform spray mixture. If spray mixture is allowed to remain without agitation for short periods of time, be sure to agitate until uniformly mixed before application.

COMPATIBILITY

2,4-DB DMA 2 may form an insoluble precipitate in very hard water. If you expect to mix 2,4-DB DMA 2 with very hard water, test compatibility by mixing a small amount of 2,4-DB DMA 2 in the proposed dilution ratios, shake and observe. A compatibility agent approved for use on growing crops such as UNITE or E-Z MIX may be tested to reduce precipitation. Whenever hard water is used to dilute 2,4-DB DMA 2, spray immediately and do not allow spray mixture to sit overnight.

GENERAL WEED LIST

2,4-DB DMA 2 will control or suppress the following weeds depending on weed height.

COMMON NAME	BOTANICAL NAME	MAXIMUM WEED HEIGHT*
velvetleaf	<i>Abutilon theophrasti</i>	1 inch
Virginia copperleaf	<i>Acalypha virginica</i>	1 inch
prickly sida (teaweed)	<i>Sida spinosa</i>	1 inch
common ragweed	<i>Ambrosia artemisiifolia</i>	1 inch
threeseed croton (goatweed)	<i>Croton lindheimeranus</i>	1 inch
lambsquarters	<i>Chenopodium album</i>	1 inch
wild mustard	<i>Sinapsis arvensis</i>	1 inch
field pennycress	<i>Thlaspi arvense</i>	1 inch
jimsonweed	<i>Datura stramonium</i>	1.5 inches
devilsclaw	<i>Proboscidea louisianica</i>	2 inches
pigweed	<i>Amaranthus ssp.</i>	3 inches
morningglory	<i>Ipomoea ssp.</i>	3 inches
cocklebur	<i>Xanthium ssp.</i>	3 inches
curled dock	<i>Rumex crispus</i>	3 inches
Russian thistle	<i>Salsola kali</i>	3 inches
yellow rocket	<i>Barbarea Vulgaris</i>	3 inches
sicklepod	<i>Cassia obtusifolia (L)</i>	2 inches
smartweed	<i>Polygonum ssp.</i>	3 inches

*Growth of taller weeds will be only suppressed.

SOYBEANS

Preplant through preemergence:

Apply 0.7-0.9 pint/A 2,4-DB DMA 2 + 0.5% V/V non-ionic surfactant for control of emerged cocklebur, annual morningglories and other susceptible broadleaf weeds before planting or before crop emergence. Apply when weeds are small and actively growing (see GENERAL WEED LIST). 2,4-DB DMA 2 may not give complete control of larger overwintered mustards.

Postemergence broadcast (over the top):

Apply 0.7-0.9 pint/A 2,4-DB DMA 2 for control of emerged cocklebur, annual morningglories and other susceptible broadleaf weeds. Apply when weeds are small and actively growing (see GENERAL WEED LIST). Apply to soybeans from 7 to 10 days before bloom up to mid-bloom when soybeans are about knee-high and growing actively. Soybean foliage should be dark green indicating that nodulation and nitrification are under way. Post emergence broadcast application at these rates prior to or after this application timing is not recommended as reduced flowering and yield may result. DO NOT APPLY 2,4-DB DMA 2 postemergence broadcast to soybeans grown in IA, IL, IN, KS, KY (except the Purchase area), MO (except the MO bootheel), MI, MN, NE, ND, OH, SD and WI.

Postemergence directed band:

Apply 0.7-0.9 pint 2,4-DB DMA 2 per broadcast acre as a directed band treatment to control emerged cockleburs and annual morningglories up to 3 inches tall. To control other susceptible broadleaf weeds up to 1 inch tall, apply 1.4 to 1.6 pints per broadcast acre as a directed band treatment. Apply no more than twice per season to minimize the potential for stunting crops. Apply when soybeans are 8 or more inches tall with sprayer nozzles mounted to insure proper placement of spray on only the lower 1/3 of the soybean plants. Do not allow spray to contact growing terminals of beans as excessive crop injury will result. Do not mount nozzles on booms with drop pipes or on cultivators without gauge wheels. Use flat fan type nozzles, 8001 or larger or the equivalent with a minimum nozzle pressure of 20 psi and at least 10 gallons of spray volume per acre.

TANK MIXTURES OF 2,4-DB DMA 2 AND OTHER HERBICIDES IN SOYBEANS

Applying tank mixtures of 2,4-DB DMA 2 preplant/preemergence or postemergence with other soybean herbicides as recommended by local weed control authorities can reduce competition from early weed populations and can improve weed control or control of mid-to-late-season weed populations, thus minimizing the likelihood of yield reduction in soybean fields with heavy broadleaf weed infestations. However, treating soybeans under stress (as from drought or disease such as *Phytophthora* root rot) or in any other manner not directed on this label can cause crop injury and yield reduction. Follow the local

recommendations of your state Cooperative Extension service, or other agricultural weed control authority. Always follow the recommendations of any product used in tank mixture with 2,4-DB DMA 2.

Gramoxone® Extra:

Apply 0.5-0.75 pint/A 2,4-DB DMA 2 + 1.5-5 pints/A Gramoxone® Extra + 0.5% V/V non-ionic surfactant for improved control of emerged annual morningglory, common cocklebur, marestail and other problem weeds. Apply to small actively growing weeds. Refer to the Gramoxone Extra label for full list of weed species controlled and specific application stage and rate recommendations. Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.

Prowl®:

Apply 0.7-0.9 pint/A 2,4-DB DMA 2 + 1.0-3.0 pints/A Prowl® + 0.5% V/V non-ionic surfactant for control of emerged cocklebur, annual morningglories and other susceptible broadleaf weeds. Apply when broadleaf weeds are actively growing and small (see GENERAL WEED LIST). 2,4-DB DMA 2 mixtures may not give complete control of larger overwintered mustards. Best results will be achieved by adding a non-ionic surfactant to the spray tank when making a preplant application. Apply up to 45 days prior to soybean planting. Do not apply a 2,4-DB DMA 2 + Prowl tank mixture at or after planting north of Interstate 80. Surface applications of Prowl tank mixtures north of Interstate 80 require at least 1 inch rainfall or mechanical incorporation prior to planting or crop injury may result. Do not apply this tank mixture after crop emergence.

Roundup® or Honcho®:

Apply 0.5-0.75 pint/A 2,4-DB DMA 2 + 1-1.5 pints/A Roundup® or Honcho® + 0.5-1.0% V/V non-ionic surfactant for improved control of emerged annual morningglories, common cocklebur and other problem weeds. Apply to small actively growing weeds. Refer to the Roundup or Honcho labels for specific rate recommendations, application stage and weed species controlled. Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.

Pursuit Plus®:

Apply 0.5-0.75 pint/A 2,4-DB DMA 2 + 2.5 pints/A Pursuit Plus® + 0.25% V/V non-ionic surfactant for improved control of emerged mustards, field pennycress and other problem weeds. Apply when broadleaf weeds are actively growing and small (see GENERAL WEED LIST). Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.

Pursuit®:

Apply 2-3 fl. oz./A 2,4-DB DMA 2 + 4 fl. oz./A Pursuit® for improved control of common and giant ragweed, morningglories, and other broadleaf weeds. Refer to the Pursuit label regarding the use of surfactants and weed species controlled. Apply this tank mixture any time after soybean emergence but no later than 85 days before harvest.

Scepter® or Scepter 70DG®:

Apply 0.5-0.75 pint/A 2,4-DB DMA 2 + 0.66 pint/A Scepter® or 2.8 oz./A Scepter 70DG® + 0.25% V/V for improved control of emerged mustards, field pennycress and other problem weeds. Apply when broadleaf weeds are actively growing and small (see GENERAL WEED LIST). Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.

Squadron®:

Apply 0.5-0.75 pint/A 2,4-DB DMA 2 + 3 pints/A Squadron® + 0.25% V/V non-ionic surfactant for improved control of emerged mustards, field pennycress and other problem weeds. Apply when broadleaf weeds are actively growing and small (see GENERAL WEED LIST). Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.

Basagran®:

Apply 2 fl oz/A 2,4-DB DMA 2 + 1.5-2 pints/A Basagran® to improve control of annual morningglories. Apply to vines up to 6 inches long in the southern states of AL, AR, FL, GA, LA, MS, NC, OK, SC, TN, TX and VA or a maximum of 10 inches long in all other states. Apply postemergence to soybeans. Under certain conditions soybean foliage may burn, crinkle and bronze following application.

Blazer®:

Apply 2 fl oz/A 2,4-DB DMA 2 + 1.5-2 pints/A Blazer to improve control of larger morningglories, cockleburs, common ragweed, jimsonweed and pigweed. Apply when weeds are actively growing and before they reach 12 inches. Apply postemergence to soybeans. Under certain conditions, soybean foliage may burn, crinkle and bronze following application. Soybean yield may be reduced. Do not add surfactant or crop oil to this mixture as increased crop injury may result.

Reflex®:

Apply 2-3 fl oz/A 2,4-DB DMA 2 + 1-1.5 pints/A Reflex® for improved control of annual morningglories, giant ragweed and cocklebur. Apply to actively growing weeds in seedling stage of growth. Apply postemergence to soybeans. Under certain conditions bronzing, crinkling or spotting of soybean foliage may occur.

Sencor DF®:

Apply 0.9 pint/A 2,4-DB DMA 2 + 0.33-0.66 lb/A Sencor DF® for improved control of cocklebur, annual morningglories and other broadleaf weeds. Apply before weeds are 3 inches tall. A non-ionic surfactant may be added to improve broadleaf weed control. Apply to soybeans as a directed band treatment only when soybeans are at least 8 inches high, with spray or nozzles mounted to insure proper placement of spray on no more than the lower 1/3 of the soybean plants. Do not apply directly to soybean plants or serious crop injury will occur. Soybean leaves contacted by spray will be killed. Follow all variety restrictions on the full Sencor DF label.

RESTRICTIONS AND LIMITATIONS FOR USE ON SOYBEANS

Beans stressed by drought or other influences should not be sprayed. Do not use this product on soybeans that show symptoms of disease such as Phytophthora root rot.

Do not graze or feed soybean hay within 60 days after application of 2,4-DB DMA 2 tank mix application.

Do not harvest soybeans within 60 days after spray application.

Do not treat soybeans with a tank mixture of 2,4-DB DMA 2 and Sevin® Carbaryl Insecticide as severe injury may result.

When preplant through preemergence treatment is followed with a 2,4-DB DMA 2 postemergence application, the cumulative rate should not exceed 1.6 pints per acre per season.

Follow all restrictions and limitations of any product used in tank mixture with 2,4-DB DMA 2.

Do not use 2,4-DB DMA 2 alone or in tank mixture as a preplant through preemergence application to soybeans in California.

PEANUTS

Apply 0.8-1.0 pint/A 2,4-DB DMA 2 for control of annual morningglories, cocklebur and other broadleaf weeds. Apply when weeds are small and actively growing (see GENERAL WEED LIST). A second application may be made for late germinating cocklebur and morningglories. Apply to peanuts 2 to 12 weeks after planting in the states of AL, AR, FL, GA, LA, NC, SC, TN, and VA. In NC, SC and VA, do not apply later than 45 days before harvest.

For control of annual morningglory and cocklebur in TX, OK and NM, apply 0.8-1.0 pint/A 2,4-DB DMA 2 before weeds are 3 inches tall. For optimum control of other susceptible broadleaf weeds (see GENERAL WEED LIST), apply 1.6 pints/A. For optimum prickly sida suppression, make a second application 14 days later. Apply to peanuts 2 to 12 weeks after planting. Do not apply later than 100 days after planting or 30 days before harvest.

RESTRICTIONS AND LIMITATIONS FOR USE ON PEANUTS

Do not feed treated peanut vines or peanut hay to livestock.

Do not apply 2,4-DB DMA 2 if peanut plants are under stress from drought as injury may occur.

SEEDLING AND ESTABLISHED ALFALFA

Apply 1-2 quarts/A 2,4-DB DMA 2 for control of emerged lambsquarters, pigweed, field pennycress, wild mustard, common ragweed, cocklebur, yellow rocket, Russian thistle and annual morningglory species less than 1 inch high. Use 2-3 quarts/A for control of these weeds up to 3 inches high. Use the higher rates in dry, low humidity growing areas. Apply 3 quarts/A for control or suppression of smartweed and curled dock up to 3 inches tall. Apply postemergence to seedling or established alfalfa. Alfalfa should be healthy and actively growing for greatest selectivity; twisting of stems and malformation of leaves may occur. Under most conditions this response is usually outgrown. A non-ionic surfactant at 0.25% V/V may improve weed control in seedling alfalfa grown in dry, low humidity areas only. The surfactant may cause some twisting of stems and malformation of leaves. This response is usually outgrown under most conditions. 2,4-DB DMA 2 may not adequately control overwintered broadleaf weeds including field pennycress and mustards.

For control of emerged susceptible broadleaf and grass weeds that are actively growing, apply 1-3 quarts/A 2,4-DB DMA 2 + 1-2.5 pints/A Poast® + 2 pints/A Crop Oil Concentrate. Refer to the recommendations above and the Poast label for weeds controlled and application timing for full use recommendations. Alfalfa should be healthy and actively growing for greatest crop tolerance. Established alfalfa is less tolerant to 2,4-DB DMA 2 than in the seedling stage of growth. Some yellowing and burning of alfalfa foliage, stem and leaf malformation may occur with this tank mixture. Alfalfa will generally outgrow this response. Balance the severity of your grass and broadleaf weed problem with the potential for crop injury. Do not add non-ionic surfactant, Dash®, UAN solution or ammonium sulfate to this tank mixture.

RESTRICTIONS AND LIMITATIONS FOR USE ON ALFALFA

Do not graze established alfalfa or feed straw or hay from treated established alfalfa to livestock within 30 days after application. Do not graze or feed seedling alfalfa within 60 days after application.

Do not apply when crop is stressed from lack of moisture. Do not spray when the temperature exceeds 90°F and/or is predicted to exceed 90°F during the three days following application.

Do not add any wetting agents or detergents to the spray solution unless as specified on this label.

Rainfall or overhead irrigation within 7-10 days following a 2,4-DB DMA 2 application can cause unacceptable crop injury. For irrigated crops, apply 2,4-DB DMA 2 as soon as possible after irrigation. Delay the next irrigation for 7-10 days after spraying.

Follow all restrictions and precautions of any product used in tank mixture of 2,4-DB DMA 2.

LIMITATION OF LIABILITY

All goods supplied by us are of high grade and we believe them to be suitable for all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods as described. No responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

Gramoxone and Reflex are registered trademarks of ICI Americans, Inc.
Prowl, Pursuit Plus, Scepter, Scepter 70DG and Squadron are registered trademarks of American Cyanamid Company.

Roundup and Honcho are registered trademarks of Monsanto Company.

Basagran, Blazer, Poast and Dash are registered trademarks of BASF AG.
Sencor is a registered trademark of Bayer AG, Germany.
Pursuit is a registered trademark of Rhone-Poulenc Ag. Company.