

WARNING

- MAY BE FATAL IF SWALLOWED. DO NOT BREATHE SPRAY MIST.
- MAY BE ABSORBED THROUGH SKIN.
- DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING.
- AVOID CONTAMINATION OF FEED AND FOODSTUFFS.
- WASH THOROUGHLY AFTER USING.

DO NOT reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

Destroy when empty.

NOTICE

Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling, and accepts the product on these conditions.

WARNING

The product is toxic to bees and should not be applied when bees are actively visiting the area.

SOUTHLAND
AGRICULTURAL
CHEMICALS

TOXAPHENE EM-6
 Emulsifiable Liquid
 (Contains 6 lbs. Toxaphene Per Gal.)
 AN AGRICULTURAL INSECTICIDE

ACCEPTED
 1975
 COMMENTS

ACTIVE INGREDIENTS:

Toxaphene*	58.50%
Xylene-Range Aromatic Hydrocarbon Solvent	38.50%
INERT INGREDIENTS:	3.00%
TOTAL	100.00%

*Technical Chlorinated Camphene (Chlorine Content 67% to 69%)

**DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME!
PROTECT FROM TEMPERATURES BELOW 0 F.**

**WARNING: KEEP OUT OF THE REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS**



Pour recommen
agitator running
mix readily, but
emulsifiable liqu
has proved them

OBSEI
U

DO NOT use
fed to dairy ani
slaughter.

This product
life. Birds feedi
Keep out of lak
where runoff is
weather conditio
NOT contamina
disposal of wa
specified on this

DO NOT use
Observe use
specific crops.

The Federal f
official toleranc
order th it residu
exceed establish
mended rates, a
fied.

DIRECTIONS FOR USE

Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing, and spraying operations. DO NOT allow mixture to stand. If mixture does not mix readily, but tends to separate as an oily layer, DO NOT use as injury to plants may result. DO NOT combine emulsifiable liquids with wettable powders in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible.

OBSERVE THE FOLLOWING USE PRECAUTIONS

DO NOT use this product on feed or forage to be fed to dairy animals or to animals being finished for slaughter.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. DO NOT contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DO NOT use on cucurbits (cantaloupe, cucumber.)

Observe use limitations given on the label for specific crops.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerance, use only at recommended rates, and DO NOT apply closer than specified.

RECOMMENDATIONS

Dosages are given in terms of pints per acre. Use in sufficient water for thorough coverage. (Approximately 100-125 gallons).

For airplane application, use approximately 3 to 10 gallons of water per acre, consistent with crop growth and good coverage.

COTTON: For control of Armyworms, Bollworm, Climbing Cutworms, Grasshoppers, apply 1-1/3 quarts per acre. DO NOT graze dairy animals or animals being finished for slaughter on cotton or cotton trash.