



ACCEPTED

JUN 9 1998
Under the Federal Insecticide,
Fungicide and Rodenticide Act,
as amended, this pesticide
registered under
EPA Reg. No. 17545-2

COTTON-AIDE HC®

Cotton Defoliant

READ ENTIRE LABEL BEFORE USING THIS PRODUCT

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution: Causes eye irritation. Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. (Coarse sprays are less likely to drift).

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. DO NOT contaminate water when cleaning equipment or disposing of equipment washwaters.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. STORAGE: DO NOT store near feed or food products. Store in original container only. Keep container tightly sealed when not in use. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

ACTIVE INGREDIENTS:

Sodium cacodylate	28.4%
Dimethylarsinic acid (Cacodylic acid)	4.9%

OTHER INGREDIENTS:

Total	66.7%
Total	100.0%

Total arsenic, all in water soluble form, expressed as elemental: 15.96%
Contains 3.25 pounds cacodylic acid equivalent per gallon.

EPA Reg. No. 17545-2

EPA Est. No. 48498-CA-1

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.
IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, DO NOT give anything by mouth and do not induce vomiting. IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists. IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

See side panel for additional precautionary statements.

CONDITIONS OF SALE AND WARRANTY

Monterey Chemical Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated when used in accordance with the use directions under normal conditions. Monterey Chemical Company neither makes, nor authorizes any agent or representative to make, any other warranties, express or implied, including fitness or merchantability.

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, presence of other materials, or other influencing factors, all of which are beyond the control of Monterey Chemical Company and Seller. Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage and use of this material not in strict accordance with directions given herewith. If no case shall Monterey Chemical Company or the Seller be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by Monterey Chemical Company and is accepted as such by the Buyer.

Attention: This product contains cacodylic acid, which is a chemical known to the State of California to cause cancer.

NET CONTENTS: 5 GALLONS / 18.93 LITERS

Manufactured for:

MONTEREY CHEMICAL COMPANY

P.O. Box 5317 • Fresno, CA 93755 • (209) 499-2100

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and domestic animals off treated areas until this material has been washed into the soil.

GENERAL INFORMATION: COTTON-AIDE HC may be used as a harvest aid for the defoliation of cotton. COTTON-AIDE HC is also a non-selective postemergence herbicide. It will kill vegetation on contact. However, its phytotoxic properties are quickly inactivated upon contact with the soil.

CARE OF EQUIPMENT: Although COTTON-AIDE HC is only moderately corrosive, do not apply with any applicator having zinc, aluminum, magnesium or alloy components. Clean equipment after use by flushing with water. Avoid spray contact with aircraft components to minimize corrosion.

CHEMIGATION: DO NOT apply this product through any type of irrigation system.

MIXING AND APPLICATION INSTRUCTIONS: Fill spray tank with desired quantity of water. While agitating, add the recommended amount of COTTON-AIDE HC. Thoroughly mix prior to application. The sprayer should be carefully calibrated for best results. Apply the spray solution at the proper rate in the proper amount of water to the properly measured area.

DEFOLIATION OF COTTON: Apply COTTON-AIDE HC before harvest when at least 50% of bolls are open. Dryland Cotton: Use a rate of 2-2½ pints COTTON-AIDE HC per acre.

Irrigated Cotton: Use a rate of 2½-3 pints COTTON-AIDE HC per acre for stand alone treatments and 1-3 pints COTTON-AIDE HC per acre for tank mix applications as specified below.

For proper defoliation, spray solution must contact all leaves. Proper equipment calibration is essential. If applied by aircraft, use 5-10 gallons of water per acre. If applied by ground equipment, use 15-20 gallons of water per acre. Addition of a non-ionic surfactant, which has been cleared for application to growing crops, may enhance coverage and overall performance. Use surfactants in accordance with their manufacturer's directions, precautions, and restrictions.

COTTON-AIDE HC may be tank-mixed with Accelerate®, DEF® 6, Dropp® 50WP, Folex® 6-EC, Ginstar®, Prep™ or other ethephon products, Defo® 6W or other sodium chlorate products, or paraquat products for cotton desiccation/defoliation according to local practices or seasonal conditions. Do so in accordance with the most restrictive label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. To assure tank mixture compatibility, mix the correct proportions of the components in a small container and let stand five minutes. If the combination remains mixed, or can be readily remixed, the tank mixture may be considered compatible.

DO NOT feed treated foliage to livestock or graze treated areas.

GENERAL WEED CONTROL: COTTON-AIDE HC will top-kill annual and perennial weeds in non-crop areas around buildings, industrial sites, storage yards and vacant lots; rights-of-ways and roadways; and fencerows and drainage ditchbanks. Apply 2½ gallons of COTTON-AIDE HC in 65-100 gallons of water per acre. Apply COTTON-AIDE HC as a broadcast treatment or directed spray to weeds around desirable plants. Wet foliage of actively growing weeds to run-off. Should growth reoccur, repeat five days after first application. Sunshine and temperatures above 75°F will accelerate herbicidal activity.

Note: COTTON-AIDE HC contains a suitable wetting agent for normal use. However, for hard-to-kill and larger weeds in certain arid regions, addition of a non-ionic surfactant may enhance weed control.

Avoid spray contact with desirable plants. DO NOT use in vegetable gardens or around crops grown for food or feed. DO NOT contaminate water used for irrigation or domestic purposes.

(Accelerate®-Elf Atochem North America Inc.; DEF®-Bayer; Defo®-Drexel Chemical Co.; Dropp® and Ginstar®-AgriEvo USA Co.; Folex® and Prep™-Rhône-Poulenc Ag Co.)