

17545-6 PA 23 1702
Proposed Labeling 8/2/88

WEED ENDER

Contact weed and grass killer for homeowner use on non-planted areas or as a liquid edger around patios, curbs, fences, walkways, driveways, trees, and shrubs. Non-staining -- Odorless.

Covers 1,125 sq. ft.

***ACTIVE INGREDIENTS:**

Sodium Cacodylate	15.10%
Dimethylarsinic Acid (cacodylic acid)	2.60%
INERT INGREDIENTS:	82.30%

TOTAL 100.00%

*Total Arsenic (as elemental) all in water soluble form 8.45%
Contains 1.60 lbs. Cacodylic Acid Equivalent per gallon.

EPA Reg. No. 17545-6
EPA Est No. 48498-CA-01

NET CONTENTS: ONE PINT

**KEEP OUT OF REACH OF CHILDREN
CAUTION:**

**PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals**

Harmful if swallowed or inhaled. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. Causes eye irritation. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep children and domestic animals off treated areas until washed into soil. Avoid contamination of feed or food products. Do not graze treated areas.


See side panel for additional Precautionary Statements

This product is manufactured by:

Monterey Chemical Company
P.O. Box 5317
Fresno, California 93755

ACCEPTED
DEC 6 1988

17545-6



2 of 2

STATEMENT OF PRACTICAL TREATMENT

If swallowed, call a physician immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
If on skin, wash with plenty of soap and water. Get medical attention if irritation persists.
If inhaled, remove victim to fresh air. Apply artificial respiration, preferably mouth to mouth, if indicated.
If in eyes, flush with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water or wetlands. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CHEMIGATION: DO NOT apply this product through any type of irrigation system

To prepare spray solution, fill spray tank with recommended amount of water. Add WEED ENDER as recommended below and mix thoroughly by shaking or other agitation. If mixed spray is allowed to stand for any length of time, it should be shaken or agitated before using.

WEED ENDER	Water	Area Covered
1/2 pint*	2 - 2 1/2 gallons	562 sq. ft.
1 pint	4 - 5 gallons	1,125 sq. ft.
2 pints	8 - 10 gallons	2,250 sq. ft.

*Equivalent to 8 fluid ounces or 16 tablespoons.

Apply WEED ENDER to top-kill actively growing, undesirable annual and perennial weeds by thoroughly wetting foliage to run-off. However, certain hard-to-kill species, such as Bermudagrass and Dallisgrass may require repeat applications. Should regrowth occur, retreat as needed at 5-day intervals. Sunshine and temperatures above 80°F will enhance effectiveness.

DO NOT allow spray to drift onto desirable plants.
DO NOT use WEED ENDER on vegetable gardens.
DO NOT track from treated areas onto turf.

NON-PLANTED AREAS: WEED ENDER may be used for general weed control or as a liquid edger around buildings, storage areas, vacant lots, patios, walkways, driveways, curbs and fences.

ORNAMENTALS: Apply WEED ENDER as a directed spray to unwanted vegetation or turf around ornamental plants, shrubs and trees. Avoid spray contact with foliage, stems or bark of desirable plants.

LAWN ESTABLISHMENT AND RENOVATION: WEED ENDER may be used to destroy weeds before lawn establishment or weeds and existing turf before lawn renovation. For best results, mow weeds or turf to about 1" height before treatment. Apply WEED ENDER according to the above table. If needed, repeat application as indicated above may be needed 5 days after first treatment.

Caution: Delay reseeding for at least 5 days after last application. Dead vegetation need not be removed unless matted turf or thatch prevents lawn seed from making soil contact. Precede reseeding by heavily watering treated area. Seed may be worked into soil, if necessary, by scarifying or raking. A light top-dressing of screened peat moss will help retain moisture during seed germination. Keep reseeded area moist.

STORAGE AND DISPOSAL

STORAGE: Do not store near feed or food products.

DISPOSAL: Do not REUSE CONTAINER. Securely wrap original container in several layers of newspaper and discard in trash. Clean sprayer with water and discard rinse water onto sprayed area.

BEFORE USING ANY HERBICIDE, STOP AND READ THE LABEL

WARRANTY - CONDITION OF SALE: DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however, impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of Monterey Chemical Company or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result. ALL SUCH RISKS ARE ASSUMED BY THE BUYER.

Monterey Chemical Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the direction for use as modified by the above. Monterey Chemical Company makes no other warranties, expressed or implied, including FITNESS or MERCHANTABILITY. In no case shall Monterey Chemical Company or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by Monterey Chemical Company and is accepted as such by the Buyer.

0188/0188(01)5.c.d

