


17545-8

12/04/2006

117

	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460		EPA Reg. Number: 17545-8	Date of Issuance: DEC - 4 2006
	NOTICE OF PESTICIDE: <input type="checkbox"/> Registration <input checked="" type="checkbox"/> Reregistration (under FIFRA, as amended)		Term of Issuance: Unconditional	
			Name of Pesticide Product: TURFLON* Ester	
Name and Address of Registrant (include ZIP Code): Monterey AgResources P.O. Box 35000 Fresno, CA 93745-5000				
Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.				
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>ff</p> <p>Based on your response to the Reregistration Eligibility Document(s), EPA has reregistered the product listed above. The label is acceptable provided the following revision is made:</p> <ol style="list-style-type: none"> 1. Add " Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in <u>Washington Toxics Coalition, et al. v. EPA, C01-0132C, (W.D. WA)</u>. For further information, please refer to http:// www.epa.gov/espp/wtc. 2. Prohibit aerial application or put the aerial drift mitigation language from the from the Triclopyr RED on the label. 3. Under Conditions of Sale and Warranty: <ol style="list-style-type: none"> a) Change " The directions for use of this product are believed to be reliable and should be followed carefully." to " The directions for use of this product are believed to be reliable and must be followed carefully." b) Change " In no case shall MAR or the seller be liable for consequential, special, indirect or incidental damages or losses resulting from the handling of this product." to " To the extent consistent with applicable law, MAR or the seller be liable for consequential, special, indirect or incidental damages or losses resulting from the handling of this product." 4. Add to Sublabel B Non-Commercial/ Residential Uses under Hazards to Humans and Domestic Animals: " Wear long sleeved shirt and long pants, shoes and socks when mixing or applying this product." 				

- 5. On both sub-labels, specify a maximum of 8 lbs/ae/A/year.
- 6. Under *Environmental Hazards*, place the following Groundwater Advisory:

This chemical has properties and characteristics associated with chemical detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination."

- 7. On the Commercial/ Agricultural Use and sub-label, under PPE section in Precautionary Statements and under early-entry in Agricultural Use Requirements box add "protective eyewear."

Enclosed is a copy of your label stamped "Accept" with comment. This action is taken under the authority of action 4(g) (2) (c) of the Federal Insecticide, Fungicide, and Renticide Act, as amended.

Reregistration under this section does not eliminated the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Signature of Approving Official. James Tompkins Product Manager 25 Herbicide Branch Registration Division (7505P)	Date: 12-4-06
---	------------------

MASTER LABEL

TURFLON* Ester **Specialty Herbicide**

ACCEPTED
with COMMENTS
In EPA Letter Dated:

DEC - 4 2006
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

17545-8

**Controls Annual and Perennial Broadleaf Weeds
and Bermudagrass in Ornamental Turf**

Active Ingredient:

Triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid, butoxyethyl ester 61.6%

Other Ingredients: 38.4%

Total: 100.0%

Triclopyr acid equivalent: 44.3% (4 lb/gal).

Contains petroleum distillates.

EPA Reg. No. 17545-8

EPA Est. No. 48498-CA-1

*Trademark of DowAgroSciences LLC

**SUBLABEL A - Commercial / Agricultural Uses (Subject to the Worker
Protection Standard)**

**SUBLABEL B - Non-Commercial / Residential Uses (Not Subject to the
Worker Protection Standard)**

Manufactured for:

Monterey AgResources

P. O. Box 35000 • Fresno, CA 93745-5000 • (559) 499-2100
www.montereyagresources.com

Controls Annual and Perennial
Broadleaf Weeds and Bermudagrass
in Ornamental Turf

TURFLON* Ester

SPECIALTY HERBICIDE

EPA Reg. No. 17545-8
EPA Est. No. 48498-CA-1

ACTIVE INGREDIENT:

Triclopyr: 3,5,6-trichloro-2-pyridinyl-
oxyacetic acid, butoxyethyl ester 61.6%
OTHER INGREDIENTS: 38.4%
TOTAL 100.0%

Triclopyr acid equivalent: 44.3% (4 lb/gal).
Contains petroleum distillates.

*Trademark of DowAgroSciences LLC

**KEEP OUT OF REACH
OF CHILDREN**
CAUTION / PRECAUCION
Si usted no entiende la etiqueta, busque a
alguien para que se la explique a usted en
detalle. (If you do not understand the label,
find someone to explain it to you in detail.)

FIRST AID
IF SWALLOWED: Call a Poison Control
Center or doctor immediately for treatment
advice. Have person sip a glass of water if
able to swallow. Do not induce vomiting
unless told to do so by a Poison Control
Center or doctor. Do not give anything by
mouth to an unconscious person.
IF IN EYES: Hold eye open and rinse slowly and
gently with water for 15-20 minutes. Remove
contact lenses, if present, after the first 5
minutes, then continue rinsing. Call a Poison
Control Center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off
contaminated clothing. Rinse skin
immediately with plenty of water for 15-20
minutes. Call a Poison Control Center or
doctor for treatment advice.

Have the product container or label with you
when calling a Poison Control Center or
doctor, or going for treatment.

Note to physician: This product may pose
an aspiration pneumonia hazard. Contains
petroleum distillates.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Causes
moderate eye irritation. Avoid contact with
eyes, skin or clothing during and after
application. Prolonged or frequently repeated
skin contact may cause allergic reactions in
some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)
Some materials that are chemical-resistant to
this product are listed below. If you want more
options, follow the instructions for category E
on an EPA chemical resistance category
selections chart.

- Applicators and other handlers must wear:**
- Long-sleeved shirt and long pants
 - Chemical-resistant gloves [such as Barrier Laminate, Nitrile Rubber, Neoprene Rubber, or Viton]
 - Shoes plus socks.

Follow manufacturer's instructions for cleaning/
maintaining PPE. If no such instructions for
washables are given, use detergent and hot
water. Keep and wash PPE separately from
other laundry.

USER SAFETY RECOMMENDATIONS:
Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS
This pesticide is toxic to fish. Do not apply
directly to water, or to areas where surface
water is present, or to intertidal areas below
the mean high water mark. Do not contaminate
water when cleaning equipment or disposing
of equipment washwaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS
Do not use or store near heat or open flame.

DIRECTIONS FOR USE
It is a violation of Federal law to use this
product in a manner inconsistent with its
labeling. Read all Directions for Use carefully
before applying. Do not apply this product in
a way that will contact workers or other
persons, either directly or through drift. Only
protected handlers may be in the area during
application. For any requirements specific to
your state or tribe, consult the agency
responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS
Use this product only in accordance with its
labeling and with the Worker Protection
Standard, 40 CFR Part 170. This Standard
contains requirements for the protection of
agricultural workers on farms, forests,
nurseries, and greenhouses, and handlers
of agricultural pesticides. It contains
requirements for training, decontamination,
notification, and emergency assistance. It
also contains specific instructions and
exceptions pertaining to the statements on
this label about personal protective
equipment (PPE), and restricted-entry
interval. The requirements in this box only
apply to uses of this product that are covered
by the Worker Protection Standard.

Do not enter or allow workers entry into
treated areas during the restricted entry
interval (REI) of 12 hours.
PPE required for early entry to treated areas
that is permitted under the Worker Protection
Standard and that involves contact with
anything that has been treated, such as
plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves [such as Barrier Laminate, Nitrile Rubber, Neoprene Rubber, or Viton]
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS
The requirements in this box apply to uses
of this product that are NOT within the scope
of the Worker Protection Standard for
agricultural pesticides (40 CFR Part 170).
The WPS applies when this product is used
to produce agricultural plants on farms,
forests, nurseries, or greenhouses.
For applications to turf other than sod
farms, do not allow entry into treated areas
until sprays have dried unless coveralls,
chemical resistant gloves, and shoes plus
socks are worn.

GENERAL INFORMATION
(Read Carefully Before Using)
TURFLON Ester herbicide is recommended for the
control of actively growing annual and perennial
broadleaf weeds and kikuyugrass in perennial
bluegrass, perennial ryegrass or tall fescue

ornamental turf including these turfgrasses in sod farms and golf courses.

In Arizona: The state of Arizona has not approved this product for use on sod farms.

DO NOT apply this product through any type of irrigation system. DO NOT apply to ditches used to transport irrigation water. DO NOT apply where runoff or irrigation water may flow onto susceptible crops.

Local conditions may affect the use of herbicides. Consult your local specialist for advice in selecting treatments from this label to best fit local conditions.

TURFLON Ester may injure certain turfgrass species. DO NOT apply to bahiagrass, bentgrass, bermudagrass, centipedegrass, St. Augustinegrass or zoysiagrass, unless turf injury can be tolerated.

DO NOT apply to golf course greens.

DO NOT apply TURFLON Ester to exposed roots of shallow rooted trees and shrubs.

DO NOT apply TURFLON Ester herbicide directly to or otherwise permit it to come in contact with or permit spray mists containing TURFLON Ester to drift onto cotton, grapes, tobacco, vegetable crops, flowers, fruit or orchard trees, shrubs or other desirable broadleaf plants.

Avoid Injurious Spray Drift: Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. DO NOT spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. Spray drift can be reduced by spraying when wind velocity is low, keeping the spray boom as low as possible, using nozzles that deliver larger spray droplets, and by using higher volume sprays and lower pressures.

Broadleaf weeds controlled by TURFLON Ester include:

Black medic	Lespedeza
Bull thistle	Matchweed
Burdock	Mustard
Canada thistle	Oxalis
Chicory	Plantain
Clover	Ragweed
Creeping beggarweed	Smartweed
Curly dock	Sweet clover
Dandelion	Vetch
Field bindweed	Wild carrot
Goldenrod	(Queen Anne's lace)
Ground ivy	Wild lettuce
Kikuyugrass**	Wild violet
Lambsquarters	Yarrow

**See control of Kikuyugrass under "Application Directions".

APPLICATION DIRECTIONS

Foliar sprays should be applied during warm weather, from early spring through fall, when weeds are actively growing. Broadleaf weeds germinate at different times. Only emerged weeds present at the time of application will be controlled. Newly seeded turf should be mowed 2 or 3 times before being treated. When making applications to mature plants, hard to control species, or during drought conditions, use higher rates. Application under drought conditions may provide less than desirable

results. Use low pressure sprays to minimize spray drift. DO NOT water for 24 hours after application.

Reseeding Precaution: DO NOT reseed for 3 weeks after application. This precaution does not apply when bermudagrass turf is overseeded with perennial ryegrass at a minimum reseeding rate of 400 lbs. per acre.

Mixing Instructions

When TURFLON Ester is mixed with water, it forms an emulsion (not a solution) and separation may occur unless the spray mixture is agitated continuously.

Add about one-half the required amount of clean water to the spray tank. Start agitation and add the recommended amount of TURFLON Ester. Provide moderate agitation while completing the addition of water and during application.

Conversion Table	
1 ounce = 2 tablespoons	
1 tablespoon = 3 teaspoons	

Tank Mixing

To improve the spectrum of activity, TURFLON Ester may be tank mixed at a rate of 1/2 to 1 pint per acre (1/4 to 1/3 fluid ounce in 2 gallons of water per 1,000 square feet) with recommended rates of low volatile amine or ester formulations of 2,4-D, MCPP or other labeled postemergence broadleaf herbicides. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

Broadcast Treatment of Ornamental Turf

Apply 1/2 to 1 quart per acre (1/3 to 3/4 fluid ounce per 1,000 square feet) of TURFLON Ester in enough water to provide uniform coverage of the target area to control actively growing broadleaf weeds growing in perennial bluegrass, perennial ryegrass, or tall fescue. DO NOT use on other turfgrass species (see "General Information" above) unless injury can be tolerated. To minimize turf injury, do not treat if turf is under heat or drought stress and make repeat applications at least 4 weeks apart.

Spot Treatment of Ornamental Turf

Mix 3/8 to 3/4 fluid ounce of TURFLON Ester per 1,000 square feet in enough water to provide uniform coverage of the target area and apply at any time broadleaf weeds are susceptible. **Note:** DO NOT apply more than 2 quarts per acre (1.5 fluid ounces per 1,000 square feet) of TURFLON Ester in a single application.

Control of Kikuyugrass

Apply TURFLON Ester at a rate of 1/2 to 1 quart per acre (1/3 to 3/4 fluid ounce per 1,000 square feet) in enough water to provide uniform coverage of the target area. To improve activity, MSMA herbicide may be tank mixed with the 1/2 quart per acre (1/3 fluid ounce per 1,000 square feet) rate of TURFLON Ester. Three to four additional applications at 4 to 6 week intervals may be required to achieve control of kikuyugrass.

Suppression of Bermudagrass

Apply TURFLON Ester at the rate of 1 quart per acre (3/4 fluid ounce per 1,000 square feet) in enough water to provide uniform coverage of the target area. Three to four additional

applications at 4 week intervals will be required to give adequate suppression of bermudagrass and allow fescue or other desired turfgrass species to dominate.

To improve suppression and control of bermudagrass, 1 quart per acre (3/4 fluid ounce per 1,000 square feet) of TURFLON Ester may be tank mixed with a post-emergence grass herbicide registered for this use pattern. Three to four additional applications of this tank mix at 4 week intervals should be made to achieve control. Reseeding following application will accelerate the transition to cool season turf (see "Reseeding Precaution" above).

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store above 28°F or agitate before use.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of according to applicable federal, state or local procedures.

CONTAINER DISPOSAL: Triple rinse (or equivalent) during mixing and loading. Recycling decontaminated containers is the best option of container disposal. The Agricultural Container Recycling Council (ACRC) operates the national recycling program. To contact your state or local ACRC recycler, visit the ACRC web page at www.acrcycle.org. Decontaminated containers may also be disposed of in a sanitary landfill.

CONDITIONS OF SALE AND WARRANTY

Monterey AgResources (MAR) warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated when used in accordance with the use directions under normal conditions. MAR neither makes, nor authorizes any agent or representative to make, any other warranties, express or implied, including fitness or merchantability.

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and plant conditions, presence of other materials, or other influencing factors, all of which are beyond the control of MAR and Seller. Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage and use of this material not in strict accordance with directions given herewith. In no case shall MAR or the Seller be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by MAR and is accepted as such by the Buyer.

NET CONTENTS:

- 1 QUART (0.95 L)
- 1 GALLON (3.79 L)

Manufactured for:



P.O. Box 35000
Fresno, CA 93745
(559) 499-2100

Controls Annual and Perennial
Broadleaf Weeds and Bermudagrass
in Ornamental Turf

TURFLON* Ester

SPECIALTY HERBICIDE

EPA Reg. No. 17545-8
EPA Est. No. 48498-CA-1

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

ACTIVE INGREDIENT:

Triclopyr: 3,5,6-trichloro-2-pyridinyl-oxyacetic acid, butoxyethyl ester 61.6%

OTHER INGREDIENTS: 38.4%

TOTAL 100.0%

Triclopyr acid equivalent: 44.3% (4 lb/gal).

Contains petroleum distillates.

*Trademark of DowAgroSciences LLC

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing during and after application. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Do not allow people (other than applicator) or pets on treated areas. Do not allow people or pets to enter the treated areas until sprays have dried. Do not apply this product in a way that will contact any person or pet, either directly or through drift.

Local conditions may affect the use of herbicides. Consult your local specialist for advice in selecting treatments from this label to best fit local conditions.

TURFLON Ester may injure certain turfgrass species. DO NOT apply to bahiagrass, bentgrass, bermudagrass, centipedegrass, St. Augustinegrass or zoysiagrass, unless turf injury can be tolerated.

DO NOT apply TURFLON Ester to exposed roots of shallow rooted trees and shrubs.

DO NOT apply TURFLON Ester herbicide directly to or otherwise permit it to come in contact with or permit spray mists containing TURFLON Ester to drift onto cotton, grapes, tobacco, vegetable crops, flowers, fruit or orchard trees, shrubs or other desirable broadleaf plants.

Avoid Injurious Spray Drift: Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. DO NOT spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. Spray drift can be reduced by spraying when wind velocity is low, keeping the spray boom as low as possible, using nozzles that deliver larger spray droplets, and by using higher volume sprays and lower pressures.

Broadleaf weeds controlled by TURFLON Ester include:

- | | |
|---------------------|---------------------|
| Black medic | Lespedeza |
| Bull thistle | Matchweed |
| Burdock | Mustard |
| Canada thistle | Oxalis |
| Chicory | Plantain |
| Clover | Ragweed |
| Creeping beggarweed | Smartweed |
| Curly dock | Sweet clover |
| Dandelion | Vetch |
| Field bindweed | Wild carrot |
| Goldenrod | (Queen Anne's lace) |
| Ground ivy | Wild lettuce |
| Kikuyugrass** | Wild violet |
| Lambsquarters | Yarrow |

**See control of Kikuyugrass under "Application Directions".

USER SAFETY RECOMMENDATIONS:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

GENERAL INFORMATION

(Read Carefully Before Using)

TURFLON Ester herbicide is recommended for the control of actively growing annual and perennial broadleaf weeds and kikuyugrass in perennial bluegrass, perennial ryegrass or tall fescue ornamental turf.

DO NOT apply this product through any type of irrigation system. DO NOT apply to ditches used to transport irrigation water. DO NOT apply where runoff or irrigation water may flow onto susceptible plants

KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a Poison Control Center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

Note to physician: This product may pose an aspiration pneumonia hazard. Contains petroleum distillates.

APPLICATION DIRECTIONS

Foliar sprays should be applied during warm weather, from early spring through fall, when weeds are actively growing. Broadleaf weeds germinate at different times. Only emerged weeds present at the time of application will be controlled. Newly seeded turf should be mowed 2 or 3 times before being treated. When making applications to mature plants, hard to control species, or during drought conditions, use higher rates. Application under drought conditions may provide less than desirable results. Use low pressure sprays to minimize spray drift. DO NOT water for 24 hours after application.

Reseeding Precaution: DO NOT reseed for 3 weeks after application. This precaution does not apply when bermudagrass turf is overseeded with perennial ryegrass at a minimum reseeding rate of 400 lbs. per acre (9 lbs. per 1,000 sq.ft.).

Mixing Instructions

When TURFLON Ester is mixed with water, it forms an emulsion (not a solution) and separation may occur unless the spray mixture is agitated continuously.

Add about one-half the required amount of clean water to the spray tank. Add the recommended amount of TURFLON Ester and continue filling the sprayer. For best results, the spray mixture should be mixed well and often.

Conversion Table

1 ounce = 2 tablespoons
1 tablespoon = 3 teaspoons

Tank Mixing

To improve the spectrum of activity, TURFLON Ester may be tank mixed at a rate of 1/4 to 1/3 fluid ounce in 2 gallons of water per 1,000 square feet with recommended rates of low volatile amine or ester formulations of 2,4-D, MCPP or other labeled postemergence broadleaf herbicides. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

Broadcast Treatment of Ornamental Turf

Apply 1/3 to 3/4 fluid ounce per 1,000 square feet of TURFLON Ester in enough water to provide uniform coverage of the target area to control actively growing broadleaf weeds growing in perennial bluegrass, perennial ryegrass, or tall fescue. DO NOT use on other turfgrass species (see "General Information" above) unless injury can be tolerated. To minimize turf injury, do not treat if turf is under heat or drought stress and make repeat applications at least 4 weeks apart.

Spot Treatment of Ornamental Turf

Mix 3/8 to 3/4 fluid ounce of TURFLON Ester per 1,000 square feet in enough water to provide uniform coverage of the target area and apply at any time broadleaf weeds are susceptible. **Note:** DO NOT apply more than 1.5 fluid ounces per 1,000 square feet of TURFLON Ester in a single application.

Control of Kikuyugrass

Apply TURFLON Ester at a rate of 1/3 to 3/4 fluid ounce per 1,000 square feet in enough water to provide uniform coverage of the target area. To improve activity, MSMA herbicide may be tank mixed with the 1/3 fluid ounce per 1,000 square feet rate of TURFLON Ester. Three to four additional applications at 4 to 6 week intervals may be required to achieve control of kikuyugrass.

Suppression of Bermudagrass

Apply TURFLON Ester at the rate of 3/4 fluid ounce per 1,000 square feet in enough water to provide uniform coverage of the target area. Three to four additional applications at 4 week intervals will be required to give adequate suppression of bermudagrass and allow fescue or other desired turfgrass species to dominate. To improve suppression and control of bermudagrass, 3/4 fluid ounce per 1,000 square feet of TURFLON Ester may be tank mixed with a post-emergence grass herbicide registered for this use pattern. Three to four additional applications of this tank mix at 4 week intervals should be made to achieve control. Reseeding following application will accelerate the transition to cool season turf (see "Reseeding Precaution" above).

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store above 28°F or shake well before use. Store unused product in original container only in a cool, dry area out of reach of children. Keep container tightly closed when not in use. Do not store near heat or open flame.

CONTAINER DISPOSAL: If empty - Do not reuse this container. Place in trash or offer for recycling if available. If Partially Filled - Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONDITIONS OF SALE AND WARRANTY

Monterey AgResources (MAR) warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated when used in accordance with the use directions under normal conditions. MAR neither makes, nor authorizes any agent or representative to make, any other warranties, express or implied, including fitness or merchantability.

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and plant conditions, presence of other materials, or other influencing factors, all of which are beyond the control of MAR and Seller. Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage and use of this material not in strict accordance with directions given herewith. In no case shall MAR or the Seller be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by MAR and is accepted as such by the Buyer.

NET CONTENTS:

- 8 FL.OZ. / 0.5 PINT (0.24 L)
- 1 PINT (0.47 L)

Manufactured for:



P.O. Box 35000
Fresno, CA 93745
(559) 499-2100