



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

MAY 18 2010

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Ms. Lynn P. Georges
Regulatory Affairs Manager
Monterey AgResources
P.O. Box 35000
Fresno, CA 93745

Dear Ms. Georges:

SUBJECT: TURFLON® Ester
EPA Registration No 17545-8
Application and Letter Dated: May 6, 2010,
Request To Amend Registration To Comply with PR
Notice 2007-4 and To Amend Labeling as Described

The proposed subject amendments to the registration of "TURFLON® Ester" have been reviewed and were found acceptable under the Federal Insecticide Fungicide and Rodenticide Act (FIFRA), as amended, provided that you:

1. Revise the ingredient statement to comply with PR Notice 97-5; i.e., replace the chemical expression for triclopyr with the CAS name: [(3,5,6-trichloro-2-pyridinyl)oxy]acetic acid.
2. Revise by deleting the "Container Reuse" heading and comply with Chapter 13 of "Label Review Manual". Add a revised "Container Handling" section. Move "Offer for recycling if available" to last statement, but with the modification: "If recycling is not available, puncture and dispose of in a sanitary landfill or incineration. Do not burn unless allowed by state and local ordinance".
3. On both labels revise the "General" Information" sections to read "Product Information".


2028

-2-

4. Submit printed labeling on one side of each page.
5. Preface all disclaimers of liability in the "Conditions of Sale and Warranty" section with the phrase: "to the extent consistent with applicable law". Note: Prior to: "Buyer and user acknowledge and assume all ", this example.
6. Submit one copy of the final printed labeling prior to shipping this pesticide product under the revised labeling.

If these condition are not complied with, this registration will be subject to cancellation in accordance with FiFRA, section 6(e). Your release for shipment of this product under the revised labeling constitutes acceptance of these conditions. A stamped copy of the accepted labeling is enclosed for your records.

Sincerely yours,


James Tompkins
Product Manager (25)
Herbicide Branch
Registration Division (7505P)

Enclosure

30f8

MASTER LABEL

Turflon[®] Ester Specialty Herbicide

ACCEPTED
with COMMENTS
In EPA Letter Dated:

MAY 18 2010

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

17545-8

**Controls Annual and Perennial Broadleaf Weeds
and Bermudagrass in Ornamental Turf**

Active Ingredient:

Triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid, butoxyethyl ester 61.6%

Other Ingredients: 38.4%

Total: 100.0%

Triclopyr acid equivalent: 44.3% (4 lb/gal).

Contains petroleum distillates.

EPA Reg. No. 17545-8

EPA Est. No. 48498-CA-1

® Trademark of DowAgroSciences LLC

SUBLABEL A - Commercial / Agricultural Uses (Subject to the Worker Protection Standard)


SUBLABEL B - Non-Commercial / Residential Uses (Not Subject to the Worker Protection Standard)

Manufactured for:

Monterey AgResources

P. O. Box 35000 • Fresno, CA 93745-5000 • (559) 499-2100

www.montereyagresources.com



4048

SUBLABEL A: Commercial / Agricultural Uses

Controls Annual and Perennial
Broadleaf Weeds and Bermudagrass
in Ornamental Turf

Turflon® Ester

SPECIALTY HERBICIDE

EPA Reg. No. 17545-8
EPA Est. No. 48498-CA-1

ACTIVE INGREDIENT:

Triclopyr: 3,5,6-trichloro-2-pyridinyl-
oxyacetic acid, butoxyethyl ester. 61.6%

OTHER INGREDIENTS: 38.4%

TOTAL 100.0%

Triclopyr acid equivalent: 44.3% (4 lb/gal).

Contains petroleum distillates.

*Trademark of DowAgroSciences LLC

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

IF SWALLOWED: Immediately call a Poison Control Center or doctor. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

For additional information on this product, you may call the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

Note to physician: Contains petroleum distillate - vomiting may cause aspiration pneumonia.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as nitrile or butyl rubber
- Shoes plus socks
- Protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to sod farm uses.

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as nitrile or butyl rubber
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to all use sites on this label except for sod farm uses.

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry restrictions for Non-WPS Uses: For applications to turf other than sod farms or turfgrass grown for seed, do not allow entry into treated areas until sprays have dried.

GENERAL INFORMATION

(Read Carefully Before Using)

Use Turflon Ester specialty herbicide for the control of actively growing annual and perennial broadleaf weeds and kikuyugrass in perennial bluegrass, perennial ryegrass or tall fescue ornamental turfgrass including these turfgrasses in sod farms and golf courses.

Local conditions may affect the use of herbicides. Consult your local specialist for advice in selecting treatments from this label to best fit local conditions.

General Use Precautions and Restrictions

Chemigation: Do not apply this product through any type of irrigation system.

Do not apply Turflon Ester directly to, or otherwise permit it to come into direct contact with, cotton, grapes, peanuts, soybeans, tobacco, vegetable crops, flowers, citrus, fruit or orchard trees, shrubs, or other desirable broadleaf plants. Do not permit spray mists containing Turflon Ester to drift onto such plants.

To minimize turfgrass injury, do not treat if turfgrass is under heat or drought stress. Reapply Turflon Ester at 4-week-intervals.

Mow newly seeded turfgrass 2 or 3 times before treating. Do not water for 24 hours after application. Do not reseed for 3 weeks after application. (This precaution does not apply when Bermudagrass turfgrass is overseeded with perennial ryegrass at a minimum reseeding rate of 400 lb per acre or 9.2 lb per 1,000 sq.ft.)

- Do not apply to ditches currently being used to transport irrigation water. Do not apply where runoff or irrigation water may flow onto agricultural land as injury to crops may result.
- Do not use on other turfgrass species such as bahiagrass, bentgrass, Bermudagrass, centipedegrass, St. Augustinegrass, or zoysiagrass unless turfgrass injury can be tolerated.
- Do not allow grazing to any area treated with Turflon Ester.
- Do not apply Turflon Ester to exposed roots of shallow rooted trees and shrubs.
- Do not apply to golf course greens.
- Do not aerially apply this product.
- Do not apply more than 8 lb ae triclopyr per acre per year (2 gallons Turflon Ester per acre per year or 5.87 fl.oz. Turflon Ester per 1,000 sq.ft. per year)

Turflon Ester is formulated as a low volatile ester. However, the combination of spray contact with impervious surfaces, such as roads and rocks, and increasing ambient air temperatures, may result in an increase in the volatility potential for this herbicide, increasing a risk for off-target injury to sensitive crops such as grapes and tomatoes.

Avoiding Injurious Spray Drift

Make applications only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants that are near enough to be injured. Spray drift can be reduced by spraying when wind velocity is low, keeping the spray boom as low as possible, using nozzles that deliver larger spray droplets, and by using higher volume sprays and lower pressures.

Broadleaf weeds controlled include:

Black medic	Lespedeza
Bull thistle	Matchweed
Burdock	Mustard
Canada thistle	Oxalis
Chicory	Plantain
Clover	Ragweed
Creeping beggarweed	Smartweed
Curly dock	Sweet clover
Dandelion	Vetch
Field bindweed	Wild carrot
Goldenrod	(Queen Anne's lace)
Ground ivy	Wild lettuce
Kikuyugrass**	Wild violet
Lambsquarters	Yarrow

**See control of Kikuyugrass under "Application Directions".

APPLICATION DIRECTIONS

Foliar sprays should be applied during warm weather, from early spring through fall, when weeds are actively growing. Broadleaf weeds germinate at different times. Only emerged weeds present at the time of application will be controlled. When making

applications to mature plants, hard to control species, or during drought conditions, use higher rates.

Mixing Instructions

Add about one-half the required amount of clean water to the spray tank. Add Turflon Ester and complete addition of water with agitation running. Mix thoroughly and continue moderate agitation while spraying. When Turflon Ester is mixed with water it forms an emulsion, not a solution; separation may occur unless the spray mixture is agitated continuously.

Tank Mixing

To improve the spectrum of activity, Turflon Ester may be tank mixed at a rate of 1/2 to 1 pint per acre (1/4 to 1/3 fl.oz. in 2 gallons of water per 1,000 sq.ft.) with recommended rates of low volatile amine or ester formulations of 2,4-D, MCPP, or other labeled postemergence broadleaf herbicides provided the tank mix product is registered for use on ornamental turfgrass (perennial bluegrass, perennial ryegrass, or tall fescue ornamental turfgrass including these turfgrasses in sod farms and golf courses). Refer to the product labels of tank mix products and follow all appropriate use directions, precautions, and limitations.

Conversion Table

- 1 ounce = 2 tablespoons
- 1 tablespoon = 3 teaspoons

Broadcast Treatment of Ornamental Turfgrass

Apply 1/2 to 1 quart per acre (1/3 to 3/4 fl.oz. per 1,000 sq.ft.) of Turflon Ester in enough water to provide uniform coverage of the target area to control actively growing broadleaf weeds growing in perennial bluegrass, perennial ryegrass, or tall fescue. DO NOT use on other turfgrass species (see "General Information" above) unless injury can be tolerated.

Spot Treatment of Ornamental Turfgrass

Mix 3/8 to 3/4 fl.oz. of Turflon Ester per 1,000 sq.ft. in enough water to provide uniform coverage of the target area and apply at any time broadleaf weeds are susceptible. **Note:** DO NOT apply more than 2 quarts per acre (1.5 fl.oz. per 1,000 sq.ft.) of Turflon Ester in a single application.

Control of Kikuyugrass

Apply Turflon Ester at a rate of 1/2 to 1 quart per acre (1/3 to 3/4 fl.oz. per 1,000 sq.ft.) in enough water to provide uniform coverage of the target area. Three to four additional applications at 4 to 6 week intervals may be required to achieve control of kikuyugrass.

Suppression of Bermudagrass

Apply Turflon Ester at the rate of 1 quart per acre (3/4 fl.oz. per 1,000 sq.ft.) in enough water to provide uniform coverage of the target area. Three to four additional applications at 4 week intervals are required

6008

for adequate suppression of Bermudagrass and allow fescue or other desired turfgrass species to dominate.

To improve suppression and control of Bermudagrass, tank mix 1 quart of Turflon Ester per acre (3/4 fl.oz. per 1,000 sq.ft.) with a postemergence grass herbicide registered for this use pattern. Three to four additional applications of this tank mix at 4-week intervals should be made to achieve control. Reseeding following application will accelerate the transition to cool season turfgrass.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store above 28°F or agitate before use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER REUSE: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure**

rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

CONDITIONS OF SALE AND WARRANTY

Monterey AgResources (MAR) warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated when used in accordance with the use directions under normal conditions. MAR neither makes, nor authorizes any agent or representative to make, any other warranties, express or implied, including fitness or merchantability.

The directions for use of this product are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and plant conditions, presence of other materials, or other influencing factors, all of which are beyond the control of MAR and Seller. Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage and use of this material not in strict accordance with directions given herewith. To the extent consistent with applicable law, neither MAR nor the Seller shall be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by MAR and is accepted as such by the Buyer.

NET CONTENTS:

- 1 QUART (0.95 L)
- 1 GALLON (3.79 L)

Manufactured for:



P.O. Box 35000
Fresno, CA 93745
(559) 499-2100

7098

Controls Annual and Perennial
Broadleaf Weeds and Bermudagrass
in Ornamental Turf

Turflon[®] Ester

SPECIALTY HERBICIDE

EPA Reg. No. 17545-8
EPA Est. No. 48498-CA-1

ACTIVE INGREDIENT:

Triclopyr: 3,5,6-trichloro-2-pyridinyl-oxycetic acid, butoxyethyl ester. 61.6%

OTHER INGREDIENTS:..... 38.4%

TOTAL 100.0%

Triclopyr acid equivalent: 44.3% (4 lb/gal).

Contains petroleum distillates.

®Trademark of DowAgroSciences LLC

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

IF SWALLOWED: Immediately call a Poison Control Center or doctor. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

For additional information on this product, you may call the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information

Note to physician: Contains petroleum distillate - vomiting may cause aspiration pneumonia.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Wear long sleeved shirt and long pants, shoes and socks when mixing or applying this product.

Do not allow people (other than the applicator) or pets on treated areas. Do not allow people or pets to enter the treated areas until sprays have dried. Do not apply this product in a way that will contact any person or pet, either directly or through drift.

USER SAFETY RECOMMENDATIONS:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage system.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

GENERAL INFORMATION

(Read Carefully Before Using)

Use Turflon Ester specialty herbicide for the control of actively growing annual and perennial broadleaf weeds and kikuyugrass in perennial bluegrass, perennial ryegrass or tall fescue ornamental turfgrass.

Local conditions may affect the use of herbicides. Consult your local specialist for advice in selecting treatments from this label to best fit local conditions.

General Use Precautions and Restrictions

Do not apply Turflon Ester directly to, or otherwise permit it to come into direct contact with, cotton, grapes, peanuts, soybeans, tobacco, vegetable crops, flowers, citrus, fruit or orchard trees, shrubs, or other desirable broadleaf plants. Do not permit spray mists containing Turflon Ester to drift onto such plants.

To minimize turfgrass injury, do not treat if turfgrass is under heat or drought stress. Reapply Turflon Ester at 4-week-intervals.

Mow newly seeded turfgrass 2 or 3 times before treating. Do not water for 24 hours after application. Do not reseed for 3 weeks after application. (This precaution does not apply when Bermudagrass turfgrass is overseeded with perennial ryegrass at a minimum reseeding rate of 400 lb per acre or 9.2 lb per 1,000 sq.ft.)

- Do not apply to ditches currently being used to transport irrigation water. Do not apply where runoff or irrigation water may flow onto susceptible plants.
- Do not use on other turfgrass species such as bahiagrass, bentgrass, Bermudagrass, centipedegrass, St. Augustinegrass, or zoysiagrass unless turfgrass injury can be tolerated.
- Do not allow grazing to any area treated with Turflon Ester.
- Do not apply Turflon Ester to exposed roots of shallow rooted trees and shrubs.
- Do not apply more than 8 lb ae triclopyr per acre per year (5.87 fl.oz. Turflon Ester per 1,000 sq.ft. per year)

Turfion Ester is formulated as a low volatile ester. However, the combination of spray contact with impervious surfaces, such as roads and rocks, and increasing ambient air temperatures, may result in an increase in the volatility potential for this herbicide, increasing a risk for off-target injury to sensitive crops such as grapes and tomatoes.

Avoiding Injurious Spray Drift

Make applications only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants that are near enough to be injured. Spray drift can be reduced by spraying when wind velocity is low, keeping the spray boom as low as possible, using nozzles that deliver larger spray droplets, and by using higher volume sprays and lower pressures.

Broadleaf weeds controlled include:

Black medic	Lepedeza
Bull thistle	Matchweed
Burdock	Mustard
Canada thistle	Oxalis
Chicory	Plantain
Clover	Ragweed
Creeping beggarweed	Smartweed
Curly dock	Sweet clover
Dandelion	Vetch
Field bindweed	Wild carrot
Goldenrod	(Queen Anne's lace)
Ground ivy	Wild lettuce
Kikuyugrass**	Wild violet
Lambsquarters	Yarrow

**See control of Kikuyugrass under "Application Directions".

APPLICATION DIRECTIONS

Foliar sprays should be applied during warm weather, from early spring through fall, when weeds are actively growing. Broadleaf weeds germinate at different times. Only emerged weeds present at the time of application will be controlled. When making applications to mature plants, hard to control species, or during drought conditions, use higher rates.

Mixing Instructions

Add about one-half the required amount of clean water to the spray tank. Add Turfion Ester and complete addition of water with agitation running. Mix thoroughly and continue moderate agitation while spraying. When Turfion Ester is mixed with water it forms an emulsion, not a solution; separation may occur unless the spray mixture is agitated continuously.

Tank Mixing

To improve the spectrum of activity, Turfion Ester may be tank mixed at a rate of 1/4 to 1/3 fl.oz. in 2 gallons of water per 1,000 sq.ft. with recommended rates of low volatile amine or ester formulations of 2,4-D, MCP, or other labeled postemergence broadleaf herbicides provided the tank mix product is registered for use on ornamental turfgrass (perennial bluegrass, perennial ryegrass, or tall fescue ornamental turfgrass). Refer to the product labels of tank mix products and follow all appropriate use directions, precautions, and limitations.

Do not apply this product through any type of irrigation system.

Conversion Table	
1 ounce = 2 tablespoons	
1 tablespoon = 3 teaspoons	

Broadcast Treatment of Ornamental Turfgrass

Apply 1/3 to 3/4 fl.oz. per 1,000 sq.ft. of Turfion Ester in enough water to provide uniform coverage of the target area to control actively growing broadleaf weeds growing in perennial bluegrass, perennial ryegrass, or tall fescue. DO NOT use on other turfgrass species (see "General Information" above) unless injury can be tolerated.

Spot Treatment of Ornamental Turfgrass

Mix 3/8 to 3/4 fl.oz. of Turfion Ester per 1,000 sq.ft. in enough water to provide uniform coverage of the target area and apply at any time broadleaf weeds are susceptible. **Note:** DO NOT apply more than 1.5 fl.oz. per 1,000 sq.ft. of Turfion Ester in a single application.

Control of Kikuyugrass

Apply Turfion Ester at a rate of 1/3 to 3/4 fl.oz. per 1,000 sq.ft. in enough water to provide uniform coverage of the target area. Three to four additional applications at 4 to 6 week intervals may be required to achieve control of kikuyugrass.

Suppression of Bermudagrass

Apply Turfion Ester at the rate of 3/4 fl.oz. per 1,000 sq.ft. in enough water to provide uniform coverage of the target area. Three to four additional applications at 4 week intervals are required for adequate suppression of bermudagrass and allow fescue or other desired turfgrass species to dominate.

To improve suppression and control of Bermudagrass, tank mix 3/4 fl.oz. of Turfion Ester per 1,000 sq.ft. with a postemergence grass herbicide registered for this use pattern. Three to four additional applications of this tank mix at 4-week intervals should be made to achieve control. Reseeding following application will accelerate the transition to cool season turfgrass.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container only. Keep in a cool, dry place inaccessible to children or animals, preferably in a locked storage area.

CONTAINER DISPOSAL: If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONDITIONS OF SALE AND WARRANTY

Monterey AgResources (MAR) warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated when used in accordance with the use directions under normal conditions. MAR neither makes, nor authorizes any agent or representative to make, any other warranties, express or implied, including fitness or merchantability.

The directions for use of this product are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and plant conditions, presence of other materials, or other influencing factors, all of which are beyond the control of MAR and Seller. Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage and use of this material not in strict accordance with directions given herewith. To the extent consistent with applicable law, neither MAR nor the Seller shall be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by MAR and is accepted as such by the Buyer.

NET CONTENTS:

- 8 FL.OZ. (0.24 L)
- 1 PINT (0.47 L)
- 1 QUART (0.95 L)
- 1 GALLON (3.79 L)

Manufactured for:



P.O. Box 35000
Fresno, CA 93745
(559) 499-2100