

DEAN'S INSECTICIDAL MINERAL

Use as a fly repellent during the summer months. Use as a feed supplement for mixing in a ration for control of lice and ticks. Use as a feed supplement for control of cattle lice for a 14-day control program for 100% control of insect cattle and non-dairy animals only.

See cow feeding directions carefully.

Pesticially Active Ingredient

Resmethal-O, O-dimethyl O-2-Ethyl-5-chlorophenyl phosphorothioate 6.0%

Inert Ingredients 94.0%

GUARANTEED ANALYSIS

Salt (NaCl), not more than 30.00%
Salt (NaCl), not less than 28.00%
Calcium (Ca), not more than 15.50%
Calcium (Ca), not less than 13.50%
Phosphorus (P), not less than 8.50%

INGREDIENTS:

Dicalcium Phosphate, Calcium Deformed Phosphate, Calcium Carbonate, Salt, Manganese Oxide, Iron Carbonate, Zinc Oxide, Copper Oxide, Ethylene Diamine Dihydrochloride, Cobalt Carbonate, Magnesium Oxide, Iron Oxide, Vitamin A Palmitate (Stability Improved), D-Activated Animal Sterol (source of Vitamin D3), dl-Alpha Tocopheryl Acetate, Cane Molasses, Natural and Artificial Flavors added.

WARNINGS

Use as a sole source of Resmethal.

Dairy cows should be treated during their dry period but **only** at the dry period (cows between the end of the fall breeding season and cows about to reach the barn).

Do not feed to dairy cows that will produce milk for human consumption. Do not use prior to freshening. If dairy cow or heifer is on a normal feeding program, use of insecticidal mineral should not be withdrawn the required 10-day period. All animal milk must not be used for food for humans during the last feeding of this product.

Withdraw for 10 days prior to slaughter.

Do not feed to sick, injured, or stressed animals, such as those that are clipped, castrated, or weaned within the last 4 weeks.

Do not use while pregnant or nursing. Do not use this product in animals seen to be pregnant within a few days of freshening. Do not use in animals seen to be nursing. If human health is a problem, consult your physician. For more information, consult your veterinarian. For more information, consult your veterinarian.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Caution: This product contains Resmethal-O, a highly toxic insecticide. It is extremely poisonous to humans and animals. It is also extremely poisonous to fish and aquatic life. Avoid contact with humans and animals.

Do not use in areas where it may be a hazard to humans or animals. Do not use in areas where it may be a hazard to fish and aquatic life. Do not use in areas where it may be a hazard to birds and other wildlife.

Do not use in areas where it may be a hazard to humans or animals. Do not use in areas where it may be a hazard to fish and aquatic life. Do not use in areas where it may be a hazard to birds and other wildlife.

Do not use in areas where it may be a hazard to humans or animals.

Do not use in areas where it may be a hazard to humans or animals.

Manufactured By
DEAN'S SPECIALTY SUPPLY CO.

Worcester, Massachusetts 01693

NET WEIGHT 1 LB.

**DIRECTIONS
FOR CONTINUOUS SELF FEEDING PROGRAM
DEAN'S INSECTICIDAL MINERAL**

When to feed: Feed Dean's Insecticidal Mineral continuously to all cattle in the herd. Make certain that the mineral is available at all times. Feed continuously in all weather conditions. Do not stop feeding continuously for more than 75 days.

How Much to Feed: Provide a feed concentration of Dean's Insecticidal Mineral containing 0.6% Rotenone in the proportion of 1 lb. weight per head per month.

How to Feed: Feed Dean's Insecticidal Mineral continuously free choice to cattle in covered mineral feeder troughs or boxes. Provide one mineral feeder box or trough space for each 15 to 20 head of cattle. Locate mineral feeder boxes and troughs near watering areas. To insure uniform and normal consumption, put out only one day supply at one time, and keep feeder boxes and troughs clean. For adequate insect control, each animal must eat its share of mineral. Small, late calves may require an additional 7- or 14-day feeding program after weaning for grub control.

WARNING

Do not combine this feed with any other feed containing Rotenone. Do not mix Dean's Insecticidal Mineral with any other feed containing salt or mineral.

**DIRECTIONS
FOR 7-DAY OR 14-DAY FEEDING PROGRAM**

Mix Dean's Insecticidal Mineral with a rough grain ration of 100 lbs. of mix with the exception of 10 lbs. of insecticidal feed. Feed continuously for 7 or 14 days and make certain that all cattle in the herd have access to the feed. Feed should be available to all cattle in the herd. Make certain that insecticidal feed is available to all cattle in the herd. Provide mineral feed for Dean's Insecticidal Mineral for each 100 lbs. of live weight of cattle for the 7- or 14-day feeding period.

NOTE: Insecticidal feed should be available to all cattle in the herd. Mineral should be available for 4 weeks of time.

When to Feed: After a long, light frost has set in, when the grass reaches the rack. For grub control treatment, feed for 7 or 14 days. Later stages of grub control should be fed to all cattle. Rotate the cattle in the feed troughs and direct to the 7- or 14-day feeding program.

7-DAY TREATMENT

When to Feed:

Southern Area: All cattle in the Southern States
Northern Area: All cattle in the Northern States

How to Feed: Mix this feed with a rough grain ration of 100 lbs. of mix with the exception of 10 lbs. of insecticidal feed. Feed continuously for 7 or 14 days and make certain that all cattle in the herd have access to the feed. Feed should be available to all cattle in the herd. Make certain that insecticidal feed is available to all cattle in the herd. Provide mineral feed for Dean's Insecticidal Mineral for each 100 lbs. of live weight of cattle for the 7- or 14-day feeding period.

14-DAY TREATMENT

When to Feed:

Southern Area: All cattle in the Southern States
Northern Area: All cattle in the Northern States

How to Feed: Mix this feed with a rough grain ration of 100 lbs. of mix with the exception of 10 lbs. of insecticidal feed. Feed continuously for 14 days and make certain that all cattle in the herd have access to the feed. Feed should be available to all cattle in the herd. Make certain that insecticidal feed is available to all cattle in the herd. Provide mineral feed for Dean's Insecticidal Mineral for each 100 lbs. of live weight of cattle for the 14-day feeding period.

