

PM 1

17975-1

17/1

DEAN'S MEAT BASE

Rat and Mouse Bait \$5.55

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. **NOTE TO PHYSICIAN** — If ingested, administer Vitamin K1 intramuscularly or orally, as indicated in bishydroxycoumarin overdoses Repeat as necessary based on monitoring of prothromin times.

ENVIRONMENTAL HAZARD:

Keep Out of Lakes, Streams or Ponds.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS — For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS —

Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

DEAN'S MEAT BASE Rat and Mouse Bait

Kills Norway Rats, Roof Rats and House Mice.

ACTIVE INGREDIENT(S):

Warfarin (3-(Alpha - acetylbenzyl) - hydroxycoumarin)	0.025%
N1 - (2 - Quinoxaliny)l) sufanilamide (Sulfaquinoxaline)	0.025%
Inert Ingredients	99.950%
TOTAL	100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS:

EPA Reg. No. 17975-1

EPA Est. No. 17975-1A-01

NET CONTENTS: 3 1/2 POUNDS

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days. **MICE:** Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS & MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

DISPOSAL

DO NOT REUSE EMPTY CONTAINER. WRAP CONTAINER AND PUT IN TRASH COLLECTION.

JERRY DEAN EXTEMINATOR

307 EAST 8TH STREET
SPENCER, IOWA 51301

ACCEPTED

JAN 21 1996

BEST AVAILABLE COPY

17975-1