

INSECTICIDAL 10-T CATTLE MINERAL

MEDICATED

Summer for Control of Hornflies and Cattle Grubs; for Mixing in Feed to Provide a 7-Day Feeding Program for Grub Control and as an Aid in the Reductation of Cattle Lice, or a 14-Day Feeding Program for Crub Control; for beef cattle and non-lactating dairy mimals only.

Follow feeling directions carefully.

GUARANTEED ANALYSIS

Calcium (Ca), not more than .				7.00%
Calcium (Ca), not less than .				6.00°
Phosphorus (P), not less than .				10.00°°
ludine (1), not less than				0.025%
Salt (NaCl), not more than				29.00 °₀
Sult (NaCl), not less than				26.00%
INGREDIEN	TS			

i Scium Phosphate, Monocalcium Phosphate, Ialta Millian Care Molasses, Steumed Bone Meal, Diatomaceous Earth, Resident Chiac, Iron Sulfate Tinc Okide Manganese Itate, Magnesium Chiae, Copper Chiae, Ethylomoviamine Dinydric dide, Copalt Carbonate, Minste, Cil, Clyceryl Monostearate, Pryodenic Tiloca, Inches Meal, Artificial Elavor Added.

CAUTION: KEEP OUT OF REACH OF CHILDREN

WARN'INGS USE as a sole source of Ronnel.

Dairy cows should be treated early in their dry period but only if the dry period occurs between the end of the adult heel fly season and before grobs reach the back.

Do not feed to dairy animals that will produce milk for human consumption for 10 days prior to freshening. If dairy cows or heifers freshen during feeding program, or if insecticidal mineral has not been withdrawn the required 10 days prior to freshening, milk must not be used for food for 10 days after the last feeding of this product.

withdraw 10 days prior to slaughter.

Do not feed to sick onimals or animals under stress, such as those just shipped, dehorned, castrated, or weaned within the last 3 weeks.

Ronnel is a cholinesterase inhibitor. Do not use this product in animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides or chemicals. Consult a veterinarian at the first sign of adverse reaction. Atropine is an antidote.

Wash hands with soap and water after handling. This material contains a cholinesterase inhibitor. Treat Symptomatically. Atropine is an antidote. Harmful if swallowed.

This product is toxic to tish. Keep out of lakes, streams, and ponds. Do not contaminate by cleaning of equipment or disposal of wastes. Use this product only as specified on this label. Do not reuse empty containers. Destroy it by burning or burying in non-crop lands away from water supplies. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

STORE IN A COCL, DRY PLACE NET WT. 30 LBS.

1025

SEECO, Inc.
Box 1014

WILLMAR, MINNESOTA 56201

EPA Reg. No. 18037 - 2

COMINUOUS SELF-FEEDING THE LET TO SECTICIDAL CATTLE THE FRANCIO-T MEDICATED

thedis a period insecticidal Cattle Mineral state of the state of the

The Feed: Recommended consumption of the Air ideal Cattle Mineral containing 6% and 100.2 lb. per 100 lb. of body weight per that per month.

now to Feed: Feed Seeco Insecticidal Cattle Minral ontinously on a self-fed basis to cattle in
more a mineral-feeder troughs or baxes. Provide
as mineral feeder box or feeding boxspace for each
with Colinead of cattle. Locate mineral-feeder
exister strongramear watering areas. To encourage
mineral and normal consumption, put out only a 7my supply at one time, and keep feeder boxes and
who obtain. For adequate insect control, each
a rail must eat its share of mineral. Small, late
all es most require an additional 7 or 14 day feedmineral strongster meaning for control.

MARNING

teeding program, do not allow cate to be a consecticidal Cattle Mineral and additional salt or mineral.

FLEERING PROGRAMS

ressing, or mix with the complete secticisal feed. Feed conting days and make certain that all Adequate feeding space should the conting of animal. Make certain that



insecticidal feed is evenly distributed on, or mixed into, the total ration. Provide a total of 0.2 lb. of Seeco Insecticidal Cattle Mineral for each 100 lb. body weight (to be fed over the 7-or 14 day feeding period).

NOTE: Insecticidal feed made from Seeco Insecticidal Cattle Mineral should be used within 4 weeks of mixing.

When to Feed: After end of additived fly season and before grubs reach the back. Formel is not effective for grub control during later stages of life cycle, when grubs are in the backs of cattle. Reduction of cattle lice will be obtained when fed as directed for the 7-day program.

7-DAY TREATMENT

When to Feed:

Southern area cattle - July through September Northern area cattle - August through October How to Feed: Mix thoroughly 2 lb. of Seeco Insecticidal Cattle Mineral containing 6% Ronnel with 19 lb. feed to produce an insecticidal feed. Prepare only enough for the 7 day feeding program. Feed 0.3 lb. of insecticidal feed per 100 lb. body weigh daily for 7 continous days.

14-DAY TREATMENT

When to Feed:

Southern area cattle - July through September.
Northern area cattle - August through DECEMBER.
How to Feed: Mix thoroughly 1 lb. of Seeco Insecticidal Cattle Mineral containing 6 is Ronnel with 20 lb. of feed to produce an insecticidal feed. Prepare only enough for the 14-day feeding program. Feed 0.3 lb. of insecticidal feed per 100 lb. body weight daily for 14 continuous days.

