

**USE DIRECTIONS
FOR CLEANING**

Lightly soiled areas — surfaces to be cleaned should be mopped or brushed with a solution containing 3 ounces of SAN-I-TERGE per gallon of water. Rinse cleaned area with fresh water.

Heavily soiled areas — mop or brush soiled surface with a solution containing 5-6 ounces of SAN-I-TERGE per gallon of water. Rinse cleaned surface with fresh water.

GENERAL PURPOSE DISINFECTATION

To disinfect walls, floors and other hard surfaces in homes, schools, restaurants, rest rooms and other public areas, mop or brush surface to be treated with a solution containing 3½ ounces of SAN-I-TERGE per gallon of water.

Surfaces that are heavily soiled should be cleaned with 2 ounces of SAN-I-TERGE per gallon of water and then disinfected. Those surfaces which may later come in contact with food must be rinsed with clear water of drinking quality.

To disinfect locker rooms, shower stalls, walkways around pools and other areas where fungicidal action is required, mop area with a solution containing 2 ounces in 1 gallon of water. This solution will destroy *Trichophyton interdigitale*, a causative agent of athlete's foot.

HOSPITAL DISINFECTION

To disinfect walls, floors, furniture and other hard surfaces in hospitals and nursing homes, mop or brush surface to be treated with a solution containing 5½ ounces of SAN-I-TERGE per gallon of water. Those surfaces that are heavily soiled with blood, fecal material or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water and then rinsed with a fresh solution containing 2 ounces per gallon of water before disinfection.

SANITIZATION

To sanitize walls, floors and equipment in food processing plants, first clean thoroughly with a solution containing 3 ounces of SAN-I-TERGE per gallon of water. If surface is heavily soiled, use SAN-I-TERGE at a rate of

SAN-I-TERGE

**DISINFECTANT—CLEANER
SANITIZER—FUNGICIDE—DEODORANT**

ACTIVE INGREDIENTS

Sodium metasilicate 3.0%
n-Alkyl (50% C₁₄, 40% C₁₂, 10% C₆) dimethyl benzyl ammonium chloride . . . 1.5%

INERT INGREDIENTS

. 95.5%

EPA Reg. No. 18305-2

ACCEPTED
JUL 9 1973

DANGER

KEEP OUT OF REACH OF CHILDREN. See antidote statement and other precautions on side panel.

Container Disposal: When empty, rinse container thoroughly with water and destroy it by crushing or burying in a safe place.

UNDER FEDERAL INSECTICIDE
FUNGICIDE AND FUNGICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 18305-2

**MANUFACTURED BY
SYNDET PRODUCTS INC.
P. O. Box 1425
Manchester, Conn. 06040**

**NET CONTENTS
1 GALLON**

5-6 ounces per gallon to insure thorough cleaning. Sanitize cleaned surface with a rinse solution containing 2 ounces of SAN-I-TERGE per gallon of water.

Apply a portable water rinse prior to use of equipment or utensils.

DEODORIZATION

To deodorize surfaces during cleaning, mop or brush surfaces with a solution containing 7 ounces of SAN-I-TERGE per gallon of water.

SAN-I-TERGE is an effective antibacterial cleaner designed for use in household, industrial and institutional cleaning. When used as directed SAN-I-TERGE will disinfect hard surfaces, such as floors, walls, counter tops and furniture. It may be relied upon to deodorize bathroom areas, garbage pails, food storage areas, kennel areas and other areas where obnoxious odors may develop. This product may be used as a general purpose antibacterial detergent in homes, schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other places where efficient cleaning and antimicrobial action is desired.

See left panel for use directions.

**DANGER
KEEP OUT OF REACH
OF CHILDREN**

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.