

OTC-10

ACCEPTED

COMPOSITION

Percent by Weight

Active Ingredient . . . . .	15.0
Poly (oxyethylene (dimethyliminio) ethylene (dimethyliminio) ethylene dichloride). . . . .	15.0
Inert Ingredient . . . . .	85.0
Weight per gallon . . . . .	8.58 pounds
Weight of active ingredient per gallon . . . . .	1.29 pounds

APPLICATION

OTC-10 is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algae growth, microbiological slime and other deposits. An initial addition of 4.0 to 10.0 fluid ounces of OTC-10 per 1000 gallons of water to provide a concentration of 32 to 80 parts per million of OTC-10 based on the total weight of water in the system is recommended. Repeat initial dosage until control is evident. Subsequent slug additions of 1.0 to 10.0 fluid ounces of OTC-10 per 1000 gallons of water will produce 8 to 80 parts per million of OTC-10. OTC-10 should be applied every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleed-off and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

CAUTION!

KEEP OFF THE REACH OF CHILDREN.

Harmful if swallowed. Avoid contact with skin and eyes. In case of contact flush with plenty of water for at least 15 minutes. If eye irritation persists get medical attention. Avoid contamination of food.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public waters. DO NOT contaminate water by cleaning or equipment or disposal of water. Use only as directed on label. Do not reuse container. Do not burn empty.