

#### ALPHA-G

A ready to use, dual purpose formulation for control of seed-borne & soil-borne diseases and for temporary protection against wireworms. A fungicide-insecticide combination seed treatment for agricultural use only

WHEAT-BARLEY -OATS -FLAX-CORN- SOYBEANS- RYE

# KEEP OUT OF REACH OF CHILDREN WARNING

Additional Warning Statements on Back

ALPHA-G protects crops against seed-borne diseases such as stinking smut of wheat,oat smut,covered and false loose smut of barley, and bunt of rye. It also gives temporary protection to cereal grains.corn,and soybeans against soil-borne diseases such as root rot and seedling blight. Lindane temporarily protects seed and seedlings against wireworms.

DIRECTIONS FOR USE Thoroughly mix ALPHA-G with seed in the planter box or with a mechanical dust treater. DO NOT  $M^{\intercal}X$  WITH HANDS

WHEAT-BARLEY-OATS-RYE: Use rate of 2 Oz. per bushel FLAX: Use rate of 4 Oz. per bushel CORN-SOYBEANS: Use rate of 2.75 Oz. per bushel For maximum benefits, use well cured, dry and clean seed at the recommended rate and avoid deep seeding and early sowing under poor field conditions.

EPA REG. No. 19382-6 EPA EST. No. 19382-ND-1 NET WEIGHT 10#

Manufactured by ALPHA RESEARCH CHEMICAL CO. Grand Forks, N.D.

White Committee is a second of the committee of the commi

registered under EPA Beg. Na.

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

May be fatal if inhaled, swallowed, or adsorbed through the skin--therefore avoid contact with this product. Do not breathed dust, and wear a mask over mouth and nose while handling. In case of contact, flush with plenty of water, and for eyes get medical attention. Wash with soap and water after handling and before eating.

Applicators who apply this product manually or without the use of a closed system treatment must wear long-sleeved shirt; long pants; gloves; and a disposable paper dust mask which covers at least one-third of the face as protective equipment. Treat outdoors or in a well ventilated place only.

#### STATEMENT OF PRACTICAL TREATMENT

- If swallowed-Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. If possible, vomiting should be induced under medical supervision. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
- If inhaled-Remove victim to fresh air. Apply respiration if needed.
- If on skin-Remove contaminated clothing and wash with soap and water.
- If in eyes-Flush eyes for at least 15 minutes with water. Call a physician if irritation persists.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label .Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all excess treated seeds by burial away from bodies of water

It is a violation of State and Federal Law to use this product in a manner inconsistent with its labeling. Read all directions before using and use only as specified or this label

## STORAGE AND DISPOSAL

- 1. PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.
- 2. PESTICIDE DISPOSAL: Pesticide that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away frow water supplies.
- 3. CONTAINER DISPOSAL: Dispose in an incinerator or landfill approved for pesticide containers.
- 4. GENERAL: Consult Federal, State or Local Disposal Authorities for approved alternative procedures such as limited open burning.

#### USE RESTRICTIONS

Treat only seeds needed for immediate planting and minimize the interval between treatment and planting. Treated seed must not be used for or mixed with food or animal feed or processed for oil. Do not store excess treated seeds beyond planting time. Do not graze or feed livestock on treated areas for six weeks after planting. Care must be exercised in handling of treated seed.

## WARRANTY

Seller warrants that this product conforms to the Ingredient Statement on this label and is reasonably fit for the purpose set forth in the Directions For Use.

Manufactured by
ALPHA RESEARCH CHEMICAL CO.
Grand Forks, N.D.

1

The state of the s

- 3. At the most open to see the following of the "L. Switter contains to present the following of the object with the first of the object of the following of the object of the following of the object of the obj
- be low bety defet to the condition, regrainly begin be induced under reduced as envision? from the "Li swellowed" so der. This statement usually around only on high to delity produces that also contain a high ownershape of lager petrologic solvent that could cause aspiration processons as a complication resulting from inducing vocating.
- is the first the net and y directly to Taken, streams or money to the not apply directly to eater. "
- . Siving the Destinate Disposal text to read:

Jectical exactor are texic. Teprosex disposal of excess posturary, sprey menture, or resate is a violation of local leve. If these visites council to dispose the local leve is a concern of the local local level of local levels of lever to the local levels of local levels of lever to the local levels of lever to levels of levels local levels of levels of levels local levels of levels of levels of levels of levels levels of leve

The Control of the Co

with the second contract that the property of the second contract of

The first transfer to the foreign, which is the first of the atom, on the control of the state o

transfer to the state of

Mar Will have been a

. ..