

TEAS - ORANGES AND CITRUS For control of the citrus thrips which feed to the detriment of the fruit, apply a systemic application of the spray containing Dursban 4E to the leaves and fruit.

CORN (FIELD) - Post Emergence Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to corn less than 4 weeks after emergence. Do not apply to corn less than 4 weeks after emergence. Do not apply to corn less than 4 weeks after emergence.

Pre-emergence - Arkansas, Louisiana, Mississippi and Tennessee Make a single application of 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to corn less than 4 weeks after emergence.

COTTON Pre-plant - Florida and California Only. Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to cotton less than 4 weeks after emergence.

Pre-emergence - Except Arizona and California Make a single application of 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to cotton less than 4 weeks after emergence.

Soil Type*	Soybean	
	Rate (lb. Dursban 4E)	Rate (lb. Dursban 4E)
Light sand	0.6	1.5
Sandy loam, loam, silty loam, and silt	1.0	1.5
Clay loam, silty loam, silty clay loam, and clay	1.25	1.5
Silt loam and silt	2.0	1.6

*Soil types listed are for general reference. For more detailed information, consult the Soil Conservation Service Manual.

BROADCAST TREATMENT Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to corn less than 4 weeks after emergence.

POST-EMERGENCE TREATMENT Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to corn less than 4 weeks after emergence.

Need Problem	Dursban 4E Herbicide	Rescue Dursban
Pre-plant	1.5 to 2.0	1.5 to 2.0
Post-plant	1.5 to 2.0	1.5 to 2.0

Late Season (May 6) Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to corn less than 4 weeks after emergence.

Subsequent Crops Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to corn less than 4 weeks after emergence.

Dursban 4E Herbicide	Type of Application	Crops That May Follow Treated Cotton
Pre-plant	Broadcast	Any crop
Post-plant	Broadcast	Cotton, soybean, grain sorghum, rice, sugarcane, and other crops
Pre-emergence	Broadcast	Cotton, soybean, grain sorghum, rice, sugarcane, and other crops
Post-emergence	Broadcast	Cotton, soybean, grain sorghum, rice, sugarcane, and other crops

NOTE Do not apply to corn less than 4 weeks after emergence.

GRAPES Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to grapes less than 4 weeks after emergence.

NOTES Do not apply to corn less than 4 weeks after emergence.

East of the Rocky Mountains Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to corn less than 4 weeks after emergence.

New York and Pennsylvania - Adoption of Federal Grades Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to corn less than 4 weeks after emergence.

West of the Rocky Mountains Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to corn less than 4 weeks after emergence.

GRASS SEED CROPS Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to grass seed crops less than 4 weeks after emergence.

Colors of the Soil Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to corn less than 4 weeks after emergence.

Western Oregon Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to corn less than 4 weeks after emergence.

NOTE Do not apply to corn less than 4 weeks after emergence.

GRAPES Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to grapes less than 4 weeks after emergence.

PREPPLE Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to corn less than 4 weeks after emergence.

SUGARCANE Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to sugarcane less than 4 weeks after emergence.

NOTES Do not apply to corn less than 4 weeks after emergence.

GRAPES Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to grapes less than 4 weeks after emergence.

NOTES Do not apply to corn less than 4 weeks after emergence.

GRAPES Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to grapes less than 4 weeks after emergence.

NOTES Do not apply to corn less than 4 weeks after emergence.

GRAPES Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to grapes less than 4 weeks after emergence.

NOTES Do not apply to corn less than 4 weeks after emergence.

GRAPES Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to grapes less than 4 weeks after emergence.

NOTES Do not apply to corn less than 4 weeks after emergence.

GRAPES Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to grapes less than 4 weeks after emergence.

NOTES Do not apply to corn less than 4 weeks after emergence.

GRAPES Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to grapes less than 4 weeks after emergence.

NOTES Do not apply to corn less than 4 weeks after emergence.

GRAPES Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to grapes less than 4 weeks after emergence.

NOTES Do not apply to corn less than 4 weeks after emergence.

GRAPES Apply 1.5 to 2.0 lbs. per acre as a broadcast application to the soil in the rows. Do not apply to grapes less than 4 weeks after emergence.

NOTES Do not apply to corn less than 4 weeks after emergence.



EPA Reg. No. 19713-21
EPA Est. No. 19713-MS-1
Net Contents lbs.