

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: HARMFUL IF SWALLOWED. Avoid Breathing of Spray. Do Not Take Internally. Avoid Contact with Skin and Eyes. Causes moderate eye irritation. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash thoroughly after handling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting because of aspiration hazard. Call a physician or Poison Control Center.

IF IN EYES: Immediately flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Immediately remove contaminated clothing, INCLUDING SHOES, and wash skin with plenty of soap and plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. This product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of waste.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

GENERAL INFORMATION

DAMQIL is an especially refined Phytotoxic Oil Contact Insecticide. It is for use on plants in foliage or the dormant stage. When using this product in combination with other pesticides as directed on this label, observe appropriate cautions, limitations for the products involved.

Do not combine Sulfur sprays with DAMQIL on foliage and do not follow a previous application of Sulfur or apply Sulfur following DAMQIL, until sufficient time has elapsed for safety (at least 30 days). If in doubt, spray a portion of one tree previously treated with Sulfur, using 1 quart to 25 gallons water. If sufficient time has not elapsed, burn will show in several days. Do not use DAMQIL when temperatures are above 90°F. When temperatures drop below 32°F (usually nighttime lows), do not spray tree and vine crops until daytime temperatures have warmed to a point above 32°F and are free from moisture due to melting frost or heavy dews. Do not use on foliage closer than 30 days after or 90 days before using Captan or Folpet products.

NOTE: This restriction does not apply to uses during the dormant period. Do not use in combination with SEVIN on crops other than citrus. Application to trees weakened by disease, drought, drying winds, or high nitrogen applications may result in injury.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Amounts Given Are For 100 Gallons of Diluted Spray.

Alfalfa (Seed Crop Only): Use 1 quart per acre in 20 to 30 gallons of water by ground application or 10 to 15 gallons by air when used in combination and at the recommended label rates with either a Malathion 5 Emulsive or a Toxaphene 8 Emulsive for control of the pests listed on the product labels. Spider Mites: Use 1 quart per acre at the dilutions noted above with Kelthane EC at the recommended label rates.

Almonds, Apricots, Cherries, Nectarines, Peaches, Plums, Prunes—Dormant or Delayed Dormant: Brown Apricot Scale, Black Scale, San Jose Scale, Olive Scale (Parlatoria Scale), Oleander Scale, Frosted Scale, Italian Pear Scale, European Fruit Lecanium, Putnam Scale, Parlatoria, Soft Brown Scale, Certain Other Scale Insects, Eggs of Aphids, Red Spider Mites—1½ to 2 gals. Summer Use—Red Spider Mites—3 to 4 qts. Summer Stages of Brown Apricot Scale, San Jose Scale—1 gal. Apples: Rust Mite—Delayed Dormant—1½ to 2 gals. Summer Use—1 to 1½ gals.

Apples, Pears—Dormant or Delayed Dormant: San Jose Scale, Italian Pear Scale, Eggs of Aphids, Red Spider Mites—1½ to 2 gals. Summer Use: Red Spider Mites—3 to 4 qts. (Concentrated Sprays)—Pears (Call only)—4 to 6 gals. in 90 to 120 gals. of water per acre. Summer Forms of San Jose and Italian Pear Scale—1 to 1½ gals. **Figs:** Fig Scale—Dormant or Delayed Dormant—Use 3 gals. DAMQIL Spray. Severe infestation—Use 2 gals. DAMQIL plus ½ pint of Phosdrin 4 Pound Emulsive.

Grapes: Mealybug—Dormant—Use 4 to 6 gals. plus 2 to 2½ qts. of ORTHOPHOS 4 Emulsive in 200 to 300 gals. of water per acre. Do not exceed 2½ qts. of ORTHOPHOS 4 Emulsive per acre. **Lemons, Grapefruit:** Red Scale, Yellow Scale, Purple Scale, Black Scale, Red Spider Mites—1 to 1½ gals. DAMQIL plus 1 lb. of a Malathion 25 Wettable

FOR WESTERN AGRICULTURE

Olive: Parlatoria Scale, Spider Mites, Oleander Scale, Black Scale—Use 1½ gals. C.L.200L combine 1 gal. D.A.C.L. with ½ pint of a Phosdrin 4 Pound Emulsive per 100 gals. water. Apply in the pre-bloom or in post-bloom up to bud set.

Nectar Orange: Fig Scale, Black Scale, Yellow Scale, Purple Scale, Red Spider Mites—3 qts. D-M-01, plus 1 lb. of a Malathion 25 Wettable. Do not use on Navel Oranges after Sept. 1.

Valencia Orange: Red Scale, Black Scale, Yellow Scale, Purple Scale, Red Spider Mites—1 gal. applying not later than Sept. 1 or 3 qts. DAMQIL plus 1 lb. of a Malathion 25 Wettable, applied not later than Sept. 15.

Pears: Pear Psylla—Delayed Dormant—2 gals. Summer Use: 1 to 1½ gals. Bester Mite—Delayed Dormant—1 or 1½ gals. per 100 gals.

Ornamentals and Shade Trees (Cedars) (see Note), Holly, Pyracantha, Sycamore, Hawthorne, Flowering Peach, Flowering Cherry, Apple, Magnolia, Maple, Dogwood, Oak, Privet, Birch: Soft Brown Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale, San Jose Scale, Red Spider Mites and Eggs—1 gal. to 100 gals. water. Dormant Spray on Deciduous Varieties—2 gals. per 100 gals. water. Dormant Spray on Other Varieties—1 gal. per 100 gals. water.

NOTE: May cause discoloration of blue spruce. Do not use on conifers or ferns not tolerant to oil sprays.

Walnuts: Late Spring to Mid-Summer Only. Summer Stages of Frosted Scale, European Fruit Scale, San Jose and other Soft-Shelled Scale, Red Spider Mites—2 to 4 quarts. The addition of one of the following propionate insecticides aids in scale control: A Trithion 25 Wettable, a Phosdrin 4 Pound Emulsive, an Ethion 25 Wettable, or a Parathion 25 Wettable. See label for recommended dosages. Do not apply after hard set.

Mixing Directions: With agitator running start filling spray tank with water. When combining with Wettable Powder Insecticide or fungicide products, add these first when tank is ¼ filled. Add DAMQIL when tank is about ½ full and while there is sufficient agitation of the spray to insure complete emulsification.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment.

STORAGE INSTRUCTIONS

Storage should be under lock and key in a ventilated room and secure from access by unauthorized persons and children. Storage should be in a cool, dry area away from any heat or ignition source. Do not stack over 2 pallets high. Move containers by hand or crane. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed, and fertilizer to avoid contamination. Store in original containers only if the contents are leaking or material is spilled follow these steps:

1. Contain spill, absorb with a material such as saw dust, clay granules or dirt.
2. Collect and place in suitable containers for disposal.
3. Wash area with water and soap to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Place a leaking container in a plastic tub and transfer contents as soon as possible to an empty original container.
6. Do not allow run off to enter sewer or contaminate water supplies.
7. Dispose of wastes as indicated below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY—CONDITION OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon best believed reliable, follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. In no case shall Oxeel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or

0%
0%
0%

ONS
-123
MS-1