



ACCEPTED

JUL 7 1981

Federal Insecticide and Rodenticide Act Under the Pesticide Label

CARBARYL 5-D

A Dusting Powder for Control of Insect Pest

**KEEP OUT OF THE REACH OF CHILDREN
CAUTION**

ACTIVE INGREDIENT: Carbaryl (1-naphthylmethylcarbamate) 5%
INERT INGREDIENT: 95%

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

HARMFUL IF SWALLOWED. Avoid contact with eyes. Avoid contact with clothing. Avoid contact with skin. Wash hands and face before eating. Wash thoroughly after handling.
NOTE: Carbaryl 5-D is a moderate to severe irritant to the eyes and skin. Avoid contact with eyes and skin. Wash hands and face before eating. Wash thoroughly after handling.

Statement of Practical Treatment

IF SWALLOWED: - Induce vomiting by drinking water or milk. Do not induce vomiting if the person is unconscious. Never give anything by mouth to an unconscious person.
IF IN EYES: - Immediately flush eyes with plenty of water for at least 15 minutes. If irritation develops, get medical attention.
IF ON SKIN: - Remove clothing. Wash skin with soap and plenty of water. If irritation develops, get medical attention.

Environmental Hazards

Avoid contact with water. Do not feed water to livestock. Do not apply when weather conditions favor drift from area treated.
BEE CAUTION: - May kill honeybees. Avoid contact with bees. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Extension Service. Do not use when the value of honeybees as pollinators is high. Do not use when insect control is not required. Do not use when bees are present in or near treated areas. Do not use when bees are present in or near treated areas.

GENERAL CAUTIONS

1. To avoid possible injury, do not use if you are allergic to any of the ingredients or if you are pregnant or expect to become pregnant during the next 30 days.
2. Do not use if you are ill.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION AND LIMITATIONS

Carbaryl 5-D is toxic to several species of bees. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

DIRECTIONS FOR USE

Apply to crops as directed. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

VEGETABLES

Beans: - Apply to beans as directed. Do not use on beans that are important bee forage plants. Do not use on beans that are important bee forage plants. Do not use on beans that are important bee forage plants.

Cabbage, Broccoli, Brussels Sprout, Cauliflower, Kohlrabi: - To control Flea Beetles, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

Collards, Garden Beets, Kale, Lettuce, Mustard Greens, Radishes, Rutabagas, Spinach, Turnips: - To control Flea Beetles, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

Corn (Sweet Field): - To control European Corn Earworm, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

Cucumber, Squash, Lantelope, Pumpkin, Melons: - To control Flea Beetles, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

Carrot: - To control Flea Beetles, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

Tomato, Eggplant, Peppers: - To control Flea Beetles, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

Potato: - To control Colorado Potato Beetle, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

SMALL FRUIT

Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries: - To control Japanese Beetle, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

Strawberries: - To control Flea Beetles, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

ORNAMENTALS

To control Flea Beetles, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

COTTON

To control Flea Beetles, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

PASTURE, FORAGE GRASSES, CLOVER, ALFALFA

To control Flea Beetles, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

HOUSEHOLD PETS

Dogs, Cats: - To control Flea Beetles, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

Sleeping Quarters (Dog, Cat)

To control Flea Beetles, use 20-40 pounds per acre. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

STORAGE AND DISPOSAL

Do not contaminate water. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

WARRANTY

The manufacturer warrants that the product is as labeled. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants. Do not use on crops that are important bee forage plants.

Net Contents



Drexel Chemical Company