



Drexel

# Toxaphene 8E

## Crop and Livestock Spray Concentrate

**ACTIVE INGREDIENTS:**  
 Toxaphene  
 (Technical Chlorinated Camphene) 71.19%  
 Xylene range aromatic solvent 23.48%

**INERT INGREDIENTS:** 5.33%  
 \*Containing 67% to 69% chlorine  
 Contains 8 lbs. Technical Toxaphene per gallon

ACCEPTED  
 JUL 1 1981  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act  
 19713 55

### Statement of Practical Treatment

**IF SWALLOWED:** Call a physician or Poison Control Center. Do not induce vomiting. See "Physician's Note." Never give anything by mouth to an unconscious person.

**IF IN EYES:** Wash eyes for at least 15 minutes, while holding eyelids open. Call a physician.

**IF ON SKIN:** Remove contaminated clothing and immediately wash the skin with soap and water.

**CALL A PHYSICIAN OR POISON CONTROL CENTER**

**PHYSICIAN'S NOTE:** If swallowed, use gastric lavage. Avoid aspiration. Toxaphene produces central nervous system excitation that

responds to barbituric acid derivatives. Prevent or control convulsions with barbiturates preferably administered intravenously.

If the barbiturate can be given before convulsions start, sodium pentobarbital is the drug of choice. Since toxaphene has a fairly long duration of action, a long-acting barbiturate such as phenobarbital may be used after initial control of convulsions. Refer to "Clinical Handbook of Economic Poisons," which was prepared by the U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE, PUBLIC HEALTH SERVICE COMMUNICABLE DISEASE CENTER, Atlanta, Georgia.

See Side Panel For Additional Precautionary statements.

Keep Out Of The Reach Of Children

### WARNING



Drexel Chemical Company Memphis, Tennessee

Net Contents

Gallons  
Liters

EPA Reg. No. 19713-55  
EPA Est. No. 19713-MS-1

# PRECAUTIONARY STATEMENTS

## HAZARD TO HUMANS AND DOMESTIC ANIMALS

**WARNING:** May be fatal if swallowed. Poisonous by skin contact and inhalation. Do not get on skin, in eyes, or on clothing. Do not breathe spray mist. Wash thoroughly after using. Wash all contaminated clothing with soap and hot water before reuse. During application, keep unprotected persons away from areas being treated or where there may be drift. Do not contaminate feed or foodstuffs.

**ENVIRONMENTAL HAZARD:** Avoid contamination of food, feed, water supplies, streams and ponds during application or when cleaning equipment. Do not apply when weather conditions favor drift from areas treated.

This product is toxic to fish, shrimp, crab, birds, and other wildlife. Keep out of any body of water. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. This product is toxic to bees and should not be applied when bees are actively visiting the area.

## PHYSICAL OR CHEMICAL HAZARDS

**Do Not Use, Pour, or Store Near Heat or Open Flame.** (In case of spillage use absorbent clay, hydrated lime, or other solids to soak up any liquid. Wash area with strong detergent and hot water.)

## DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is to be diluted in water for ground equipment application on the listed crops. The amount of water required per acre will depend upon the type of equipment used. This varies from 5 to 100 gallons of water per acre.

### DO NOT USE ON CUCURBITS

**ALFALFA Restrictions:** For Seed Production Only. Do not feed straw and threshings to livestock. If applied during bloom, do so only when bees and other pollinators are not visiting plants.

Insect	Pints/Acre
Alfalfa caterpillar	2.5 - 3
Alfalfa leathopper	1.5 - 3
Alfalfa looper	2 - 3.5
Alfalfa weevil	2 - 3.5
Armyworm (including true, fall, and yellow-striped)	2 - 3.5
Clover leaf weevil	2
Corn earworm	2.25
Cutworm (including climbing and surface-feeding)	2 - 3
Grasshopper	2 - 3.5
Lygus bug	1.5 - 2
Pea aphid	3
Spittlebug	1.5
Sweetclover weevil	1.5
Thrips	1.5 - 2
Webworm	2.25 - 3

**BANANAS Restrictions:** Repeat applications as necessary but do not apply less than one day before harvest. Do not graze treated area to dairy animals or animals being finished for slaughter.

Insect	dosage
Leaf-eating caterpillar	1/2 pt/gal of water
Ceratitis	1/2 pt/gal of water
Carpenter fly	1/2 pt/gal of water
Osmia	1/2 pt/gal of water
Spine	1/2 pt/gal of water

**BEANS (Green, Snap, and Lima) Restrictions:** Sprays not to be applied to shelled beans as a pre-harvest treatment. Do not apply sprays to green snap beans after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter.

If the above restrictions apply to the crop, use the following rates for the listed insects. Do not apply more than 3 applications after the first application and harvest 21 days or less after the last application. Do not apply within 30 days of harvest.

Insect	Pints/Acre
Armyworm	1 - 2
Bean beetles (including Mexican bean leaf beetle, cucumber beetle, wax bean beetle, blister beetle, flea beetle)	2 - 4
Bean leathopper	1
Bean leaf roller	1
Bean leaf skeletonizer	3 - 4
Bean Lycaenid	2.5 - 3.5
Corn earworm	1 - 2
Cowpea curculio	1 - 4
Cutworm (including climbing and surface-feeding)	2
Darkling ground beetle	3 - 4
Garden webworm	3 - 4
Grasshopper	2 - 3
Lygus bug	2 - 4
Pea weevil	3 - 4
Salt-marsh caterpillar	1 - 4
Serpentine leaf miner	2 - 3
Thrips	1.2 - 1.6

**BROCCOLI Restrictions:** Do not apply after edible parts begin to form or within 30 days of harvest.

Insect	Pints/Acre
Armyworm	1.2 - 1.6
Cabbage aphid	1.2 - 1.6
Cabbage looper	1.5 - 4
Corn earworm	1.5 - 4
Cutworm (including climbing and surface-feeding)	1.2 - 1.6
Diamondback moth	1.5 - 4
Fall armyworm	1.2 - 1.6
Flea beetle	3
Green peach aphid	1.5 - 2
Imported cabbageworm	2
Serpentine leaf miner	2
Thrips	2

**BRUSSELS SPROUTS Restrictions:** Do not apply after edible parts begin to form or within 30 days of harvest.

Insect	Pints/Acre
Armyworm	2 - 4
Cabbage aphid	1.2 - 1.6
Cabbage looper	1.5 - 4
Corn earworm	1.5 - 4
Cutworm (including climbing and surface-feeding)	1.2 - 1.6
Diamondback moth	1.5 - 4
Fall armyworm	1.2 - 1.6
Flea beetle	3
Green peach aphid	1.5 - 2
Imported cabbageworm	2
Serpentine leaf miner	2
Thrips	2

**CABBAGE Restrictions:** Do not apply after head starts to form. Do not feed treated parts to dairy animals or animals being finished for slaughter.

Insect	Pints/Acre
Armyworm	2
Cabbage aphid	1.2 - 1.6
Cabbage looper	2 - 3
Corn Earworm	1.5 - 4
Cutworm (including climbing and surface-feeding)	1.2 - 1.6
Diamondback moth	2 - 3
Fall Armyworm	1.2 - 1.6
Flea beetle	3
Green peach aphid	1.5 - 2
Imported cabbageworm	2 - 3
Serpentine leaf miner	2
Thrips	2

**CARROTS Restrictions:** No time limitations. Apply as a foliar treatment only. Do not feed treated foliage to dairy animals or animals being finished for slaughter.

Insect	Pints/Acre
Cutworm (including black cutworms)	2 - 3
Flea beetle	3
Grasshopper	2
Lygus Bug (seed crop)	3.5 - 4

**CAULIFLOWER Restrictions:** Do not apply after edible parts start to form or within 30 days of harvest.

Insect	Pints/Acre
Armyworm	2
Cabbage aphid	1.2 - 1.6
Cabbage looper	1.5 - 4
Corn earworm	1.5 - 4
Cutworm (including climbing and surface-feeding)	1.2 - 1.6
Diamondback moth	1.5 - 4
Fall armyworm	1.2 - 1.6
Flea beetle	3
Green peach aphid	1.5 - 2
Imported cabbageworm	2
Serpentine leaf miner	2

**CELERY Restrictions:** Do not apply after plants start to bunch or after plants are half mature.

Insect	Pints/Acre
Cabbage aphid	1.2 - 1.6
Celery leaf tier	4 - 5
Cutworm (including climbing and surface-feeding)	1.2 - 1.6
Fall armyworm	1.2 - 1.6
Green peach aphid	1.2 - 1.6
Imported cabbageworm	2
Serpentine leaf miner	1.2 - 1.6
Thrips	1.2 - 1.6
Vegetable weevil	4 - 5

**CITRUS (nonbearing) Restrictions:** Do not feed forage to dairy animals or animals being finished for slaughter.

Insect	Pints/Acre
Cutworm	6
Grasshopper	4

**COLLARDS Restrictions:** Do not make more than a total of 4 applications nor more than 3 applications after the first application. Minimum interval between applications is 7 days. Over a 21-day period.

Insect	Pints/Acre
Armyworm	2 - 4
Cabbage aphid	1.2 - 1.6
Cabbage looper	1.5 - 4
Corn earworm	1.5 - 4
Cutworm (including climbing and surface-feeding)	1.2 - 1.6
Diamondback moth	1.5 - 4
Fall armyworm	1.2 - 1.6

Flea beetle	3
Green peach aphid	1.5 - 2
Imported cabbageworm	2
Serpentine leaf miner	2
Thrips	2

**CORN Restrictions:** Do not feed treated forage to dairy animals or animals being finished for slaughter. No limitation on use of grain.

Insect	Pints/Acre
Armyworm	2.5
Corn billbug	2
Chinch bug	2 - 3
Corn lantern fly	3 - 4
Cutworm	2 - 3
Flea Beetle	2
Grasshopper	2 - 3
Lesser cornstalk borer	4
Webworm	2 - 2.5

**CORN (Preplant or Preemergence Only) Armyworms, Cutworms—Use 2 1/2 to 3 pints of TOXAPHENE BE suitably diluted with water per acre on a broadcast basis. Do not use as a postemergence treatment. Do not graze dairy animals on stover. Do not graze meat animals on stover within 4 weeks of slaughter. No limitation on the use of grain.**

**COTTON Restrictions:** Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season. Do not apply more than a 4 pint/acre rate after buds open.

Insect	Pints/Acre
Armyworm (including true, fall, beet, and yellow striped)	2 - 4
Beet weevil	2 - 4
Beetle	2 - 4
Cotton leaf perforator	2 - 4
Cotton leafworm	2 - 4
Cutworms	2 - 4
Flea beetles	2 - 4
Fleshopper	1 - 3
Garden webworm	3 - 4
Grasshopper	2 - 4
Plant bug (including tarnished and rapid plant bugs, lygus bugs, and superb plant bugs)	2 - 4
Salt-marsh caterpillar	3 - 5
Stink bugs	2 - 4
Thrips	1.5 - 2
Yellow-striped armyworm	2.5

**COWPEA Restrictions:** No limitation on use of shelled peas as human food. Do not apply to cowpeas to be used as green snaps after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter.

Insect	Pints/Acre
Bean leathopper	1
Bean leaf roller	1
Cowpea curculio	1.5 - 3
Cutworm	2 - 3
Darkling ground beetle	3
Lygus bug	2 - 3
Southern green stink bug	2 - 3

**CRANBERRIES Restrictions:** Do not apply after fruit starts to form.

Insect	Pints/Acre
Armyworm	2.5 - 4
Grasshopper	2.5 - 3

**EGG PLANT Restrictions:** Do not apply within 5 days of harvest.

Insect	Pints/Acre
Armyworm	1.2 - 3
Blister beetle	2 - 3
Colorado potato beetle	1 - 1.5
Cucumber beetle	3
Cutworm (including climbing and surface-feeding)	2 - 3
Flea beetle	1.2 - 3
Leathopper	1.2 - 3
Serpentine leaf miner	2 - 3

**GRAIN SORGHUM Restrictions:** Do not apply more than once after heads have begun to form. Do not apply within 28 days of harvest when using up to 2 pints/acre. Do not apply within 40 days of harvest when using over 2 pints/acre. Do not feed treated forage to dairy animals or animals being finished for slaughter. Do not ensile treated forage.

Insect	Pints/Acre
Armyworm	2.5
Corn billbug	2
Chinch bug	2 - 3
Corn lantern fly	3
Cutworm	2 - 3
Flea beetle	2
Grasshopper	2 - 3
Lesser cornstalk borer	3
Webworm	2 - 2.5

**GRAINS: BARLEY, OATS, RICE, RYE, WHEAT Restrictions:** Do not feed treated forage to dairy animals or animals being finished for slaughter. No restrictions on the use of the grain.

Insect	Pints/Acre
Armyworm	2 1/2 - 3
Cutworm (including climbing and surface-feeding)	2 1/2 - 3
Grasshopper	2 1/2 - 3
Chinch bug	3
Rye stink bug	3

**GRAPEFRUIT (see CITRUS) GRASS Restrictions:** For use along highways, power line rights-of-way, and other nonagricultural land.

Insect	Pints/100 gal
Armyworm (including true and fall)	2 - 3
Chiggers	1 - 2
Cutworm (including climbing and surface-feeding)	2
Grasshopper	1.5 - 2
Webworm (including sod and garden webworms)	2 - 3

**HICKORY NUTS Restrictions:** Do not graze livestock in treated groves. No time limitation.

Insect	Pints/100 gal
Pecan weevil	2 - 2.4

**HORSERADISH Restrictions:** Apply as a foliar treatment only. Do not use treated tops as food or feed.

Insect	Pints/Acre
Grasshopper	2 - 3
Striped flea beetle	1.5

**KALE Restrictions:** Do not make more than a total of 4 applications nor more than 3 applications after the first application. Where 2 pints/acre or less is applied, use a 35-day preharvest interval. Where rates above 2 pints/acre are used, use a 30-day preharvest interval.

Insect	Pints/Acre
Armyworm	2
Cabbage aphid	12 - 16
Cabbage looper	15 - 4
Corn earworm	15 - 4
Cutworm (including climbing and surface feeding)	2
Diamondback moth	15 - 4
Fall armyworm	12 - 16
Flea beetle	3
Green peach aphid	15 - 2
Imported cabbage worm	2
Serpentine leaf miner	2
Thrips	2

**KOHLRABI**  
Restrictions:  
Do not apply after edible parts begin to form or within 30 days of harvest

Insect	Pints/Acre
Armyworm	20
Cabbage aphid	12 - 16
Cabbage looper	15 - 4
Corn earworm	15 - 4
Cutworm (including climbing and surface feeding)	12 - 16
Diamondback moth	15 - 4
Fall armyworm	12 - 16
Flea beetle	3
Green peach aphid	15 - 2
Imported cabbage worm	2
Serpentine leaf miner	2
Thrips	2

**LETTUCE**  
Restrictions:  
Do not apply after seeding stage on leaf lettuce. Do not apply after heads begin to form on head lettuce

Insect	Pints/Acre
Armyworm	2
Cabbage looper	3 - 4
Cucumber beetle	15 - 2
Cutworm	2 - 4
Field cricket	2 - 4
Flea beetle	15 - 3
Grasshopper	2 - 4
Imported cabbage worm	15
Salt-marsh caterpillar	3 - 5

**OKRA**  
Restrictions:  
Do not use after pods begin to form

Insect	Pints/Acre
Serpentine leaf miner	2 - 3

**ONIONS**  
Restrictions:  
Do not apply to green or spring onions

Insect	Pints/Acre
Onion thrips	2

**ORNAMENTALS**  
Restrictions:  
See specific crop for restrictions that apply when food crop is taken from ornamentals such as flowers and shrubs

Insect	Pints/100 gals.
Bagworm	2 - 3
Boxelder bug	3
Cankerworm	2 - 3
Catnip worm	3
Emerald ash borer	4
Fall armyworm	4
Fall webworm	4 - 3
Giant cutworm	15 - 2
May apple sawfly	15 - 2
May apple sawfly	15 - 2
Peach weevil	15 - 2
Tent caterpillar	15 - 2
Worm chrysalis	15 - 2

**PARSNIPS**  
Restrictions:  
Apply after harvest

Insect	Pints/Acre
Grasshopper	15
Striped flea beetle	15

**PEAS**  
Restrictions:  
Do not apply after pods begin to form if pods are to be used as food or within 7 days of harvest if pods are not to be used as food. Do not feed vines to dairy animals or animals being finished for slaughter

Insect	Pints/Acre
Bean leafhopper	1
Bean leaf roller	1
Cowpea curculio	15 - 3
Cutworm	2 - 4
Darkling ground beetle	3 - 4
Lygus bug	2 - 3
Southern green stink bug	2 - 4

**PEACHES**  
Restrictions:  
Do not apply within 40 days of harvest or after second cover spray

Insect	Pints/100 gal.
Grasshopper	2 - 3
Plum curculio	7 - 15

**PEANUTS**  
Restrictions:  
Do not feed treated forage to dairy animals or animals being finished for slaughter

Insect	Pints/Acre
Corn earworm	3 - 4
Cutworm	15 - 4
Fall armyworm	2 - 4
Flea hopper	4
Potato leafhopper	2 - 4
Red-necked peanutworm	4
Southern green stink bug	4 - 5
Thrips	2
Velvetbean caterpillar	3

**PEARS**  
Restrictions:  
Do not apply within 40 days of harvest or after second cover spray

Insect	Pints/100 gal.
Pear psylla	1 - 12
Red-banded leaf roller	12

**PECANS**  
Restrictions:  
No time limitations

Insect	Pints/100 gal.
Peanut nut casebearer	1
Pecan weevil	2 - 2 1/2

**PEPPERS**  
Restrictions:  
Do not apply within 5 days of harvest

Insect	Pints/Acre
Armyworm	2 - 3
B. stier beetle	2 - 3
Colorado potato beetle	2 - 3
Flea beetle	12 - 3
Fruitworm	3
Hornworm	2 - 3
Pepper weevil	3
Serpentine leaf miner	2 - 3

**PIMENTOS**  
Restrictions:  
Do not apply within 2 days of harvest

Insect	Pints/Acre
Flea beetles	1 - 4
Hornworm	1 - 4
Cutworms	1 - 4
Pepper weevils	1 - 4

**PINEAPPLE**  
Restrictions:  
Apply after harvest

Insect	Pints/Acre
Pineapple guinea worm	220
(Batrachedra sp.) (Lep. thripa)	220

**QUINCE**  
Restrictions:  
Do not apply within 40 days after harvest or after second cover spray

Insect	Pints/100 gal.
Eastern tent caterpillar	15
Grasshopper	2
Red-banded leaf roller	1 - 12

**RUTABAGAS**  
Restrictions:  
Apply as a foliar treatment only. Do not use treated tops for food or feed

Insect	Pints/Acre
Cutworm	2
Flea beetles	25 - 4
Imported cabbage worm	25 - 4
Spinach leaf miner	25 - 4
Sugarbeet webworm	25 - 4

**SOYBEANS**  
Restrictions:  
May be applied with aerial or ground sprayers. With aerial application the product should be applied in 3 to 4 gallons water per acre. Avoid spray drift. Do not apply less than 3 weeks before harvest. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry. Do not apply within 21 days of harvest

Insect	Pints/Acre
Armyworm	225 - 4
Bean leaf beetle	225 - 4
B. stier beetle	225 - 4
Bollworm	225 - 4
Corn earworm	225 - 4
Cricket	2
Flea beetle	15
Grasshopper	25 - 4
Green cloverworm	225 - 4
Thrips	1 - 15
Velvetbean caterpillar	225 - 4

**STRAWBERRIES**  
Restrictions:  
Do not apply after fruit starts to form

Insect	Pints/Acre
Cutworm	2 - 3
Strawberry crown borer	2 - 3
Strawberry weevil	2 - 3

**SUNFLOWERS**  
Restrictions:  
Apply the specified rate in 5 to 15 gallons of water per acre by ground equipment or aircraft. Do not apply after yellow ray petals have formed on the flower heads. Do not ensile or graze livestock on treated sunflower plants. Do not apply more than 2 times per growing season

Insect	Pints/Acre
Sunflower beetle	15 - 2
Painted lady (the butterfly)	15 - 2
Caterpillars	15 - 2
Cutworms	2
Grasshoppers	2

**SWEET CORN**  
Restrictions:  
Use Restricted Use formulation for the fresh market. Formulate with methyl parathion. Note that methyl parathion is a Restricted Use Pesticide

**Restrictions:**  
Do not apply within 2 days of harvest. Do not use treated tops for 48 hours after application. Do not feed treated tops to dairy animals or animals being finished for slaughter. Do not use treated tops for animal bedding.

**Ground Application (For insects in silks or ears)**  
Apply at 2 day intervals or as needed until fully diluted with water

Insect	Pints Toxaphene/Acre	Pounds Methyl Parathion/Acre
Earworms (Corn ear worm, Fall armyworm)	1 1/2	1/2

**Aerial Application (For insects in whorl or bud)**  
Apply twice a week or as needed until the plant has tasseled. Apply in 3 to 5 gallons of water per acre

Insect	Pints Toxaphene/Acre	Pounds Methyl Parathion/Acre
Budworms (Fall armyworm, corn earworm)	1 1/2	1/2

**Aerial application (For insects in silks and ears)**  
Apply at 1 to 3 day intervals from the time silks appear until the silks have dried. Use 3 to 5 gallons of water per acre

Insect	Pints Toxaphene/Acre	Pounds Methyl Parathion/Acre
Earworms (Corn earworm or fall armyworm)	2	1 1/2

**TOMATOES**  
Restrictions:  
Do not apply within 1 day of harvest at 2 pint/acre rate or within 3 days of harvest at rates over 2 pts/acre

Insect	Pints/Acre
Armyworm	2 - 5
B. stier beetle	2 - 5
Cabbage looper	2 - 3
Colorado potato beetle	2 - 5
Cutworm (including climbing and surface feeding)	2 - 3
Flea beetle	2 - 3
Grasshopper	25 - 4
Green chrysanthemum aphid	2 - 5
Hornworms	2 - 5
Pea worm	2 - 5
Russell mite	2 - 5
Serpentine leaf miner	2 - 5
Thrips	3
Tomato fruitworm	2 - 5

**DIRECTIONS FOR USE ON ANIMALS**  
For controlling horn flies, cattle lice, ticks and sarcoptic mange on beef cattle and controlling sheeplice, sheep scab and goat lice on sheep and goats  
Mixing rate: Dilute with water and use as a dip or spray. For a 0.5% toxaphene water spray, mix 2 1/2 quarts Toxaphene Conc. in 100 gallons water. Mix in 33 gallons water 1 pint in 18 gallons water or 1 cup (6 oz) in 6 gallons. For controlling horn flies, cattle lice, ticks, lone star tick, winter tick, cattle fever tick, Gulf coast tick, sp nose ear tick on beef cattle and

sheep lice, sheep ticks, ticks, goat lice, sheep or goats.

**Dip or Spray:** When dipping make sure the animals held in vat at least two minutes and submerge each animal twice during dipping. For sheep tick apply during warm weather soon after shearing or before wool is 2 or 3 months long.

When spraying use about 2 quarts of the solution per beef animal and proportionately less for smaller animals. If ticks are not controlled repeat application in 3 weeks for cattle. NOTE: Do not apply to dairy animals or small animals under 6 months of age. Do not treat meat animals within 28 days of slaughter.

**HOGS**  
Lice—Spray sows 30 days before farrowing to prevent pigs from being infested. Do not treat pigs under 6 months of age. Do not treat meat hogs within 28 days of slaughter. Do not treat sows within two weeks of farrowing nor for 4 weeks thereafter.

**SARCOPTIC MANGE AND SHEEP SCAB**  
(Beef cattle—sarcoptic mange, sheep and goats—sheep scab) Dip or spray cattle, sheep and goats, for thorough animal coverage. Apply to herd when mange is found. Repeat treatment in 15 to 21 days if necessary. NOTE: Do not apply to dairy animals or small animals under 6 months of age. Do not treat meat animals within 28 days of slaughter.

**BACKRUBBER USE (BEEF CATTLE ONLY)**  
To control hornflies and aid in reducing lice dilute 1/2 gallon Toxaphene Concentrate with 11 gallons kerosene, fuel oil, or diesel oil. Apply this solution to canvas or burlap cable at the rate of one gallon to 20 feet of cable. Retreat at 2 to 3 week intervals or as necessary to keep cable dampened. Locate backrubber to give beef cattle and hogs free access to it. Best locations are usually near watering places, salt licks, or feed racks. NOTE: Do not allow calves under 6 months of age, or dairy animals to treat themselves. Do not permit meat animals access to this treatment within 28 days of slaughter.

**WARRANTY—CONDITION OF SALE:**  
DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of Drexel Chemical Company, or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result.

**ALL SUCH RISKS ARE ASSUMED BY THE BUYER.**  
Drexel warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use as modified by the above. Drexel makes no other warranties, express or implied, including FITNESS or MERCHANTABILITY. In no case shall Drexel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by Drexel Chemical Company and is accepted as such by the Buyer.

**STORAGE AND DISPOSAL**

**Do not contaminate water, food or feed by storage or disposal.**

**Pesticide Disposal:** Pesticide spray mixture or residue that cannot be used according to label instructions must be disposed of according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

**Container Disposal:** Triple rinse (or equivalent) and offer for recycling or reconditioning, or burn in a certified incinerator, or by other approved State and local procedures.