

19713-60

6-30-08

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Luz Chan
Drexel Chemical Company
1700 Channel Avenue
P.O. Box 13327
Memphis, TN 38113-0327

6-30-08

Subject: EPA Reg. 19713-60 Drexel Simazine 4L Supplemental Label Amendment

Dear Ms. Chan:

The label referred to above is acceptable. If you have any questions please contact Erik Kraft at 703-308-9358 or Kraft.Erik@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Jim Tompkins", written over a circular stamp or mark.

Jim Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)


SUPPLEMENTAL LABELING

2/3



Simazine 4L Herbicide

(EPA Reg. No. 19713-60)



ACCEPTED
with **COMMENTS**
In EPA Letter Dated:
6-30-08

FOR USE AS HERBICIDE IN TURF

READ THE ENTIRE DREXEL SIMAZINE 4L LABEL BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABEL

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION

19713-60

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This labeling and the EPA registered labeling must be in the possession of the user at time of pesticide application.

All applicable directions, restrictions and use precautions on Drexel SIMAZINE 4L label(s) are to be followed. This labeling contains supplemental instructions and restrictions for use of this product which may not appear on the Drexel SIMAZINE 4L label. Follow the instructions carefully.

TURFGRASSES FOR SOD (FL Only)

Centipedegrass, St. Augustinegrass, Zoysiagrass: Apply 2 to 4 quarts per acre, according to soil texture as indicated below.

Muck or Peat	4 qts.	Old beds	Within 2 days after lifting of sod
		New beds	3 to 4 days after sprigging or plugging
Sandy soil	2 qts.	Old beds	Within 2 days after lifting sod
		New beds	7 to 10 days after sprigging or plugging

Note: On newly sprigged turfgrass, temporary slowing of growth may follow application.

If weeds regrow, apply an additional 2 quarts of this product on muck or peat, or 1 quart on sandy soil.

Use Precautions and Restrictions:

- Do not apply more than 4 quarts of this product per acre per application. Maximum number of applications per calendar year is two. Maximum total amount per calendar year is 6 quarts of this product per acre.
- Do not apply within 30 days prior to cutting or lifting.
- Do not graze or feed turf clippings to animals.
- Do not apply in combination with surfactants or other spray additives.
- Product must be watered-in immediately after application if applied at a rate greater than 2 qts. per acre.
- Use only on turfgrass reasonably free of infestations of insects, nematodes and diseases.

TURFGRASSES FOR SOD (In States Except FL)

For use on turfgrasses for Sod in all States except Florida, refer to the following section on "TURFGRASSES FOR FAIRWAYS, ORNAMENTAL LAWNS, AND SIMILAR AREAS" for directions for use.

For use directions on turfgrass for Sod in Florida only, refer to the above section on "TURFGRASSES FOR SOD (FL Only)" on this label.

TURFGRASSES FOR FAIRWAYS, ORNAMENTAL LAWNS, AND SIMILAR AREAS

Bermudagrass, Centipedegrass, St. Augustinegrass, Zoysiagrass: Apply this product after September 1 (after October 1 for annual Bluegrass) before emergence of winter annual weeds. Fall applications of this product will control annual Bluegrass, Burclover, common and Mouseear chickweed, Corn speedwell, Henbit, Hop clover, Lawn burweed, Paisley-piert, and Spurweed. This product will also control these weeds soon after emergence.

For control of summer annual weeds listed in the "General Information" section of the main label, apply this product in late winter before the weeds emerge. Apply in a minimum of 15 gallons of water per acre. Irrigate with one-half inch of water if rainfall does not occur within 10 days after preemergence treatment. Where annual Bluegrass is the major weed, use 1 quart of this product per acre (0.75 fl. ozs. per 1,000 sq. ft.). Use 1 to 2 quarts per acre (0.75 to 1.5 fl. ozs. per 1,000 sq. ft.) for control of other weeds. However, do not exceed 1 quart per acre per treatment on newly sprigged Turfgrass or on hybrid Bermudagrass such as Tiflawn, Tifway, and Ormond.

For control of summer annuals which emerge after the initial application, apply an additional 1 quart per acre at least 30 days after the initial application, but not after June 1. However, do not make more than 2 applications per year.

Note: On newly sprigged turfgrass, hybrid Bermudagrass, nondormant Bermudagrass or nondormant Zoysiagrass, temporary slowing or growth and yellowing may occur following application.

Use Precautions and Restrictions:

- Do not apply more than 2 quarts of this product per acre per application. Maximum number of applications per calendar year is two. Maximum total amount per calendar year is 3 quarts of this product per acre.
- Do not apply within 30 days prior to cutting or lifting.
- Do not graze or feed turf clippings to animals.
- Do not use on golf greens.
- Use only on turfgrass reasonably free of infestations of insects, nematodes and diseases.

3/3

- 6) Do not use north of North Carolina (except where used in the VA coastal) or on soils with pH above 7.8.
- 7) Do not use on muck or alkaline soils.
- 8) Do not apply this product to newly seeded grasses until they have overwintered and have a well developed rhizome system.
- 9) Do not apply over the rooting area of trees or ornamentals not listed on the Drexel Simazine 4L label(s).
- 10) Do not seed or overseed with desirable turfgrass within 4 months before or 6 months after treatment.
- 11) Do not exceed 2 quarts of this product per acre within 12 months of seeding of grasses.

FOR HOMEOWNER USE ON TURFGRASSES

Bermudagrass, Centipedegrass, St. Augustinegrass, Zoysiagrass: Apply this product after September 1 (after October 1 for annual Bluegrass) before emergence of winter annual weeds. Fall applications of this product will control annual Bluegrass, Burclover, common and Mouseear chickweed, Corn speedwell, Henbit, Hop clover, Lawn burweed, Paisley piert, and Spurweed. This product will also control these weeds soon after emergence.

For control of summer annual weeds listed in the "General Information" section of the main label, apply this product in late winter before the weeds emerge. Apply in sufficient water. Irrigate with one-half inch if rainfall does not occur within 10 days of preemergence treatment.

Where annual Bluegrass is the major weed, apply 0.75 fl. ozs. of this product per 1,000 sq. ft. per application. Apply 0.75 to 1.24 fl. ozs. of this product per 1,000 sq. ft. per application for control of other weeds. Do not exceed 1.24 fl. ozs. of this product per 1,000 sq. ft. per application on newly sprigged turfgrass or on hybrid Bermudagrass such as Tiflawn, Tifway, and Ormond.

For control of summer annuals which emerge after the initial application, apply an additional 1.24 fl. ozs. per 1,000 sq. ft. at least 30 days after the initial application, but not after June 1.

Note: On newly sprigged Turfgrass, Bermudagrass, nondormant Bermudagrass or nondormant Zoysiagrass, temporary slowing or growth and yellowing may occur following application.