



**ATRAZINE PLUS LINURON
WETTABLE POWDER**

ACCEPTED
MAR 15 1985
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 19713-79

Weed Killer for Selective Weed Control in Corn

Active Ingredients:

Linuron	3-(3,4-dichlorophenyl)-1-methoxy-1-methyluera	30.8%
Atrazine	2-chloro-4-(ethylamino)-6-isopropylamino)-s-triazine	29.3%
Related Compounds		1.5%
Inert Ingredients:		38.4%

**KEEP OUT OF THE REACH OF CHILDREN
CAUTION**

Statement of Practical Treatment

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.



IF SPLASHED IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes and get medical attention.

IF SPILLED ON SKIN: Immediately remove contaminated clothing, INCLUDING SHOES, and wash skin with soap and plenty of water. If irritation develops, send for a physician.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 19713-79
EPA Est. No. 19713-MS-1

Net Contents. _____ Lbs. _____ Kilograms



Precautionary Statements

Hazard to Human and Domestic Animals

CAUTION: HARMFUL IF SWALLOWED. Do not get in eyes, on skin or on clothing. Do not breathe spray mist or dust. Do not wear contaminated clothing. Keep away from heat and open flame. Do not use or store near food or feed products. Do not use around home or recreation areas.

Environmental Hazards

Do not apply to any body of water. Avoid contamination of food, feed, water supplies, streams and ponds when cleaning equipment. Toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed.

Required clothing and equipment for mixing/loading and applying

One-piece overalls which have long sleeves and long pants constructed of finely woven fabric as specified in the USDA/EPA Guide for Commercial Applicators.

Wide-brimmed hat.

Mid-forearm to elbow-length Chemical-resistant gloves.

Any article worn while applying linuron must be cleaned before reusing. Clothing which has been drenched or has otherwise absorbed concentrated pesticide must be disposed in a sanitary landfill, or incinerated, or if allowed by state and local authorities, by burning.

Instead of clothing and equipment specified above, the applicator can use an enclosed tractor cab which provides a filtered air supply (as described by Taschenberg and Bourke, 1975).

Non-aquatic

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Rotational Crop Restrictions

Do not rotate crops used for food or feed, which are not registered for use with linuron onto areas previously treated with this chemical.

Reentry Restriction

Do not reenter treated areas for 24 hours following application unless protective clothing is worn.

Precautionary Statements

Hazard to Human and Domestic Animals

CAUTION: HARMFUL IF SWALLOWED. Do not get in eyes, on skin or on clothing. Do not breathe spray mist or dust. Do not wear contaminated clothing. Keep away from heat and open flame. Do not use or store near food or feed products. Do not use around home or recreation areas.

Environmental Hazards

Do not apply to any body of water. Avoid contamination of food, feed, water supplies, streams and ponds when cleaning equipment. Toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed.

Required clothing and equipment for mixing/loading and applying

One-piece overalls which have long sleeves and long pants constructed of finely woven fabric as specified in the USDA/EPA Guide for Commercial Applicators.

Wide-brimmed hat.

Mid-forearm to elbow-length Chemical-resistant gloves.

Any article worn while applying linuron must be cleaned before reusing. Clothing which has been drenched or has otherwise absorbed concentrated pesticide must be disposed in a sanitary landfill, or incinerated, or if allowed by state and local authorities, by burning.

Instead of clothing and equipment specified above, the applicator can use an enclosed tractor cab which provides a filtered air supply (as described by Taschenberg and Bourke, 1975).

Non-aquatic

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Rotational Crop Restrictions

Do not rotate crops used for food or feed, which are not registered for use with linuron onto areas previously treated with this chemical.

Reentry Restriction

Do not reenter treated areas for 24 hours following application unless protective clothing is worn.

GENERAL INFORMATION

DO NOT apply with aerial equipment.

IMPORTANT: Use Atrazine Plus Linuron Weed Killer only as specifically recommended. Overdosage caused by incorrect preparation of spray mixture, improper calibration of equipment, or application of a speed slower than used for calibration should be avoided as injury to the crop may result.

DO NOT apply, drain or flush equipment on or near desirable trees and plants or on areas where their roots may extend or locations where the chemical may be washed or moved into contact with their roots. DO NOT use on lawns, walks, driveways, tennis courts, or similar areas.

Prevent drift of dry powder or spray to desirable plants. Keep from contact with fertilizers, insecticides, fungicides and seeds.

Atrazine Plus Linuron Weed Killer in a wettable powder to be mixed in water and applied as a spray for the selective control of annual weeds and grasses in field corn. It is non-volatile and non-flammable.


Treatment provides control of germinating annual weed seedlings such as foxtail (including giant foxtail), barnyardgrass, crabgrass, buttonweed, lambsquarters, pigweed, purslane, mustard, ragweed, and smartweed. Treatment will not control established nutgrass, Johnsongrass, Bermuda-grass or other perennial weeds.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Make a single application as a band (over the row) or broadcast spray after planting but before crop emergence. For accuracy and convenience, planting and spraying should be combined in one operation. Apply with a properly calibrated fixed-boom power sprayer, using sufficient water (equivalent to 25 to 40 gallons per acre on a broadcast basis) to provide thorough and uniform coverage of the ground. Avoid overlapping, and shut off spray boom while starting, turning, slowing or stopping, or injury to the crop may result. Continuous agitation is required to keep the material in suspension.

Best results are obtained when Atrazine Plus Linuron Weed Killer is moved by moisture (rainfall or overhead irrigation into the upper layer of soil within 2 weeks after application). Lack of sufficient moisture (1 to 2 inches is generally sufficient) following treatment will generally result in unsatisfactory weed control.



Soils should be well prepared and free as possible from trash and clods. Seeds should be planted to a depth of at least 1 3/4 inches or crop injury may result. If soil becomes crusted before corn emerges a shallow rotary hoeing may be made. If rainfall is insufficient to activate the chemicals following treatment, rotary hoeing or stand cultivation should be used before weeds become well established; otherwise avoid cultivating or disturbing the sprayed area, such as covering treated band with soil while cultivating middles. Fields treated with Atrazine Plus Linuron Weed Killer may be planted to the same crop if initial seeding fails to germinate or produce a stand. Thoroughly re-work soil before planting. Do not retreat field with a second application during the same crop year as injury to the crop may result.

Do not replant treated areas with crops other than corn within 6 months after application or crop may be injured.

Do not follow treated corn with sugar beets, tobacco or vegetables in rotation.

Prior to replanting to any crop following treated corn, thorough seedbed preparation including fall or spring plowing is recommended.

APPLY THE SPRAY AT THE FOLLOWING RATES

Soil Type*	Lbs. of Atrazine Plus Linuron Weed Killer in 25 to 40 gallons water per acre (of ground actually sprayed.)	Example-Band treatment Lbs/Crop Acre 14" Band on 42" Rows.
<u>Light colored-Low organic matter 2% or less</u>		
Sandy Loam	1.6	.5
Silt Loam	2.0	.7
Clay Loam	2.5	.8
<u>Dark colored-moderate organic matter</u>		
Silt Loam	4.0	1.3
Clay Loam	5.0	1.7

*DO NOT USE ON SAND OR LOAMY SAND SOILS AS CROP INJURY MAY RESULT.

Braille characters on the left margin.

Soils should be well prepared and free as possible from trash and clods. Seeds should be planted to a depth of at least 1 3/4 inches or crop injury may result. If soil becomes crusted before corn emerges a shallow rotary hoeing may be made. If rainfall is insufficient to activate the chemicals following treatment, rotary hoeing or stand cultivation should be used before weeds become well established; otherwise avoid cultivating or disturbing the sprayed area, such as covering treated band with soil while cultivating middles. Fields treated with Atrazine Plus Linuron Weed Killer may be planted to the same crop if initial seeding fails to germinate or produce a stand. Thoroughly re-work soil before planting. Do not retreat field with a second application during the same crop year as injury to the crop may result.

Do not replant treated areas with crops other than corn within 6 months after application or crop may be injured.

Do not follow treated corn with sugar beets, tobacco or vegetables in rotation.

Prior to replanting to any crop following treated corn, thorough seedbed preparation including fall or spring plowing is recommended.

APPLY THE SPRAY AT THE FOLLOWING RATES

Soil Type*	Lbs. of Atrazine Plus Linuron Weed Killer in 25 to 40 gallons water per acre (of ground actually sprayed.)	Example-Band treatment Lbs/Crop Acre 14" Band on 42" Rows.
<u>Light colored-Low organic matter 2% or less</u>		
Sandy Loam	1.6	.5
Silt Loam	2.0	.7
Clay Loam	2.5	.8
<u>Dark colored-moderate organic matter</u>		
Silt Loam	4.0	1.3
Clay Loam	5.0	1.7

*DO NOT USE ON SAND OR LOAMY SAND SOILS AS CROP INJURY MAY RESULT.



EQUIPMENT

Use a tractor mounted power sprayer properly calibrated to a constant speed and rate of delivery. (SEE CALIBRATION SECTION). Sprayer should be designed to treat the same number of rows as the planter used.

Make certain spray equipment is clean (free of scale, rust, dirt, oil, and pesticide deposits). Screens should be 50-mesh or coarser. Booms and hoses should be maximum of 1/2" diameter. MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION. Agitate by mechanical or hydraulic means. With hydraulic system, use a separate agitation line in addition to bypass line; both lines should terminate at bottom of tank to minimize foaming. Spray booms must be shut off while starting, turning, slowing or stopping or injury to crop may result.

CALIBRATION: The following is designed for application of indicated number of gallons spray per acre of actually sprayed; it is based on width of band treated and is not effected by row spacing.

STEP 1: To calibrate, use tractor gear and throttle setting which will be used for actual application (full throttle is preferred to insure maximum pump capacity). With only water in the tank, determine number of seconds required for tractor to travel in the field the measured distance shown below--opposite desired band width and under proper spray volume.


BAND WIDTH	GALLONS SPRAY PER ACRE OF GROUND ACTUALLY SPRAYED			
	25	30	35	40
10 in.	523 Ft.	436 Ft.	363 Ft.	326 Ft.
12 "	436 "	363 "	311 "	272 "
14 "	374 "	311 "	266 "	232 "
16 "	327 "	272 "	233 "	204 "
18 "	291 "	242 "	207 "	181 "
20 "	262 "	218 "	187 "	163 "

Adjust height of boom to obtain desired band width. With only water in the tank and using the same throttle setting as in STEP 1, but with tractor standing still adjust the pressure so that EACH NOZZEL delivers ONE QUART of spray in the exact number of seconds as determined in STEP 1 for the selected band width.

STEP 2: Equipment is now ready to apply the required spray volume, provided nozzle height, gear and throttle settings are not changed.

EQUIPMENT CLEANING

Thoroughly clean all traces of Atrazine Plus Linuron Weed Killer from application equipment immediately after use otherwise crop injury may result when equipment is used again. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).



STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal.

STORAGE INSTRUCTIONS

Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. Do not allow storage in a damp or humid area. Do not stack over 2 pellets high. Move bags carefully so as not to tear or puncture. Do not move from one area to another unless they are securely sealed to prevent dust from escaping. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Avoid contamination with acids and alkalies. Store in original containers only. If the contents are leaking or material is spilled follow these steps while wearing protective equipment:

1. Collect and place in suitable containers for disposal.
2. Wash area with water and soap to remove remaining pesticide.
3. Follow washing with clear water rinse.
4. Do not allow run off to enter sewer or contaminate water supplies.
5. Dispose of waste as indicated below:

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY - CONDITION OF SALE:

..... DIRECTIONS FOR USE of this product are based on field use and tests
..... believed reliable and should be followed carefully. It is however
..... impossible to eliminate all risks associated with use of this product.
..... Because such factors as weather conditions, foreign material and manner
..... of use for application are all beyond the control of the Seller of this
..... product, such things as crop injury, ineffectiveness or other unintended
..... consequences may result.

..... ALL SUCH RISKS ARE ASSUMED BY THE BUYER

..... Seller warrants that this product conforms to the chemical description
..... on the label and is reasonably fit for the purposes referred to in the
..... directions for use. Seller makes no other warranties, express or implied,
..... including FITNESS OR MERCHANTABILITY. In no case shall the Seller be liable
..... for consequential, special or indirect damages resulting from the use or
..... handling of this product. The foregoing is a condition of sale by Seller
..... and is accepted as such by the Buyer.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal.

STORAGE INSTRUCTIONS

Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. Do not allow storage in a damp or humid area. Do not stack over 2 pellets high. Move bags carefully so as not to tear or puncture. Do not move from one area to another unless they are securely sealed to prevent dust from escaping. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Avoid contamination with acids and alkalis. Store in original containers only. If the contents are leaking or material is spilled follow these steps while wearing protective equipment:

1. Collect and place in suitable containers for disposal.
2. Wash area with water and soap to remove remaining pesticide.
3. Follow washing with clear water rinse.
4. Do not allow run off to enter sewer or contaminate water supplies.
5. Dispose of waste as indicated below:

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY - CONDITION OF SALE:

DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of the Seller of this product; such things as crop injury, ineffectiveness or other unintended consequences may result.

ALL SUCH RISKS ARE ASSUMED BY THE BUYER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. Seller makes no other warranties, express or implied, including FITNESS OR MERCHANTABILITY. In no case shall the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by Seller and is accepted as such by the Buyer.