

RESTRICTED USE PESTICIDE
(GROUND WATER CONCERNS)

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

THIS PRODUCT IS A RESTRICTED USE HERBICIDE DUE TO GROUND WATER CONCERNS. USERS MUST READ AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE TO REACH GROUND WATER.

DREXEL


ATRAZINE 5L
HERBICIDE

For Season-Long Weed Control in Corn, Established Orchardgrass Grown for Seed (Pacific Northwest Only), Conifers, Fallowland, Guava, Macadamia Nuts, Sorghum, Sugar Cane, Lawns, Turf, Chemical Fallow, Forest and Christmas Tree Plantations, Perennial Ryegrass Grown for Seed (Pacific Northwest Only), and Nonselective Weed Control on Noncrop Sites.

ATRAZINE 5L CONTAINS 5 LBS. ACTIVE INGREDIENTS PER GALLON.

KEEP OUT OF REACH OF CHILDREN

CAUTION



ACTIVE INGREDIENTS:

Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine)	50.81%
Related Compounds	2.67%

INERT INGREDIENTS: 46.52%

TOTAL 100.00%

EPA Reg. No. 19713-80
EPA Est. No. 19713-MS-1

NET CONTENTS: 2.5 Gallons / 9.45 Liters

4-5-90

SHAKE WELL BEFORE USING



MANUFACTURED BY:

DREXEL CHEMICAL COMPANY
P.O. BOX 9306, MEMPHIS, TENN 38109-0306

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

HARMFUL IF SWALLOWED. Avoid Breathing of Spray. Do Not Take Internally. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating or drinking. Users are required to wear long sleeve shirts and long pants or equivalent, chemical resistant gloves, and boots (waterproofed). In addition, persons involved in mixing/loading operations are required to use chemical resistant rubber or neoprene gloves and a face shield or goggles. Change to clean clothing daily.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact your local Poison Control Center, hospital or physician. If the patient is unconscious, maintain breathing and heartbeat (cardiopulmonary resuscitation). If the patient is conscious and alert, induce vomiting (syrup of ipecac or stimulate the back of the throat with finger). Never Give Anything by Mouth to an Unconscious Person!

IF ON SKIN: Remove contaminated clothing and wash skin with soap and plenty of water. If irritation develops, get medical attention.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: There is no specific antidote for atrazine. If this product is ingested, induce emesis or lavage stomach. The use of an aqueous slurry of activated charcoal may be considered.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil, and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loam sand soils where the water table (groundwater) is close to the surface and where

these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product may not be mixed/loaded, or used within 50 feet of all wells including abandoned wells, drainage wells, and sink holes.

FARMWORKER SAFETY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to worker who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must inform workers of areas or fields that may not be entered without specific protective clothing until sprays have dried, and appropriate actions to take in case of accidental exposure, as described under Precautionary Statements on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must contain the following information: CAUTION. Area treated with Atrazine on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, flush eyes or skin with plenty of water. Call physician if irritation persists. Remove and wash contaminated clothing before reuse.

GENERAL INFORMATION

Atrazine 5L Herbicide will control most annual broadleaf and grass weeds in corn, sorghum, sugarcane, and ~~in certain~~ other crops as specified on this label, such as barnyardgrass, witchgrass (Echinochloa capillare), yellow foxtail, green foxtail, wild oats, large (hairy) crabgrass, giant foxtail, velvetleaf, morning-glory, lambsquarters, pigweed, ~~amaranth~~, nightshade, purslane and mustard. ~~It is~~

BEST AVAILABLE COPY

also effective in non-crop areas and industrial sites for control of most annual and many perennial broadleaf and grass weeds. Atrazine 5L may be applied before or after weeds emerge.

In each case where a range of rates is given, the lower rate should be used on coarse textured soil and soils low in organic matter, and the higher rate should be used on fine textured soils and soils high in organic matter.

Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected and weeds controlled by this product are expected to be present along with resistant biotypes, we recommend the use of those products in combinations or in sequence with other registered herbicides which are not triazines. If only resistant biotypes are expected to be present, use a registered non-triazine herbicide. Consult with your state Agricultural Extension Service for specific recommendations.

Since Atrazine 5L acts mainly through root absorption, its effectiveness depends on rainfall or irrigation to move it into the root zone. Should weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control.

Atrazine 5L is non-flammable.

Care should be taken to avoid using Atrazine 5L near adjacent desirable plants or in greenhouses or injury may occur.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

NOTE: The Seller does not recommend the use of Atrazine 5L in combination with other herbicides or oils except as specifically described on the label or in literature published by the Seller.

FAILURE TO FOLLOW ALL PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

APPLICATION PROCEDURE:

Ground Applications: Use conventional ground sprayers, equipped with nozzles that provide accurate and uniform

BEST AVAILABLE COPY

application. Be certain that nozzles are uniformly spaced and are the same size. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 10 gallons of spray mixture/A for all preplant incorporated, preplant surface, preemergence, and postemergence applications (with or without oil surfactant) with ground equipment.

Use a pump with capacity to (1) maintain 35-40 psi at nozzles, (2) provide sufficient agitation in tank to keep mixture in suspension, and (3) to provide a minimum of 20% bypass at all times. Use centrifugal pumps which provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gallons/minute/100 gallon tank size circulated through a correctly positioned sparger tube or jets.


Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For band application, calculate the amount to be applied per acre as follows:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

Aerial Application

Use aerial application only where broadcast applications are specified. Apply in a minimum of 1 qt. of water for each quart of Atrazine 5L applied per acre. For post-emergence treatments on corn and sorghum, apply recommended rate in a minimum of 2 gal. of water/A. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.



Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin, and should wash thoroughly before eating and at the end of each day's operation.

ATRAZINE 5L IN WATER APPLICATIONS

ATRAZINE 5L a liquified formulation, should be mixed with water and applied as a spray. Four ATRAZINE 5L into the tank during or after filling. Hydraulic (jet) or mechanical agitation is recommended during mixing and application to keep the materials in suspension. All return lines to the tank must discharge below liquid level and agitation should not be so violent as to cause air bubbles to form in the liquid. Wash sprayer thoroughly after use.

ATRAZINE 5L IN LIQUID FERTILIZER APPLICATIONS

Nitrogen solutions or complete liquid fertilizers may replace all or part of the water as a carrier for pre-emergence and preplant applications of ATRAZINE 5L on corn and sorghum. Mixing should be accomplished as described under water applications. Check the compatibility of this product with liquid fertilizer and/or nitrogen solution as shown below before use. Do not apply in nitrogen solution or complete liquid fertilizer after corn or sorghum emerges, except as noted under Lay-by treatment for corn, or crop injury may occur.

Compatibility Test: Since liquid fertilizers can vary, even within the same analysis, always check compatibility with herbicide(s) each time before use. Be especially careful when using complete suspension or fluid fertilizers as serious compatibility problems are more likely to occur. Commercial application equipment may improve compatibility in some instances. The following test assumes a spray volume of 25 gal per acre. For other spray volumes, make appropriate changes in the ingredients. Check compatibility using this procedure:

1. Add 1 pint of fertilizer to each of 2 one-quart jars with tight lids.
2. To one of the jars add 1/4 tsp. of 1.2 milliliters of a compatibility agent approved for this use, such as **Coatex** or **Unite** (1/2 tsp. is equivalent to 2 pt per 100 gal spray). Shake or stir gently to mix.

3. To both jars add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix. The appropriate amount of herbicides for this test follows:

Dry herbicides: For each pound to be applied per acre, add 1.5 level teaspoons to each jar.

Liquid herbicides: For each pint to be applied per acre, add 0.5 teaspoon or 2.5 milliliters to each jar.

4. After adding all ingredients, put lids on and tighten, and invert each jar ten times to mix. Let the mixture stand 15 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (A) slurry the dry herbicide(s) in water before addition, or (B) add 1/2 of the compatibility agent to the fertilizer and the other 1/2 of the emulsifiable concentrate or flowable herbicide before addition to the mixture. If incompatibility is still observed, do not use the mixture.

ATRAZINE 5L PLUS EMULSIFIABLE OIL OR OIL CONCENTRATE

Adding emulsifiable oil (petroleum-derived, petroleum-derived oil concentrate, or single or mixed crop-derived oil concentrate) to postemergence water-based sprays in corn and sorghum may improve weed control. However, under certain conditions, the use of either type of oil may seriously injure the crop. To minimize this possibility, observe the following directions:

Use one of the following properly emulsified:

1. A suitable oil concentrate containing at least 1% but not more than 20% suitable emulsifier or surfactant blend.
2. Petroleum-derived oil containing at least 1% suitable emulsifier.

Note: In the event of a compatibility problem when mixing oil with this product and water, a compatibility agent such as Complex or Daple should be used. Any of the above oils combined with the water or other materials can cause compatibility problems and/or crop injury.

BEST AVAILABLE COPY

MIXING PROCEDURES - all uses: (1) Be sure sprayer is clean and not contaminated with any other materials or crop injury or sprayer clogging may result. (2) Fill tank 1/4 full with clean water, nitrogen solution, or complete liquid fertilizer. (3) Start agitation. (4) Be certain that the agitation system is working properly and creates a rippling or rolling action on the liquid surface. (5) Pour product directly from container into tank. (6) Continue filling tank until 90% full. Increase agitation if necessary to maintain surface action. (7) Add emulsifiable oil, oil concentrate, or tank mix herbicide (s) after this product is thoroughly suspended. (8) Finish filling tank. (9) Empty tank as completely as possible before refilling to prevent buildup of oil or emulsifiable concentrate residue. Maintain agitation to avoid separation of materials. (10) If an oil or emulsifiable concentrate film starts to build up in tank, drain it and clean with strong detergent solution or solvent. (11) Clean sprayer thoroughly immediately after use by flushing system with water containing a detergent.

INSTRUCTIONS FOR SAFE HANDLING

Avoid application directly to animals or humans. Flagmen or loaders should avoid inhalation of spray mist or contact with skin and should wash thoroughly before eating and at the end of each day's operation. See Precautionary Statements for Further Information.

DIRECTIONS FOR USE

It is a violation for Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Postemergence application to corn and sorghum must be made before corn and sorghum reaches 12 inches in height.

The maximum application rate for corn and sorghum is 3 lbs. a.i. per acre per calendar year. Applications for quackgrass suppression in corn and sorghum are restricted to a spring application only. No fall applications are permitted.

Application rates to non crop land for industrial weed control cannot exceed a maximum of 10 lbs. active ingredient per acre per calendar year.



ATRAZINE 5L APPLIED ALONE - CORN OR GRAIN SORGHUM

Preplant Surface-Applied, Preplant Incorporated, or Pre-emergence (or Post-emergence at 3.2 ot/A with Oil)

Broadleaf or Grass Weeds Controlled

barnyardgrass (watergrass)	cocklebur	nightshade
giant foxtail	groundcherry	pigweed
green foxtail	jimsonweed	purslane
large (hairy) crabgrass	kochia	ragweed
wild oats	lambsquarters	velvetleaf (buttonweed)
witchgrass (Panicum capillare)	annual morning-glory	
yellow foxtail	mustards	

Post-emergence with Emulsifiable Oil or Oil Concentrate in Water (2 pt. Oil plus 1.9 pt. product/A)

Broadleaf Weeds Controlled

annual morning-glory	pigweed
cocklebur	ragweed
jimsonweed	smartweed
lambsquarters	wild buckwheat
mustards	velvetleaf

CORN

ATRAZINE 5L may be applied either before planting, at planting, or after planting as indicated below.

Preplant Surface-Applied (Broadleaf and grass control): Use on medium-and fine-textured soils with minimum-tillage or no-tillage systems only in CO, IL, IN, IA, KS, KY, MN, MO, MT, NE, ND, SD, WI, WY. Apply 2/3 the recommended rate of ATRAZINE 5L shown in Table 1 for the soil texture as a split treatment 30-45 days prior to planting and the remainder at planting. Applications made less than 30 days prior to planting may be as either a split or single treatment. On coarse-textured soils, do not apply more than two weeks prior to planting.

If weeds are present at the time of treatment, apply in tank mix combination with a contact herbicide (for example, paraquat or glyphosate). Observe directions for use, precautions, and restrictions on the label of the contact herbicide.

NOTE: To the extent possible, do not move treated soil out or the top or move untreated soil to the surface during planting, or weed control will be diminished.

Preplant Incorporated (Broadleaf and Grass Weed Control):

Broadcast in the spring after plowing at the rate indicated in Table 1. Application may be made before, during or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation of ATRAZINE 5L. Best results have been obtained when ATRAZINE 5L is applied within two weeks prior to planting.

PRE-EMERGENCE (Broadleaf and Grass Weed Control):
Apply during or shortly after planting prior to weed emergence at the rate indicated in Table 1.

POST-EMERGENCE (Broadleaf and Grass Weed Control):
Apply before weeds exceed 1.5 inches in height, at the rate indicated in Table 1.

TABLE 1

For Control of Broadleaf and Grass Weeds*

SOIL TEXTURE	ATRAZINE 5L BROADCAST RATE/A
COARSE	
Sand, loamy sand, sandy loam	3.2 pts.
MEDIUM	
Silt and clay loam low in organic matter	3.8 pts.
FINE	
Silt and clay loam with medium to high organic matter, clay (including the dark prairie soils of the Corn Belt)	4.8 pts.
Peat, muck, high organic clay (post-emergence only)	4.8 pts.

*Broadleaf control (eastern CO, western KS, western NE, NM, OK Pan Handle, west TX, and eastern WY): On sand, loamy sand, sandy loam, mild to strongly alkaline soil and all recently leveled soil, apply no more than 1.9 pt/A, either preplant or preemergence. On other soils in these areas, apply rate in Table 1 for broadleaf and grass control.

LAY-BY TREATMENT (Broadleaf and Grass Control):
Broadcast 1.4 to 3.2 pt/A before weeds are 1.5 inches tall and corn is 12 inches tall. When using nitrogen solutions,

direct the spray to lower 3-4 inches of cornstalks to avoid corn foliage injury.

POST-EMERGENCE WITH EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER:

Add the following volume of one of the type oils indicated for aerial or ground application unless the oil label specifies otherwise:

Type Oil	Ground Application	Aerial Application
Oil concentrate (Crop or petroleum-derived)	1 qt/A	.5 - 1 qt/A
Petroleum-derived oil	2.4 qts/A	2 qt/A

NOTE: Crop-derived or petroleum-derived oil concentrates should contain at least 1% but not more than 20% suitable emulsifier or surfactant blend. Petroleum-derived oils should contain at least 1% suitable emulsifier.

BROADLEAF AND GRASS CONTROL: For post-emergence control of those weeds listed under Preplant Incorporated and Pre-emergence. Broadcast 3.2 pt/A plus emulsifiable oil or oil concentrate after weed emergence, but before weeds reach 1.5 inches in height.

BROADLEAF CONTROL: For post-emergence control of those weeds listed under Post-emergence with emulsifiable oil or oil concentrate in water. Broadcast 1.9 pt/A plus emulsifiable oil or oil concentrate before pigweed and lambsquarters reach 6 inches in height and before all other weeds reach 4 inches in height. A cultivation may be necessary if all weeds are not controlled or if weeds regrow.

BEST AVAILABLE COPY

PRECAUTIONS FOR APPLICATIONS OF ATRAZINE 5L PLUS EMULSIFIABLE OIL IN WATER TO CORN

1. Do not use oil in ATRAZINE 5L sprays when corn is under stress from prolonged cold, wet weather, poor fertility, or other factors, or when corn is wet and succulent from recent rainfall as crop injury may occur.
2. Do not use oils in sprays when treating inbred lines or any breeding stock as injury may occur.
3. Adding other insecticides, herbicides, liquid fertilizers or other materials is not recommended with ATRAZINE 5L and emulsifiable oil in water because they cause compatibility problems or crop injury.
4. Store and handle emulsifiable oil carefully. Oil contaminated with even a small amount of water may not emulsify properly when added to the tank.
5. Do not make more than one application of ATRAZINE 5L and emulsifiable oil in water per season (except as recommended for control of yellow nutsedge, quackgrass, and Canada thistle on this label).

PROBLEM WEEDS

FOR CONTROL OF YELLOW NUTSEGE AND CANADA THISTLE

ATRAZINE 5L will control yellow nutsedge (*Cyperus esculentus*) and Canada thistle (*Cirsium arvense*) when applied as indicated below. For best results, ATRAZINE 5L should be used each year until yellow nutsedge or Canada thistle is eliminated or reaches a level of infestation where neither weed species is a problem. If regrowth of yellow nutsedge or Canada thistle occurs following the last application of ATRAZINE 5L during the season, cultivate once. When ATRAZINE 5L is applied post-emergence to the weeds, 1 gallon of emulsifiable crop oil should be applied per acre.

For control of yellow nutsedge and Canada thistle with ATRAZINE 5L several alternative methods of use are available. These methods are listed in order of preference below. If other weed species, such as annual grasses, are also present, use alternative 2 or 4.

1. Broadcast 3.2 pints plus 1 gal. of emulsifiable oil or 1 qt. of oil concentrate. Apply after crop and yellow nutsedge or Canada thistle emerge, but before yellow nutsedge reaches a height of 3 inches or Canada thistle reaches a height of 2 inches. Broadcast application before

13/52

lay-by (12 inches), 10-20 days after the first application.

2. Broadcast 2.4 pt/A preplant incorporated for control of yellow nutsedge, or broadcast 2.4 pt/A during or shortly after planting, but prior to crop or weed emergence for control of Canada thistle. Follow with an application of 3.2 pt/A plus 1 gal. of emulsifiable oil or 1 qt. of oil concentrate after corn and weeds emerge, but before yellow nutsedge or Canada thistle reaches height of 6 inches.
3. Broadcast 4.8 pt/A plus 1 gal. of emulsifiable oil or 1 qt. of oil concentrate after crop emerges, but prior or lay-by (12 inches). Apply after yellow nutsedge and Canada thistle emerge, but before yellow nutsedge is 3 inches tall or Canada thistle is 6 inches tall.
4. Broadcast 4.8 pt/A before, during or shortly after planting, but before crop or weed emergence (yellow nutsedge control only).

NOTE: Do not use emulsifiable oil or oil concentrate when corn is wet or under stress, especially when using 4.8 pt/A of this product, or crop injury may occur. See "Precautions for applications with emulsifiable oil or oil concentrate in water" for additional directions.

FOR QUACKGRASS CONTROL ON LAND GOING INTO CORN PRODUCTION

Split Application: Broadcast 2.4 pts. acre in the spring and plow 1 - 3 weeks later. Broadcast a second application at the rate of 2.4 pts. per acre in the spring before, during or after planting, but before weeds are 1.5 inches high. This split application will control both quackgrass and most annual broadleaf and grass weeds.

Single Application: Broadcast 4.8 pts per acre in the spring. Plow 1-3 weeks after application.

TANK MIXTURES WITH ATRAZINE 5L FOR CORN

This product may be tank mixed with these herbicides for control of certain broadleaf and grass weeds in corn:

metolachlor	alachlor
metolachlor + paraquat	alachlor + glyphosate
metolachlor + glyphosate	alachlor + paraquat
metolachlor + simazine	paraquat
metolachlor + simazine + paraquat	glyphosate
metolachlor + simazine + glyphosate	simazine
promachlor	simazine + paraquat
	simazine + glyphosate
	butylate 4.7E

BEST AVAILABLE COPY

Use tank mix directions appearing on the labels of the above herbicides when tank mixing with this product. Observe all precautions and limitations on labeling of products used in a particular tank mix.

NOTE: When the labels of the above herbicides refer to Atrazine 80W, use equivalent rates of ATRAZINE 5L. One lb of 80W equals 1.28 pt of 5L.

ATRAZINE 5L + PARAQUAT

For kill of Existing Vegetation and Residual Weed Control where Corn Will Be Planted Directly Into a Cover Crop, Established Sod or in Previous Crop Residues.

Apply 3.2-4.8 pts. ATRAZINE 5L plus 1 to 2 pts. paraquat per acre, as a broadcast spray. Refer to both labels for directions, limitations and cautions. Apply 20 to 60 gallons diluted spray per acre. Add Surf-Ac 820 or another non-ionic surfactant at the rate of 8 fl. oz. per 100 gallons diluted spray. Add Atrazine to spray tank first mixing thoroughly in water. Then add paraquat and surfactant. Constant agitation recommended at all time.

Atrazine plus paraquat may be applied in liquid nitrogen and/or complete liquid fertilizer solutions when 25 or more gallons of dilute solution are applied per acre in combination with 1 or 2 pts. of Surf-Ac 820 or another non-ionic surfactant per 100 gallons. Equipment must be adjusted and calibrated to obtain thorough coverage of undesirable vegetation. Inadequate coverage and/or absence of a non-ionic surfactant will result in severe reduction of paraquat activity. Phosphate containing liquid fertilizer solutions will reduce the activity of paraquat. When liquid fertilizer solutions containing phosphorous are used with paraquat, and a rapid kill is desired, use the high paraquat rate.

Refer to the paraquat label for further directions, limitations and cautions.

ATRAZINE 5L PLUS ALACHLOR 4 EC PLUS PARAQUAT

For kill of Existing Vegetation and Residual Weed Control Where Corn Will Be Planted Directly Into a Cover Crop, Established Sod or in Previous Crop Residues.

Apply 1.6-3.2 pts. ATRAZINE 5L plus 4 to 5 pts. alachlor 4 EC plus 1 to 2 pts. paraquat per acre prior to, during, or after planting, but before corn emergence. Use 20 to 60 gal. of diluted spray per treated area. Add Surf-Ac 820 or 8 oz. per 100 gal. of diluted spray. Add the atrazine to the spray tank while agitating, then add the alachlor.

BEST AVAILABLE COPY

paraquat and Surf-Ac 820. Continue agitation during application. Refer to alachlor label for caution and for specific rates for a given soil type.

ATRAZINE 5L PLUS METOLACHLOR and ATRAZINE 4L PLUS METOLACHLOR PLUS SIMAZINE 4L or 80W

Preplant Incorporated or Pre-emergence: In addition to the weeds controlled by ATRAZINE 5L alone, metolachlor + ATRAZINE 5L or metolachlor + ATRAZINE 5L + Simazine 4L applied preplant surface, preplant incorporated, or pre-emergence, also controls the following weeds: browntop panicum, cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf.

Apply metolachlor + ATRAZINE 5L or metolachlor + ATRAZINE 5L + Simazine 4L preplant surface, preplant incorporated, or pre-emergence.

BEST AVAILABLE COPY

16/52

Preplant Surface Applied: Follow instructions for use of metolachlor alone under Application Procedures and under application instructions for metolachlor alone on corn. Apply metolachlor + ATRAZINE 5L or metolachlor + ATRAZINE 5L + simazine 4L on medium soils (2.5 pts./A metolachlor + 2.6-3.2 pts./A of ATRAZINE 5L or simazine 4L or ATRAZINE 5L + simazine 4L combined) and on fine soils 2.5-3 pts./A metolachlor + 3.2-4 pts./A of ATRAZINE 5L or ATRAZINE 5L + simazine 4L combined in minimum-tillage and no-tillage systems in CO, IL, IN, IA, KS, KY, MN, MO, MT, NE, ND, SD, WI, WY. Apply the tank mixtures as a split or single treatment in those states and as indicated in the metolachlor alone Preplant Surface Applied section of the metolachlor label for corn. On coarse soils, apply 2 pts./A of metolachlor and 2.6 pts./A of ATRAZINE 5L or ATRAZINE 5L + simazine 4L combined.

When using simazine 80W use equivalent rates.

Preplant Incorporated or Pre-emergence: Follow instructions for use of metolachlor alone under Application Procedures. Apply metolachlor + ATRAZINE 5L or metolachlor + ATRAZINE 5L + simazine 4L using the appropriate rates from Table 2.

Note: Do not make more than one application per year.


TABLE 2

Broadcast Rates Per Acre

	Less than 3% Organic Matter		3% organic matter or greater	
	Atrazine [®] 5L + Metolachlor	Atrazine 5L ^{**} OR + Metolachlor + Simazine ^{**} 4L or 80W	Atrazine [®] 5L + Metolachlor	Atrazine 5L ^{**} OR + Metolachlor + Simazine ^{**} 4L or 80W
SOIL TEXTURE COARSE: Sand, loamy sand, silty loam	1.6 - 3.2 pts + 1.25 - 1.5 pts	0.8 - 1.6 pts + 1.25 - 1.5 pts + 1.1-2 pts Sim 4L -or- .7 - 1.2 lbs Sim 80W	1.9 - 3.2 pts + 1.5 pts	1.0 - 1.6 pts + 1.5 pts + 1.25 - 2 pts Sim 4L -or- .3 - 1.2 lbs Sim 80W
MEDIUM: Loam, silt loam silt	1.9 - 3.2 pts + 1.5 - 2 pts	1.0 - 1.6 pts + 1.5 - 2 pts + 1.25 - 2 pts Sim 4L -or- .8 - 1.2 lbs Sim 80W	2.6 - 3.2 pts + 2 pts	1.3 - 1.6 pts + 2 pts + 1.6 - 2 pts Sim 4L -or- 1 - 1.2 lbs Sim 80W
FINE: Silty clay loam, sandy clay loam, silty clay, sandy clay, clay loam, clay	2.6 - 3.2 pts + 2 pts	1.3 - 1.6 pts + 2 pts + 1.6 - 2 pts Sim 4L -or- 1 - 1.2 lbs Sim 80W	2.6 - 3.2 pts ^{***} + 2 - 2.5 pts ^{***}	1.3 - 1.6 pts ^{***} + 2 - 2.5 pts + 1.6 - 2 pts Sim 4L ^{**} -or- 1 - 1.2 lbs Sim 80W

Muck or peat

DO NOT USE



*On soils having between 6% and 20% organic matter, metolachlor may be used up to 3.5 pts./A in tank mix combination with 4.8 pts./A of ATRAZINE 5L.

**When using the tank mixture of ATRAZINE 5L +metolachlor + simazine 4L or 80W, use equal rates of each as shown when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use a 1:2 ratio of active ATRAZINE + simazine instead of 1:1 ratio given in Table 2 above.

***After cocklebur, yellow nutsedge, and velvetleaf control on fine-textured soils above 3% organic matter: Apply 3.2 pts. ATRAZINE 5L or the same total amount of Atrazine + simazine with 2-2.5 pts. of metolachlor/A.

ATRAZINE 5L PLUS METOLACHLOR POST-EMERGENCE

WEEDS CONTROLLED:

- | | | | |
|---------------|----------------|---------------|-----------|
| barnyardgrass | green foxtail | lambsquarters | ragweed |
| (watergrass) | yellow foxtail | mustard | smartweed |
| crabgrass | pigweed | velvetleaf | |
| crowfootgrass | jimsonweed | prickly sida | |
| fall panicum | kochia | purslane | |
| giant foxtail | | | |

Weeds Partially Controlled: cocklebur, morning-glory, yellow nutsedge.

Apply early post-emergence using the appropriate rates from Table 3.

Apply this tank mixture before grass and broadleaf weeds pass the 2-leaf stage and before corn exceeds 5 inches in height. Application to weeds larger than the 2-leaf stage will generally give unsatisfactory control. Occasionally some corn leaf burn may result, but this should not affect later growth or yield. Do not apply this post-emergence tank mixture in fluid fertilizer, as severe crop injury may occur.

BEST AVAILABLE COPY

Table 3: ATRAZINE 5L plus Metolachlor

POST-EMERGENCE CORN

SOIL TEXTURE	BROADCAST RATES PER ACRE	
	ATRAZINE 5L	+ METOLACHLOR
COARSE: Sand, loamy sand, sandy loam	1.9 pts.	1.5 pts.
MEDIUM: Loam, silt, silt loam	2.6 pts.	2 pts.
FINE: Sandy clay loam, silty clay loam, clay loam, sandy clay, silty clay, clay	2.6 to 3.2 pts.*	2-2.5 pts.

*For better control of cocklebur, morning-glory, velvetleaf and yellow nutsedge on fine-textured soils above 3% organic matter, apply 3.2 pts. ATRAZINE 5L with 2-2.5 pts. of metolachlor per acre.

ROTATIONAL CROPS:

Follow the crop rotation instructions on the metolachlor label for metolachlor plus Atrazine Preplant Incorporated or Pre-emergence - Corn.

ATRAZINE 5L + METOLACHLOR + SIMAZINE 4L or 80W WITH PARAQUAT OR GLYPHOSATE FOR MINIMUM - TILLAGE OR NO - TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides paraquat or glyphosate may be added to a tank mix of ATRAZINE 5L + metolachlor or ATRAZINE 5L + metolachlor + simazine 4L or 80W. When used as directed, the paraquat portion of the tank mixture controls most emerged annual weeds and suppresses many perennial weeds. Glyphosate combinations will control emerged annual and perennial weeds when applied as directed on the glyphosate label. The Atrazine 5L + metolachlor or Atrazine 5L + metolachlor + simazine 4L or 80W portion of the tank mixture provides pre-emergence control of the weeds listed on the metolachlor

label in the tank mixture section for metolachlor + Atrazine or metolachlor + Atrazine 5L + simazine Preplant Incorporated or Preemergence..

Application: Apply before, during or after planting, but before the corn emerges, at the rate in Table 4. Add paraquat or glyphosate at the following broadcast rates:

Paraquat: 1 - 2 pts.* per acre plus 8 oz. Surf-Ac 820 per 100 gals. of spray mixture. Use the lower rate for control of annual weeds less than 4 inches tall and the higher rate for weeds 4-6 inches tall. This treatment will not consistently control weeds taller than 6 inches.

*Based on products containing 2 lbs. active ingredient per gal.

NOTE: Do not apply combinations containing paraquat in suspension type liquid fertilizers as the activity of paraquat will be reduced.


Glyphosate: 1.5 qts. per acre for existing annual weeds, or 2 - 4 qts. per acre for existing perennial weeds. See the glyphosate label for weeds controlled and recommended rates for specific weeds.

Apply in 20 - 60 gals. of water or fluid fertilizer per acre with ground equipment.

TABLE 4

Broadcast Rates Per Acre

	ATRAZINE 5L* + METOLACHLOR	OR	ATRAZINE 5L** + METOLACHLOR + SIMAZINE 4L or 80W
SOIL TEXTURE			
Coarse: Sand,	1.9 pts.		1 pt.
Loamy sand, Sandy	+		+
Loam	1.5 pts.		1.5 pts.
			+
			1.25 pts. or 8 lbs.
Medium: Loam, silt	2.6 pts.		1.5 pts.
Loam, silt	+		+
	2 pts.		2 pts.



+
1.6 pts. simazine 4L
or
1 lb. simazine 80W

Fine:*** Silty clay loam, sandy clay loam,	2.6 to 3.2 pts.	1.3 to 1.6 pts.
	+	+
silty clay, sandy clay, clay loam, clay	2 to 2.5 pts.	2 to 2.5 pts.
		+
		1.6 to 2 pts. simazine 4L or
		1.0 to 1.2 lbs. simazine 80W

*Use simazine in preference to Atrazine when heavy infestations of crabgrass or fall panicum are expected.

**When using tank mixture of ATRAZINE 5L plus metolachlor plus simazine, use equal rates of Atrazine and simazine as shown when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use 1:2 ratio of ATRAZINE 5L + simazine (4L or 80W) instead of the 1:1 ratio given above.

***For cocklebur, yellow nutsedge, and velvetleaf control on ATRAZINE 5L fine-textured soils above 3% organic matter, apply 3.6 pts./A of ATRAZINE 5L, or equivalent, or the same total amount of active ATRAZINE 5L + simazine with 2-2-1/2 pts./A of metolachlor.

ATRAZINE 5L PLUS ALACHLOR 4EC

Use as tank mixture for control of many annual broadleaf and grass weeds in corn (field, hybrid seed, silage, sweet, popcorn), including barnyardgrass, carpetweed, crabgrass, fall panicum, Florida pusley, foxtails (giant, green, yellow), goosegrass, jimsonweed, kochia, lambsquarters, mustard, nightshade, pigweed, purslane, ragweed, signalgrass (Brachiaria), smartweed, and witchgrass. This tank mix will reduce competition from the hard to control annual weeds - annual morning-glory, cocklebur, velvetleaf (buttonweed), and sandbar.

For pre-emergence or early post-emergence broadcast application, use rates in Table 5. Apply tank mixture any time from immediately after planting until weeds reach 2-leaf stage and corn is no more than 5 inches tall. On sweet corn, apply before crop and weeds emerge. Application after the 2-leaf weed stage will not give satisfactory control.

With post-emergence application, occasional corn leaf burn should not affect growth or yield. If applied with liquid fertilizers, spray before crop emerges.

BEST AVAILABLE COPY

TABLE 5

Tank Mixtures of ATRAZINE 5L Plus Alachlor 4EC on Corn
(Surface Application)

Soil Texture	Surface Application Rate/A			
	Less than 3% organic matter		3% or more organic matter	
	ATRAZINE 5L	Alachlor 4EC	ATRAZINE 5L	Alachlor 4EC
COARSE loamy sand, sandy loam	.6 to .8 qts.	2 qts.	.8 to 1.2 qts.	2 qts.
MEDIUM Loam, silt loam, silt	.8 to 1.0 qts.	2 qts.	1 to 1.2 qts.	2 qts.
FINE Silty clay loam, sandy clay loam, silty clay, sandy clay, clay	.96 to 1.2 qts.	2.25 qts.	1 to 1.4 qts.	2.5 qts.

NOTE: Use the higher rates in the recommended ranges in areas of heavy weed infestations, or for fields under irrigation. Use minimum of 2.5 quarts of alachlor 4EC per acre in this tank mixture on coarse soils and 3 to 4 quarts on medium or fine textured soils to reduce competition for hard-to-control weeds.

ATRAZINE 5L PLUS ALACHLOR 4EC ON CORN
(Preplant Incorporated Application)

BEST AVAILABLE COPY

TABLE 6

Preplant Incorporated Application Rate per Acre

Soil Texture	Less than 3% organic matter		3% or more organic matter	
	ATRAZINE 5L	Alachlor 4EC	ATRAZINE 5L	Alachlor 4EC
MEDIUM Loam, silt loam, silt	1.0 qts.	2.5 qts.	1.2 qts.	2.5 qts.
FINE Silty clay loam, sandy clay loam, silty clay, sandy clay, clay	1.2 qts.	2.75 qts.	1.4 qts.	2.75 qts.

NOTE: Use 3 to 4 quarts of alachlor 4EC per acre for control of yellow nutsedge or to reduce competition from the hard-to-control weeds listed on this label.

NOTE: Two pass shallow incorporation is recommended to reduce competition from shattercane. Use the 4 quart rate for shattercane.

For preplant incorporated application, use rates in Table 6. Broadcast within 7 days before planting and incorporate 2 inches deep.

Apply tank mixture in a minimum of 20 gals. of water/A. Liquid fertilizer may replace all or part of the water. Add ATRAZINE 5L to spray tank, thoroughly mix with water, then add alachlor 4 EC. Follow mixing procedures in the section, "Application Procedures" of this label.

At least 14 weeks must elapse following treatment with ATRAZINE 5L Plus alachlor 4EC before immature corn forage can be harvested or fed to cattle.

Refer to Alachlor 4EC Label for information concerning other directions, limitations and cautions.

BEST AVAILABLE COPY

ATRAZINE 5L PLUS ALACHLOR 4EC OR 4EC PLUS PARAQUAT or ATRAZINE 5L PLUS ALACHLOR 4EC PLUS GLYPHOSATE

MINIMUM OR CONSERVATION TILLAGE SYSTEMS TANK MIXTURES.
FIELD CORN OR SILAGE CORN ONLY.

When applied as directed under the conditions described, these tank mixtures control many emerged annual weeds, suppress many emerged perennial weeds and give pre-emergence control of many annual weeds when corn will be planted directly into a cover crop, established sod or in previous crop residues. These tank mixtures will not control regrowth from perennial weeds. Do not apply by air.

For mixing instructions, see the "Mixing Instructions" section of this label.

Apply these tank mixtures in 15 to 60 gals. of water per acre immediately before, during or after planting, but BEFORE CROP EMERGENCE. As density of stubble crop residue or weeds increases, spray gallonage and rate should be increased within the recommended ranges to insure complete coverage.

Add the Atrazine to the spray tank while agitating then add the alachlor 4EC, paraquat and Surf-Ac 820. If glyphosate herbicide is used in this tank mixture, add it to the tank last instead of the paraquat and Surf-Ac 820.

CONTROL OR SUPPRESSION OF EMERGED WEEDS

Add paraquat or glyphosate to the rates of ATRAZINE 5L and alachlor 4EC given in Table 5.

PARAQUAT

When used as directed, paraquat in tank mixture with Atrazine and alachlor 4EC controls many emerged annual weeds and suppresses many emerged perennial weeds. In the absence of emerged vegetation, the paraquat should be deleted, since this product does not provide pre-emergence weed control.

Broadcast treatment - Apply 1 to 2 pints of paraquat in this tank mixture immediately before, during or after planting but BEFORE CROP EMERGENCE. Use the 2 pint rate if any weeds are 4 to 6 inches tall. This mixture may not control weeds taller than 6 inches. As density of stubble, crop residue or weeds increases, spray gallonage should be increased within the recommended range from complete coverage. Add Surf-Ac 820 surfactant at 1 oz. per 100 gals. of diluted spray. Refer to the paraquat label for precautionary statements.

BEST AVAILABLE COPY

GLYPHOSATE

Annual Weeds - Use 1.5 quarts of glyphosate herbicide per acre in tank mixture with Atrazine plus alachlor 4EC for control of many emerged annual grass and broadleaf weeds. For emerged annual weeds controlled, see the "Weeds Controlled" section of the label for glyphosate.

Perennial Weeds - At normal application dates in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. See the "General Information" section of the label for glyphosate for the proper stage of growth for perennial weeds. Use of 2 to 4 quarts of glyphosate herbicide per acre in the above tank mixtures under these conditions provides top kill and reduces competition from many emerged perennial grass and broadleaf weeds. For emerged perennial weeds controlled see the "Weeds Controlled" section of the label for glyphosate. To obtain control, follow label recommendations for stage of growth and rate of application for specific perennial weeds. To obtain the desired stage of growth, it may be necessary to apply glyphosate herbicide alone in the late summer or fall and then follow with a label approved seedling weed control program at planting.

NOTE: When using this tank mixture, do not exceed 4 quarts of glyphosate herebicide per acre.

BEST AVAILABLE COPY

USE OF THIS MIXTURE FOR BERMUDAGRASS OR JOHNSONGRASS CONTROL IS NOT RECOMMENDED.

ATRAZINE 5L PLUS PROPACHLOR 65W

Use a tank mixture of ATRAZINE 5L plus propachlor 65W for control of most annual broadleaf and grass weeds in corn (field, hybrid seed, silage, and sweet corn only), such as annual ryegrass, barnyardgrass, (watergrass), carpetweed, crabgrass, fall panicum, Florida pusley, giant foxtail, green foxtail, yellow foxtail, goosegrass, groundsel, jimsonweed, lambsquarters, mustard, nightshade, pigweed, purslane, ragweed, smartweed and sunflower. This combination also reduces competition from buttonweed (velvetleaf), cocklebur and annual morning-glory. Broadcast 1.7 to 2.6 pts. of ATRAZINE 5L plus 3.8 to 6.0 lbs. of propachlor 65W per acre on the surface any time from immediately after planting until broadleaf and grass weeds reach the two-leaf stage. Use the lower rates of Atrazine 5l and propachlor 65W on the light-textured soils low in organic matter. Use the higher rates on heavy-textured soils high in organic matter.

A minimum of 2.3 pts. per acre of ATRAZINE 5L in the tank mixture will give better control of annual morning-glory, button weed (velvetleaf), cocklebur and sunflower.

Apply in a minimum of 20 gals. of water per acre. Fluid fertilizer may replace all or part of water used as a carrier for applications applied to the soil surface before crop and weeds emerge. Add the Atrazine 5L to the spray tank first and thoroughly mix with water. Cut the propachlor 65W bag and fill with water to the fill line. Grasp the neck of the bag firmly and shake vigorously and add to the spray tank.

Refer to propachlor 65W label for further directions, limitations and cautions.

ATRAZINE 5L PLUS SIMAZINE 4L OR SIMAZINE 50W

In addition to the weeds listed under ATRAZINE 5L Applied Alone - Corn and Grain Sorghum - Preplant Surface - Applied, Preplant Incorporated, or Pre-emergence, this combination also controls crabgrass, fall panicum, and carpetweed.

Broadcast tank mix before planting, at planting, or after planting, but before crop and weeds emerge, at rates in Table 2. Use the label rate for control of most weeds. Use the 1.2 rate for control of bermudagrass, johnsongrass, and fall panicum. Do not be lulled by weed control.

Preplant Surface - Applied: Use on medium and fine textured soils with minimum-tillage or no-tillage systems only in CO, IL, IN, IA, KS, KY, MN, MO, MT, NE, ND, SD, WI, and WY. Apply 2/3 the recommended rate of ATRAZINE 4L and simazine shown in Table 7 for the soil texture as a split treatment 30-45 days prior to planting and the remainder at planting. Applications made less than 30 days prior to planting may be as either a split or single treatment. On coarse-textured soils, do not apply more than two weeks prior to planting. Refer to the ATRAZINE 5L Alone - Preplant Surface - Applied section of the corn label for additional details.

If weeds are present at time of treatment, apply in a tank mix combination with a contact herbicide (for example, paraquat or glyphosate). Observe directions for use, precautions and restrictions on the label of the contact herbicide.

NOTE: To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

Preplant Incorporated: Apply to the soil and incorporate in the spring before, during, or after final seedbed preparation. Avoid deep incorporation. For best results, apply within two weeks prior to planting.

Pre-emergence: Apply during or shortly after planting, but before crop and weeds emerge.

TABLE 7
TANK MIXTURES WITH SIMAZINE 4L OR 80W ON CORN

Soil Texture	Broad cast rate/A					
	1:1 Ratio*			1:2 Ratio**		
	Atrazine 5L	Simazine 80W	Simazine 4L	Atrazine 5L	Simazine 80W or 4L	Simazine
Sand, loamy sand, sandy loam	1.6 pts	1.25 lb	2 pts	1.1 pts	1.67 lb	2.6 pts
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with low organic matter	1.92 pts	1.5 lb	2.4 pts	1.3 pts	2 lb	3.2 pts
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with medium to high organic matter, and clay (including dark prairie soils of the Corn Belt)	2.4 pts	1.8 lb	3 pts	1.5 pts	2.4 lb	3.84 pts

*For control of most weeds.
**For control of expected heavy infestations of crabgrass and fall panicum.

Refer to Corn sections of this label and to simazine 4L or simazine 80W label for further directions, limitations and cautions.

ATRAZINE 80W PLUS SIMAZINE 4L OR 80W PLUS PARAQUAT

For Kill of Existing Vegetation and Residual Weed Control Where Corn Will Be Planted Directly Into a Cover Crop, Established Sod or in Previous Crop Residues.

Apply 1.6-3.2 pts. ATRAZINE 5L plus 2-4 pts. simazine 4L or 1.25 to 2.5 lbs. simazine 80W plus 1 to 2 pts. paraquat per sprayed acre prior to, during or after planting - but before crop emergence. Use the low rates of Atrazine and simazine on light soils, and the higher rates on heavy clay soils. Use to 20 to 60 gals. of diluted spray per sprayed acre. Add Surf-Ac 820 or another non-ionic surfactant at the rate of 8 fl. oz. per 100 gals. diluted spray. Add Atrazine to the spray tank while agitating, then add the simazine, paraquat and Surf-Ac 820. Continue agitation during application. Use the 2 pts. rate of paraquat if existing vegetation is 4-6 inches tall. This mixture will not control weeds taller than 6 inches. Refer to paraquat and simazine labels for further limitations and precautions.


ATRAZINE 5L PLUS SIMAZINE 4L OR 80W PLUS GLYPHOSATE

When applied as recommended under the conditions described, this tank mixture controls many emerged weeds, and gives pre-emergence control of many annual weeds when corn will be planted directly into a cover crop, established or in previous crop residues.

Apply this tank mixture in 20 to 60 gallons of water per acre immediately before, during or after planting, but BEFORE CROP EMERGENCE. As density of stubble, crop residue or weeds increases, spray gallonage and rate should be increased within the recommended ranges to insure complete coverage.

Annual Weeds - apply atrazine and simazine to actively growing grasses and broadleaf weeds that are 6 inches tall at the rates given in Table 8 plus 1 quart of gl phosphate per acre. If the weeds are over 6 inches tall, increase the rate of gl phosphate to 1.5 qts.

For Annual Weeds - all annual broadleaf weeds, annual grasses, tillage systems, perennial weeds; may not be at the proper



stage of growth for control. See the "General Information" section of the glyphosate label for the proper stage of growth for perennial weeds. Apply the Atrazine and simazine at the rates given in Table 8 plus 2-4 qts. of glyphosate per acre. This combination will provide top kill and reduce competition from many emerged perennial grass and broadleaf weeds.

NOTE: Do not use this tank mix on sand or loamy sand soils. Do not exceed 4 qts. of glyphosate per acre in this tank mixture. Not recommended to control bermudagrass or johnsongrass in minimum tillage systems.

TABLE 8; ATRAZINE 5L plus Simazine 4L or 80W for mixture with Glyphosate.

Soil Texture	Broadcast Rate Per Acre	
	ATRAZINE 5L*	Simazine* 4L or 80W
Coarse: sandy loam only	1.6 pts.	2 pts. or 1.25 lbs.
Medium: loam, silt loam, silt, sandy clay loam	1.6 to 2.3 pts.	2-2.75 pts/or 1.25-1.75 lbs.
Fine: silty clay loam, clay loam, sandy clay, silty clay, clay	1.9 to 2.6 pts.	2.35-3.15 pts. or 1.25-2 lbs.

*Use the higher rate of Atrazine and simazine in the recommended ranges in soil with greater than 3% organic matter.

ATRAZINE 5L PLUS BUTYLATE 6.7E

A tank mix combination of ATRAZINE 5L and butylate 6.7E can be applied and mixed into the soil before planting for weed control in corn. This combination will control a broader spectrum of weeds than will either material used alone. In addition, potential hazard of ATRAZINE residues affecting certain succeeding crops is reduced because a lower rate of ATRAZINE can be used when it is combined with butylate 6.7E than when it is used alone.

FOR THE FULL PRECAUTIONS AND WARNINGS THAT APPLY ON THE BUTYLATE 6.7E LABEL, SUPPLEMENTARY BUTYLATE 6.7E PRECAUTIONS AND WARNINGS ON THIS LABEL.

BEST AVAILABLE COPY

MAKE ONLY ONE APPLICATION PER CROP. AFTER TREATMENT WITH ATRAZINE 5L/BUTYLATE 6.7E COMBINATION. DO NOT PLANT ANY CROP EXCEPT CORN UNTIL THE FOLLOWING YEAR OR INJURY MAY OCCUR.

DO NOT USE ON MILO OR SORGHUM.

DO NOT USE ON CORN SEED STOCK (BREEDERS, FOUNDATION, OR INCREASE).

For weeds controlled by ATRAZINE 5L/butylate 6.7E see the butylate 6.7E Label.

The ATRAZINE 5L/butylate 6.7E combination can be mixed with water or combined with fluid (solution, slurry or suspension) fertilizers and applied as a spray, or can be impregnated on dry bulk fertilizers.

See Appendix III on butylate 6.7E label for Tank Mix Directions.

ATRAZINE 5L/BUTYLATE 6.7E FLUID FERTILIZER TANK MIX.

The ATRAZINE 5L/butylate 6.7E combination can be combined with fluid fertilizers for applications made before planting. However, the physical compatibility with these fluid fertilizers must be determined before combining in the spray tank, and even though found to be compatible, constant agitation is necessary to keep the ATRAZINE 5L/butylate 6.7E evenly mixed with the fluid fertilizer during application.

In some areas disking 2 to 3 times spaced at 4 to 6 week intervals during fall and winter can be done. If this cannot be done, start with fall plowing followed by several diskings in the spring before planting. The disking cuts rhizomes into small pieces and brings them to the surface, where both the weather and butylate 6.7E will kill or weaken them.

Rhizome Bermudagrass: The rhizomes and stolons of Bermudagrass must be cut up thoroughly so that four or less nodes remain on a strand. For control of bermudagrass, the disk must be set to cut 6 inches deep.

BEST AVAILABLE COPY

For Suppression of Broadleaf Signalgrass: For early and mid-season suppression, use 7-1/3 pts. of butylate 6.7E and .8 to 1.2 quarts ATRAZINE 5L per acre. Cultivation or application of a post-emergence herbicide recommended for broadleaf signalgrass will aid in extending control to late season.

For Suppression of Hemp Dogbane: For early and mid season suppression, use 7-1/3 pints of butylate 6.7E and .8 to 1.2 quarts ATRAZINE 5L per acre. Cultivation or application of a post-emergence herbicide recommended for hemp dogbane may aid in extending control beyond mid-season.

*Use the higher rate of ATRAZINE 5L on soils with over 5% organic matter.

Fall Application: For use only in Minnesota, Northern Iowa (North of Interstate 80), Northeast Nebraska (North of Interstate 80 and East of State Highway 14), South Dakota (East of Missouri River), Wisconsin and Northern Illinois (North of Interstate 80). Apply and incorporate 7-1/3 pints butylate 6.7E and .8 to 1.2 qts. ATRAZINE 5L per acre in the late Fall before the ground freezes.

See Appendix IV on the butylate 6.7E label for directions on use of dry bulk fertilizer applied in the Fall with this combination.

NOTE: Do not use Fall Application for suppression of wild cane.

CULTURAL PRACTICES FOLLOWING APPLICATION

Should weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control. When cultivating for any reason, it should be shallow; i.e., no more than 1/2 the depth the herbicide was incorporated.

RECOMMENDATIONS

NOTE: The ATRAZINE 5L/butylate 6.7E tank mix combination is not recommended on corn grown in Arizona or in the ten Southernmost California counties.

For all uses, cultivation is recommended in addition to the ATRAZINE 5L/butylate 6.7E treatment in fields with moderate to heavy infestations of wild cane, nutgrass, sandbar, johnsongrass seedlings, rhizome bermudagrass, Texas panicum or volunteer sorghum.

BEST AVAILABLE COPY

**BROADCAST RATES PER ACRE CONVENTIONAL SPRAY - ATRAZINE
5L/BUTYLATE 6.7E**

For control of annual grasses and broadleaf weeds: Use 3-3/4 to 4-3/4 pints butylate 6.7E and 1 to 1.2 quarts ATRAZINE 5L per acre. Use the higher rate of butylate 6.7E on medium (medium fine textured-silt loam, clay loam, etc.) to fine (fine textured-clay) soils and for sandbur and johnsongrass seedling control.

For control of nutsedge and Texas panicum (in addition to the broadleaf weeds and other annual grasses listed): Use 4-3/4 to 7-1/3 pints of butylate 6.7E and .8 to 1.2 qts. ATRAZINE 5L per acre. Use the higher rates butylate 6.7E on medium (medium fine to textured-silt loam, clay loam, etc.) soils, fine (fine textured-clay) soils and for heavy infestations.

For partial control (suppression) of Wild Cane (and for control of the broadleaf weeds and annual grasses listed): When applied and incorporated immediately before planting, butylate 6.7E at the rate of 7-1/3 pints and .8 - 1.2 qts. ATRAZINE 5L per acre will reduce competition from wild cane. On continuous corn acreage with high cane speed population, rotation to another crop is recommended.

NOTE: See preceding butylate 6.7E section for wild cane supplemental cultural and tillage practices.

For control of Johnsongrass and Bermudagrass (in addition of the broadleaf weeds and annual grasses listed): Apply and disc-incorporate butylate 6.7E just before planting at the rate of 4-3/4 to 7-1/3 pints plus .8 to 1.2 qts. of ATRAZINE 5L per acre. Use the higher rate of butylate 6.7E on medium (medium fine textured-silt loam, clay loam, etc.) soils, fine (fine textured-clay) soils, for rhizome johnsongrass suppression and for heavy infestations. Disk a second time at right angles to the first. Cultivate at least once, preferably twice.

Johnsongrass: Using a program of plowing and disking followed by butylate 6.7E and ATRAZINE 5L will give good control of seedlings and greatly reduce competition from plants coming from rhizomes.

PLANTING

Do not plant seed deeper than 2 inches. For a BUTYLATE 5L/butylate 6.7E application made before planting, seedling can be done immediately, or it may be delayed up to 2 weeks. To avoid removing BUTYLATE 6.7E from the soil with loss of weed control in the corn, do not move or stir soil after treatment, such as planting or deep plowing. If a corn cover crop is used, the soil following application

3/15/57

and incorporation but before planting, do not work the soil deeper than depth the herbicides were incorporated.

Refer to Appendix II on butylate 6.7E label regarding the procedure for testing the compatibility of ATRAZINE/butylate 6.7E combination with fluid fertilizers.

IMPREGNATION ON DRY BULK FERTILIZERS

Dry bulk fertilizers (except single superphosphate, treble superphosphate and nitrate fertilizers) may be impregnated or coated with ATRAZINE 5L/butylate 6.7E. However, uniform distribution of ATRAZINE 5L/butylate 6.7E on fertilizer particles and uniform application are necessary to assure good results. See Appendix IV on butylate 6.7E label for information and directions for impregnation and use for these combinations.

APPLICATION AND SOIL INCORPORATION DIRECTIONS

SPRAY APPLICATION AND MECHANICAL INCORPORATION

Apply the recommended dosage in 10 to 50 gals. of water per acre (use the higher gallonage in arid areas) using a properly calibrated, low-pressure type nozzle (flood, fan or swirl chamber) on boom-type sprayer having good, but not excessive agitation. Check frequently during application to be sure nozzles are free from clogging and are delivering a uniform spray pattern. Read directions on this label concerning nozzle and screen sizes.

The herbicides must be incorporated into the soil immediately after application to prevent loss of butylate 6.7E. Whenever possible, application and incorporation should be done in the same operation. Apply the herbicides to soil dry enough to permit good soil mixing.

See Appendix I on butylate 6.7E label for Soil Mixing (incorporation) before Planting Directions.

Loss of weed control will result from any delay in incorporation if butylate 6.7E is applied to a moist soil surface.

If there must be a delay (less than one hour) in incorporation of herbicides when application and incorporation are done in separate operations, butylate 6.7E must be applied to a dry soil surface (at least 1/2 inch deep) and free from dew and incidental moisture.

IMPORTANT: Thorough mixing of butylate 6.7E in the soil is necessary. The soil must be in good tilth. Incorporation should be done as soon as possible after application. Use directions for control.

BEST AVAILABLE COPY

PRECAUTIONS FOR ALL APPLICATIONS TO CORN

- 1. To avoid crop injury and illegal residues do not apply more than 4.8 pts. per acre of ATRAZINE 5L to corn in any one year.
- 2. Following harvest of a treated crop, plow (moldboard or disk-plow) and thoroughly till the soil in the fall or spring to minimize possible injury to rotational spring-seeded crops, regardless of the rate used.
- 3. Do not graze treated area or feed treated forage to livestock for 21 days following application or illegal residues may result.

For post-emergence applications plus emulsifiable oil or oil concentrate, see additional precautions under that section of this label.

SUGGESTIONS FOR ROTATIONAL CROPS

- 1. Land treated with ATRAZINE 5L should not be planted to any crop except corn or sorghum until the following year or injury may occur.
- 2. If ATRAZINE 5L is applied after June 10, do not rotate with crops other than corn or sorghum the next year or injury may occur.
- 3. If the broadcast rate or equivalent band application rate exceeds 4.8 pt./A of Atrazine 5L plus simazine 4L (or equivalent of another formulation), a crop of untreated corn or sorghum should precede the next rotational crop.
- 4. In the High Plains and Intermountain area of the West where rainfall is sparse and erratic or where irrigation is required, use Atrazine 4L to control weeds in corn only when corn is to follow corn or a crop of untreated corn, or sorghum is to precede other rotational crops.
- 5. In Western Minnesota and Eastern parts of the Dakotas, Nebraska and Kansas, corn treated with ATRAZINE 5L should not be followed with soybeans if the broadcast rate applied was more than 3.2 pts. per acre (or comparable rate in a band) or injury may occur.
- 6. Injury may occur to soybeans planted in North Central Iowa and South Central Minnesota the year following an ATRAZINE 5L application on Harpe, Canisteo, Stroden or other soils having a calcareous surface layer.
- 7. Do not plant sunflowers, tobacco, vegetables (including dry beans), spring-seeded small grain or small-seeded

BEST AVAILABLE COPY

legumes and grasses the year following ATRAZINE 5L application or injury may occur.

SORGHUM AND SORGHUM-SUDAN HYBRIDS (GRAIN AND FORAGE TYPES)

Apply before planting, at planting, or after planting. See Corn section for weeds control.

Preplant Surface-Applied (Broadleaf and grass control): Use on medium-and-fine-textured soil with minimum-tillage or no-tillage systems only in CO, IL, IN, IA, KS, KY, MN, MO, MT, NE, ND, SD, WI, WY. Apply 2/3 the recommended rate of Atrazine shown in Table 9 for the soil texture as a split treatment 30-45 days prior to planting and the remainder at planting. Applications made less than 30 days prior to planting may be as either a split or single treatment. Under dry conditions, irrigation after application is recommended to move Atrazine into the soil.

If weeds are present at time of treatment, apply in a tank mix combination with a contact herbicide (for example, paraquat or glyphosate). Observe directions for use precautions and restrictions on the label of the contact herbicide.

NOTE: To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

PREPLANT INCORPORATED (BROADLEAF AND GRASS CONTROL)

Broadcast in spring after plowing at rate in Table 9. Apply before, during, or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation. For best results, apply within two weeks prior to planting.

PRE-EMERGENCE (BROADLEAF AND GRASS CONTROL)

Apply during or shortly after planting, but prior to weed or crop emergence at rate in Table 9.




TABLE 9
Preplant Surface, Preplant Incorporated,
and Pre-emergence Applications*
For Broadleaf and Grass Weed Control in Sorghum

Soil Texture	Organic Matter	Broadcast rate/ATRAZINE 5L
COARSE		
Sand, loamy sandy, sandy loam	any level	DO NOT USE (except on pre-emergence use on bedded sorghum in AZ and CA as indicated below)
MEDIUM AND FINE		
Loam, silt loam, silt, clay loam, sandy clay loam, silt clay loam, sandy clay, silty clay, clay	less than 1% 1-1.5% more than 1.5%	2.6 to 3.2 pts. 3.2 to 3.8 pts.

*Do not apply preplant in AL, AR, FL, GA, LA, MS, NC, NM, OK, SC, TN, or TX. Do not apply pre-emergence in NM, OK, or TX, except in northeast OK and the TX Gulf Coast.

In case of planting failures, sorghum can be replanted into soil previously treated with ATRAZINE 5L. Do not make a second broadcast application or injury may occur. If Atrazine 80W is applied in a band and sorghum is replanted in the untreated row middles, ATRAZINE 5L can be applied in a band to the second planting.

PRE-EMERGENCE BROADLEAF WEED CONTROL IN FURROW IRRIGATED BEDDED SORGHUM (AZ and CA only):

For pre-emergence control of broadleaf weeds such as ground cherry, lambsquarters, morning-glory, mustard, pigweed and purslane, broadcast 1.3 to 1.9 pts. per acre. Use the lower rate on coarse-textured soils and soils low in organic matter and use the high rate on fine-textured soils and soils high in organic matter. Make application after bed preparation, during or after planting, but before sorghum and weeds have emerged and before the first furrow irrigation. Several irrigations should follow the application, making sure that all soil is thoroughly wet.

PRECAUTIONS FOR PRE-EMERGENCE APPLICATIONS OF ATRAZINE 5L TO FURROW IRRIGATED BEDDED SORGHUM GROWN IN AZ AND CA:

To avoid possible sorghum injury, do not use on sand or loamy sand soils or on sorghum planted in the furrow. Additionally, applications made to sorghum growing on alkali soils or where cuts, fills or erosions have exposed calcareous or alkali subsoils, may result in crop injury. In case of crop failure, do not replant sorghum for eight months following application. Corn may be planted immediately.

POST-EMERGENCE (BROADLEAF AND GRASS WEED CONTROL):

Apply before weeds exceed 1.5 inches in height at the rate indicated in Table 10. Application may be made up the sorghum reaching 12 inches in height.

TABLE 10

Post emergence Broadleaf and Grass Weed Control in Sorghum

Soil Texture	Minimum Height of Sorghum at Treatment	Broadcast Rate/ATRAZINE 5L
Sandy or loamy sand		DO NOT USE
Sandy loam		See directions for broadleaf and weed control below.
Silt loam to sandy clay loams	Completely emerged	3.2 to 3.8 pts.
Ulton and Pullman clay soils	At least 6 in. high	3.2 to 3.8 pts.
Silty clay loams and heavier soils	Completely emerged	4.8 pts.

POST-EMERGENCE BROADLEAF WEED CONTROL WITH ATRAZINE 5L PLUS EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER:

Broadcast 1.7 pts. per acre for control of broadleaf weeds such as annual morning-glory, cocklebur, lambsquarters, mustard, pigweed, ragweed, smartweed and wild buckwheat. Application should be made before pigweed and lambsquarters reach six inches in height and before all other weeds reach four inches in height. In CO, western KS, NM, Ok, TX, and desert regions of AZ and CA, apply when sorghum is 6-10 inches in height, but before it reaches boot stage. In all other areas, apply after sorghum reaches the three-leaf stage. Add emulsifiable oil at the rate of 1 gal. per acre for ground applications and 0.5 gal. per acre for aerial.

BEST AVAILABLE COPY

applications, or add 1 qt. per acre of oil concentrate for ground application. A cultivation may be necessary if all weeds are not controlled or if regrowth of weeds occurs.

For the list of weeds controlled, see ATRAZINE 5L Applied Alone - Corn or Grain Sorghum - Post-emergence with Emulsifiable Oil or Oil Concentrate in Water.

PRECAUTIONS FOR APPLICATION OF ATRAZINE 5L PLUS EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER TO SORGHUM - See "Precautions for applications with emulsifiable oil or oil concentrate in water" in Corn section.

POST-EMERGENCE BROADLEAF WEED CONTROL WITH ATRAZINE 5L PLUS SURFACTANT (CO, western KS, NM, OK, TX, and desert regions of AZ and CA only):

Broadcast 1.9 pts. of ATRAZINE 5L plus .75-1.50 pts. of surfactant per acre after sorghum reaches 6 inches in height, but before weeds reach 1.5 inches in height. Apply only on sandy loams and finer-textured soils.

TANK MIXTURES FOR GRAIN SORGHUM


ATRAZINE 5L + METOLACHLOR
on Grain Sorghum (Cyometrinil-Treated Seed)

The ATRAZINE 5L + metolachlor tank mixture provide control of weeds listed on this label under the ATRAZINE 5L + metolachlor on corn tank mix section.

Apply ATRAZINE 5L + metolachlor pre-plant incorporated or pre-emergence using the appropriate rates from Table 12 Preplant Incorporated or Pre-emergence. Follow the use instructions on the metolachlor label immediately preceding Table 12.

Table 12: ATRAZINE 5L + Metolachlor - Grain Sorghum*

Soil Texture	Broadcast rates per acre	
	Less than 1.5% organic matter	1.5% organic matter or greater
	ATRAZINE 5L + Metolachlor	ATRAZINE 5L + Metolachlor
Coarse sand, loamy sand, sandy loam	60-80 fl. oz.	60-80 fl. oz.



Medium: loam, silt, silt loam	Do Not Use	1.9 pts. 5L + 1.5 pts.
Fine: silty clay loam, sandy clay loam, silty clay sandy clay, clay loam, clay	1.9 pts. 5L + 1.5 pts.	2.3 to 2.6 pts. 5L + 1.75-2 pts.

*Do not use in NM, OK, or TX except in northeast OK and the Texas Gulf Coast areas. Do not apply preplant incorporated in AZ or the Imperial Valley in CA.


Precautions:

1. Applications of Atrazine + metolachlor on highly alkaline soils or on eroded areas where calcareous subsoils are exposed may cause sorghum injury.
2. If sorghum seed is not properly treated with Cyometrinil, Atrazine + metolachlor will severely injure the crop.
3. Under high soil moisture conditions prior to sorghum emergence, injury may occur following the use of Atrazine + metolachlor. The crop will normally outgrow this effect.
4. Do not use Atrazine + metolachlor on sorghum grown under dry mulch tillage or injury may occur.

Rotational Crops: Refer to the crop rotation instructions on the metolachlor label for metolachlor + Atrazine tank mixtures on corn.

PRECAUTIONS FOR ALL APPLICATIONS TO SORGHUM

1. Heavy rains immediately following application tend to result in excessive concentrations of herbicide in seed furrow, resulting in possible crop injury. Applications to furrow-planted sorghum should not be made until furrows are leveled (Flow-in). Deep planter marks or seed furrows should also be leveled before application.
2. Application made to sorghum growing under stress caused by minor element deficiency or to sorghum growing on highly calcareous soils may result in crop injury.
3. Do not graze or feed forage from treated areas for 21 days following application or illegal residues may result.
4. Following harvest of a treated crop, plow (coldblood or disk plow) and thoroughly till the soil in the fall or



spring to minimize possible injury to rotational spring-seeded crops, regardless of rate used.

5. Injury may occur if both this herbicide, preplant surface, preplant incorporated, or pre-emergence, and an at-planting systemic insecticide applied in-furrow are used.

For applications to furrow-irrigated bedded sorghum in Arizona and California and for post-emergence applications

plus emulsifiable oil, see additional precautions under these sections of this label.

Suggestions for Rotational Crops:

See "Suggestions for Rotational Crops" at the end of Corn Section.

WINTER WEED CONTROL IN TEXAS

For post emergence control of winter weeds only, such as henbit, seedling dock, and annual thistle on fall bedded land in the Gulf Coast and Blacklands of Texas, apply 1.2-1.6 pts./A post-emergence to the weeds in November or December to land that will be planted to corn, grain sorghum or forage sorghum the following spring. For best results add a suitable surfactant such as Surf Ac 820 at the rate of 0.5% of the spray volume, an emulsifiable oil at the rate of 1.0% of the spray volume, or an oil concentrate at the rate of 1 qt per acre.

Normal weed control programs may be used in the following corn, grain sorghum or forage sorghum crop.

Note: Do not plant any crops except corn, grain sorghum, or forage sorghum the spring following this treatment, or illegal residues may result.

CHEMICAL FALLOW ATRAZINE 5L ALONE

WHEAT-SORGHUM-FALLOW

This treatment controls annual broadleaf and grass weeds following wheat harvest and in the following sorghum crop which grows under summer fallow.

Apply 0.50 pt. to one acre immediately following wheat harvest. If weeds are present, remove them with a power plow or other suitable implement after application or use an approved contact herbicide such as paraquat or other application of ATRAZINE 5L. If weeds are present, apply 0.50 pt. to one acre immediately following wheat harvest. If weeds are present, remove them with a power plow or other suitable implement after application or use an approved contact herbicide such as paraquat or other application of ATRAZINE 5L.

BEST AVAILABLE COPY

present at planting, remove them with a sweep plow or other suitable implement before planting.

For the list of weeds controlled, see ATRAZINE 5L Applied Alone - Corn or Grain Sorghum - Preplant Surface-Applied, Preplant Incorporated, or Pre-emergence.

PRECAUTIONS:

1. Use only on silt loam or finer textured soil.
2. Wheat-sorghum-fallow cropping sequence must be followed.
3. Do not apply following sorghum harvest.

NOTE: To avoid illegal residues, do not graze or feed forage from treated area to livestock. To avoid illegal residues and crop injury, do not plant any crop other than those on this label within 18 months following treatment.

WHEAT-CORN-FALLOW (KS, NE): This product controls cheatgrass (downy brome, chess), kochia, mustards, pigweed, Russian thistle, wild lettuce, wild sunflower, and volunteer wheat during period after wheat harvest. Weed control may extend into following corn crop grown under minimum tillage.

Follow directions for use, notes, and precautions in the "Wheat-Sorghum-Fallow" section above, substituting corn for references to sorghum.

WHEAT-FALLOW-WHEAT (CO, KS, MT, NE, ND, SD, and WY): For pre-emergence and control of cheatgrass (downy brome, chess), common lambsquarters, field pennycress, kochia, mustard, Russian thistle, wild lettuce, and suppression of volunteer wheat during fallow period of a wheat-fallow-wheat rotation, broadcast .8-1.6 pt/A on all soils except those listed under "Precautions". For control of pigweed and wild sunflower, use the higher rate. Apply to stubble ground. Treat only once during same fallow period.

TANK MIXTURES FOR CHEMICAL FALLOW

WHEAT-SORGHUM-FALLOW OR WHEAT-CORN-FALLOW (KS, NE)

ATRAZINE 5L PLUS PARQUAT

If weeds are present at application, a tank mix with paraquat may be used. Broadcast 1.2-1.6 pt. of this product plus 1-2 pt. of paraquat in 20-60 gals. of water/A by ground equipment. Add 0.5-1 pt. of a non-ionic surfactant, such as Surf 62 (20% non-ionic) or 1 qt. of spray mixture. Add the paraquat to spray tank first and thoroughly mix before then add atrazine followed by surfactant. Use the 1.2 pt.

BEST AVAILABLE COPY

rate of paraquat if weeds are 4-6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground. Treat only once during same fallow period. Refer to paraquat label for further directions, precautions, and limitations.

If weeds are present at application and this product is used alone, use either an approved contact herbicide before or after treatment, or tillage after treatment.

Use tillage to control weeds which escape during fallow period. Till before planting. For this product applied alone or in tank mixture with paraquat, plant at least 2 inches deep and 12 months or more after application.

PRECAUTIONS:


1. Do not use on sand soil.
2. Do not treat eroded hillsides, caliche and rocky outcroppings, or exposed calcareous subsoil.
3. Do not treat soils of the Rosebud and Canyon Series in western NE and adjoining counties in CO and WY.
4. Do not treat soils with calcareous surface layers.
5. Avoid spray overlap.

NOTE: Do not graze treated areas within 6 months after application or illegal residues may result.

AERIAL APPLICATION: In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft. using low drift nozzles at a maximum pressure of 40 psi. and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply Atrazine 4L alone by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

ROADSIDES

To control certain annual weeds in established perennial grasses along roadsides in CO, KS, MI, NE, ND, SD, and WY, including cheatgrass (downy brome, chess), common (annual) broomweed, little barley, medusahead, safflower, and tumble mustard, broadcast 1.5 pt/A in a mixture of 10 gallons of water by ground equipment in the fall before ground freezes or after thawing in the spring, but before the established grasses green up and before weeds emerge. Examples of desirable established grasses include big bluestem, bluegrass, creosote, buffalograss, crested wheatgrass, Indiangrass, little bluestem, and wild prairie, and timothy and Kentucky bluegrass. Apply and use as directed.



Temporary discoloration or other form of injury to the desirable perennial grasses may occur following application.

NOTE: To avoid illegal residues:

1. Do not cut or feed roadside grass hay
2. Do not allow livestock to graze treated areas.

SUGAR CANE

For control of many broadleaf and grass weeds, such as crabgrass, jungle rice, wire grass, foxtail, amaranths, Flora's paintbrush, fireweed and similar plants. Broadcast 1.6-3.2 qts./A at time of planting or ratooning, but before the cane emerges. Broadcast aerially in a minimum of 5 gal. of spray per acre, or broadcast or band by ground equipment in a minimum of 20 gal. per acre, unless indicated otherwise. One additional application may be made over the cane as it emerges, and two additional applications may be made interline after emergence, as a directed spray. Repeat treatments, where needed, may be applied broadcast, band, or interline as suggested with the final application being prior to "close-in." Do not exceed the rate of herbicide suggested for any one crop of sugarcane.

NOTE: Where high rates of Atrazine are used alone, apply in a minimum of 1 qt. of water for each .8 qts. of ATRAZINE 5L applied per acre.

AERIAL APPLICATION: In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft. using low drift nozzles at a maximum pressure of 40 psi. and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply ATRAZINE 5L alone by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

FLORIDA

For control of emerged Pellitory Weeds (artillery weed). Apply .6-1.0 pts/A in 40 gals of water per acre as a directed spray, by ground equipment prior to "close-in." Add 2 qts. of surfactant for each 50 gals of spray and be sure weed foliage is thoroughly covered.

For control of early emerging grasses, large crabgrass, pellitory (artillery) weeds, and spiny amaranth, use one of the following methods of planting or ratooning:

1. Apply seed of a grass or grasses. Follow with one or two applications, as needed, per label directions suggested and



weeds, at 3.2 pt/A. Treat before weeds exceed 1.5 inches in height.

2. Apply 1-3 times, as needed, at 3.2 pt/A post-emergence to sugarcane and weeds. Treat before weeds exceed 1.5 inches in height.

LOUISIANA

For control of annual weeds during the summer fallow period, apply 3.2 pt/A to weedfree beds immediately after bed formation. Follow normal weed control after planting.

Precautions: To avoid crop injury:

1. Do not apply more than 8 qts. to any one crop of sugarcane.
2. If making 3.2 pt/A application during summer fallow period do not exceed 6.4 qts. during the remainder of the growing season, or illegal residues may result.

TEXAS

Use ATRAZINE 5L for control of barnyardgrass, pigweed, purslane and sunflower, in plant or ratoon sugarcane.

Apply 6.4 pt/A of ATRAZINE 5L pre-emergence. Follow with one or two applications, as needed, at 4.8 pt/A post-emergence to sugarcane and weeds.

For best results when weeds are emerged, add a nonionic surfactant, such as Surf Ac 820 at a concentration of 2 qts/100 gals to the spray and apply before weeds exceed 1.5 inches in height.

PRECAUTIONS:

1. Do not apply ATRAZINE 5L after "close-in."
2. Do not apply more than 16 pts. of ATRAZINE 5L to any one crop of cane.
3. Injury to sugarcane may occur when under moisture stress, when soil is of low absorptive capacity, or when land is first cropped to sugarcane.

COFFEE

For control of annual broadleaf and grass weeds prior to transplanting, after transplanting or in established controls including Douglas fir, grand fir, noble fir, white fir, Austrian pine, larch pine, Jeffrey pine, loblolly pine, slash pine, longleaf pine, shortleaf pine, white pine, yellow pine, red pine, Scotch pine, blue pine, loblolly pine,

BEST AVAILABLE COPY

and Sitka spruce): Broadcast 3.2-6.4 pts. in a minimum to 5 gals of water per acre by air or 10 gals by ground before weeds are 1.5 inches tall. Apply to established trees between fall and early spring while trees are dormant. For applications prior to transplanting allow sufficient precipitation to activate Atrazine before transplanting. In areas where spring and summer rainfall is inadequate to activate Atrazine, apply during fall prior to spring transplanting.

For the list of weeds controlled, see ATRAZINE 5L Applied Alone - Corn or Grain Sorghum - Preplant Surface Applied, Preplant Incorporated or Pre-emergence.

QUACKGRASS CONTROL: Broadcast 6.4 pt in a minimum of 5 gals of water per acre by air or 10 gals by ground between fall and early spring while trees are dormant and before quackgrass is more than 1.5 inches tall.

PRECAUTIONS: Temporary injury to trees may occur following the use of Atrazine on coarse-textured soil. To avoid injury:

- 1. Do not apply to seedbeds
- 2. Apply only once per year.

AERIAL APPLICATION: In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft. above vegetation, using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply ATRAZINE 5L by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

PERENNIAL RYEGRASS GROWN FOR SEED (PACIFIC NORTHWEST ONLY)

For the control of annual broadleaf and grass weeds such as annual ryegrass, rattail fescue, and similar species. Broadcast 1.7 pints per acre, after first fall rain.

PRECAUTIONS: (To Avoid Crop Injury & Illegal Residues)

- 1. Apply only to perennial ryegrass stands from which a seed crop has been harvested.
- 2. Do not apply Atrazine to ryegrass more than 2 years in succession.
- 3. Do not graze treated land within 10 days of application.
- 4. Do not plant treated land to the other crop for at least 12 months.



NONSELECTIVE WEED CONTROL ON NONCROP LAND

For use on fence rows, around utility poles, and industrial sites, highway medians and shoulders, railroad right-of-way, lumberyards, petroleum tank farms, and in noncrop areas on farms, such as around buildings, equipment, and fuel storage areas, along fence and lanes. Do not use near adjacent desirable plants or in greenhouses, or injury may occur. Apply ATRAZINE 5L at rates recommended below before or soon after weeds begin growth. Post-emergence applications should be made when weeds are young and actively growing. Use higher rates on heavy clay and muck soils. Do not contaminate domestic or irrigation water supplies, or lakes, streams, or ponds. Use sufficient water to assure thorough coverage. Use at least 1 gal. of water for each quart of ATRAZINE 5L, more if practical. Mechanical by-pass or jet agitation is necessary to keep ATRAZINE 5L in suspension during application.

Note: Do not allow livestock to graze treated areas, or illegal residues may result.

Annual broadleaf and grass weeds (including barnyardgrass, cheatgrass, crabgrass, lambsquarters, foxtail, ragweed, puncturevine, and turkey mullein): Broadcast 3.8 to 8 qts/A.

Hard-to-kill annual and many perennial broadleaf and grass weeds (including bluegrass, burdock, Canada thistle, dogfennel, orchardgrass, plantain, quackgrass, purpletop, redtop, and smooth brome): Broadcast 8 qt./A.

Hard-to-kill biennial and perennial weeds (including bullthistle and sowthistle): Broadcast 8 qt/A.

For longer residual control in regions of high rainfall and a long growing season, broadcast 8 qt/A.

For small areas, 3.6 oz. per 1,000 sq. ft is equivalent to 8 qt/A.

ST AUGUSTINE GRASS, CENTIPEDE GRASS, AND DORMANT BERMUDA GRASS

ATRAZINE 5L controls spurweed, Florida betony, annual bluegrass, and many other problem weeds in home lawns, ornamental, and recreational turf and other noncrop areas such as highway right-of-way and similar areas. ATRAZINE 5L may be applied with any pump-up or compressed air type sprayer or with a boom or type sprayer. ATRAZINE 5L will control BOTH existing weeds and weeds from seeds. It is of water and can be applied during or after rain. The effectiveness is increased by soil moisture. However, the control of

BEST AVAILABLE COPY

weeds from seeds, rainfall or watering is necessary within 7 to 10 days after treatment.

CAUTION: Do not apply within the active root zone of vegetables or desirable ornamental plants such as camellias, azaleas, boxwoods, etc. However, ATRAZINE 5L treatments should not normally cause injury to medium or large shrubs or trees in the landscape. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result.

WEEDS CONTROLLED OR SUPRESSED

Annual Bluegrass (Poa Annua), Chickweed (Common and Mouseear), Crabgrass, Cransbill, Cudweed, Dichondra, Florida Betony, Henbit, Knotweed, Lespedeza, Moneywort, Mustards, Narrowleaf Vetch, Parsley-Piert, Sandspur, Smutgrass, Spurge, Spurweed, Swinecrest, Woodsorrel and various annual clovers.

TIMING OF APPLICATION

The timing of applications to achieve maximum control may vary quite a bit with different weed species. The following application times are suggested for certain problem weeds.

Spurweed: The best control of spurweed can be obtained by applying ATRAZINE 5L when spurweed has emerged (December and January).

Florida Betony: This weed emerges in the fall so an application of ATRAZINE 5L in mid to late October followed with a second application in mid to late February would give best control.

Dichondra, Moneywort: The best control of these weeds can be obtained by apply ATRAZINE 5L in early April followed with a second application in July. Do not apply to growing Bermuda Grass.

Applications for Spurweed or Florida Betony generally will give control or suppression of the other weeds listed. However, as a general rule ATRAZINE 5L will give the best control when applied to young tender weeds or just prior to weed emergence.

SO, ABOUT THE USE OF ATRAZINE 5L

ATRAZINE 5L can be applied to established SL, from time and Centipede grasses during both the dormant and the growing season. Best results are generally obtained in the early spring or dormant period, when weeds are small or newly emerged. Follow rates below.

BEST AVAILABLE

DORMANT BERMUDA GRASS

ATRAZINE 5L may be applied to Bermuda grass during the dormant season only. Applications to Bermuda grass during the growing season will cause injury. Follow rates below.

RATE OF APPLICATION

Determine the total area to be sprayed and base rate of application on the chart below. Avoid overlapping spray pattern while treating. Shake contents well before mixing.

Area to Be Treated	Amount of ATRAZINE 5L	Suggested Amount of Water
500 Sq. ft.	.6 oz.	1 gallon
1,000 " "	1.2 oz.	2 gallons
3,000 " "	3.6 oz.	8 gallons
5,450 " "	6.4 oz. (.4 pt.)	10 gallons
10,900 " "	12.8 oz. (.8 pt.)	20 gallons
(0.25 acre)		
1 Acre	3.2 pts.	40
gallons		

NOTE: Do not graze or feed turf clippings to animals, or illegal residues may result.

BEST AVAILABLE COPY

TURFGRASS FOR FAIRWAYS, LAWNS, SOD PRODUCTION* AND SIMILAR AREAS

*In states other than Florida. For use on turfgrass for sod in Florida, see "Turfgrass for Sod (Florida only)" section above.

Bermudagrass, Centipedegrass, St. Austustinegrass, and Zoysia Grass.

Apply ATRAZINE 5L after October 1 before emergence of winter annual weeds of control of annual bluegrass, burclover, carpet burweed, chickweed, corn speedwell, henbit, hop clover, and spurweed. ATRAZINE 5L will control annual bluegrass even if it is emerged at time of treatment. For control of summer annual weeds listed in the preemergence portion of the ATRAZINE 5L Applied Alone - Corn or Grain Sorghum section on this label, also apply ATRAZINE 5L in late winter before the weeds emerge. Apply in a minimum of 15 gals. of water per acre or 1 gal. per 1,000 sq. ft.

Where annual bluegrass is the major weed, use 1.6 pts per acre (or .6 oz. per 1,000 sq. ft.) Use 3.2 pts. per acre (or 1.2 oz. per 1,000 sq. ft.) for control of the other weeds named above. Do not exceed 1 qt. per acre per treatment on newly sprigged turfgrass or on hybrid bermudagrass such as Tiflawn, Tifway, or Ormond.

For continued summer annual weed control, apply another 1.6 pts. per acre at least 30 days after the previous application, but not after April 15. However, do not make more than two applications of this product per year.

Precautions: On newly sprigged turfgrass and hybrid bermudagrass, temporary slowing of growth and yellowing may occur following application. To avoid turf injury, 1) Use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases. 2) Do not use on golf greens. 3) Do not use north of NC (except may be used in Virginia Coastal Plains) or west of the high rainfall areas of eastern OK and eastern TX. 4) Do not use on muck or alkaline soils. 5) Do not apply over the rooting area of trees or ornamentals not listed on this label. 6) Do not overseed with desirable turfgrass within 4 months before or 6 months after treatment.

Notes: Do not mow or treat turf clippings to animals, or allow residues to result.

TURF GRASS FOR SOD PRODUCTION

Use only on bermudagrass, centipedegrass, and zoysia grass. Do not use on St. Augustinegrass, St. Augustinegrass, or any other grass not listed on this label.

BEST AVAILABLE COPY

control those weeds listed under Atrazine 5L Applied Alone - Corn or Grain Sorghum - Preplant Surface Applied, or Preplant Incorporated, or Preemergence.

Muck Old Beds: Within 2 days after lifting sod.
 or 6.4 pts.
 Peat New Beds: 3-4 days after sprigging or plugging.

Sandy Old Beds: Within 2 days after lifting sod.
 Soil 3.2 pts.
 New Beds: 7-10 days after sprigging or plugging

If weeds regrow, apply an additional 3.2 pts./A on muck or peat, or 2 pt./A on sandy soil.

PRECAUTIONS: To avoid crop injury

1. Do not apply within 30 days prior to cutting or lifting.
2. Do not apply in combination with surfactants or other spray additives.
3. Use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases.
4. On newly sprigged turfgrass, temporary slowing of growth may follow application.

MACADAMIA NUTS

For preemergence control of many broadleaf and grass weeds, including crabgrass, foxtail, wiregrass, Flora's paintbrush, spanishneedles, and fireweed, broadcast 3.2-6.4 pts./A before harvest and before weeds emerge. Repeat as necessary. Do not spray when nuts are on ground during harvest period. Do not apply by air.


GUAVA

Use only on established plantings which are at least 18 months old. Apply as a directed spray at 3.2-6.4 pts./A ATRAZINE 5L in 20-50 gallons of spray mix preemergence or early post-emergence to weeds. When applying post-emergence, the use of a surfactant and greater spray volume (80-100 gallons of spray mix per acre) may enhance weed control.

This product controls many annual broadleaf and grass weeds, including fireweed, rumsline, scarlet pimpernel, spanishneedles, and scabchistle.

Notes: To avoid illegal residues

1. Do not allow spray to contact foliage or fruit.
2. Do not apply more frequently than at 4 month intervals.
3. Do not apply more than 1.58 pts. atrazine 5L per year.



ESTABLISHED ORCHARDGRASS GROWN FOR SEED (PACIFIC NORTHWEST ONLY)

Atrazine aids in control of annual ryegrass, rattail fescue, and volunteer orchardgrass. Broadcast 3.2 pts/A of ATRAZINE 5L in at least 15 gallons of water to field which has been burned after first fall rain.

PRECAUTIONS:

1. Apply only to orchardgrass from which seed crop has been harvested.
2. Do not apply to orchardgrass more than two years in succession.
3. Some yellowing or crop stunting can occur.

NOTE: To avoid illegal residues.

1. Do not graze treated land within 6 months of application.
2. Do not plant treated land to other crops for 16 months.

CompexR trademark of Kalo Agricultural Chemicals, Inc.
UniterR trademark of Hopkins Agricultural Chemical Company

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE INSTRUCTIONS.

Store in a cool dry location.
Avoid storage at high temperatures.
Keep container tightly sealed.
Avoid contamination with acids or alkalies.
Do not stack more than 2 pallets high to prevent crushing.
Keep containers away from any source of puncture.
Store in original container only.
Pesticides should be separated during storage to prevent cross contamination of other pesticides, fertilizer, food and feed.
Storage area should preferably be locked to prevent admittance by unauthorized or unknowledgeable persons.
If the container is damaged and leaking or material has been spilled follow these procedures:

1. Cover spill with absorbent material.
2. Sweep into disposal container.
3. Wash area with detergent and water and follow with clean water rinse.
4. Do not allow to contaminate water supplies.

BEST AVAILABLE COPY

5. Dispose of according to instructions below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY - CONDITION OF SALE:

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall Drexel or the Seller be liable for consequential, special or indirect damages resulting from the use and/or handling is not not in strict accordance with directions given herewith. The foregoing is a condition of sale by Drexel Chemical Company and is accepted as such by the Buyer.

BEST AVAILABLE COPY