

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with this label.

STORAGE AND DISPOSAL

Do not store near fertilizer, seeds or other pesticides. Do not contaminate water, food or feed by storage or disposal. Emptying is prohibited. Pesticide, spray mixture or rinseate that cannot be used according to label instructions must be disposed of according to Federal, state or local procedures under the Federal, State and Environmental Pesticide Control Act of 1972 (or equivalent) and approved for use in a military landfill, only if approved by the appropriate procedures.

DO NOT apply, allow to drift or drain or flush equipment onto or near croplands, lawns, trees, or other desirable plants or to areas where their roots may extend.

DO NOT allow contact with fertilizers, insecticides, fungicides, or seeds. Application equipment should be thoroughly cleaned with a detergent and water to prevent corrosion. Equipment which has been used to apply this product should not be used for applying insecticides, fungicides, fertilizers, or other agricultural chemicals as plant injury may result.

USE IN INDUSTRIAL PLANTS, LUMBER YARDS, TANK FARMS, STORAGE AREAS AND PARKING LOTS for control of most annual and perennial grasses and broadleaved weeds. The effectiveness and length of time that control will last will depend on type of soil, amount of rain and other conditions.

RATE: Under ordinary conditions use at the rate of 3 gals. per 1000 sq. ft. Under conditions where the weeds are difficult to control, use at the rate of 6 gals. per 1000 sq. ft.

HOW TO APPLY: Use a sprayer which will distribute the required amount evenly over the area to be treated. Use a coarse spray to reduce drift. Application may be made when top growth is dense, but better control will be obtained when weeds are short or when excess growth is mowed and removed before applying.

WHEN TO APPLY: Best results will be obtained if applied shortly before or at the time plant growth begins. Rain or thorough watering after application will cause the material to move to the root zone and thus speed up control.

LIMITED WARRANTY

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

DREXEL DYNAMYTE 3

EPA REGISTRATION NO. 19713-82

Additional Uses for Lentils: ✓

LENTILS: Pre-emergence - Use 1 gal. of DYNAMYTE 3 Herbicide in 20 to 30 gallons of water per acre as a broadcast application after planting but before emergence of crop. Use 1 gal. of DYNAMYTE 3 Herbicide in 5 to 10 gallons of water per acre for aircraft application.

*DON'T
RECOMMEND
DO NOT
PROSE*

ACCEPTED
with comments
APR 8 1983
19713-82