

Apply only when conditions for desiccation are poor.

SUNFLOWER—To facilitate harvest and reduce moisture in the fully mature seed heads, apply 1 gallon Defol 6 in sufficient water to give good coverage 7 or more days before normal harvest when sunflower heads are fully mature. Apply in 5 to 10 gallons of water per acre by air or in 20 to 30 gallons of water per acre by ground with a high clearance sprayer. Do not apply later than 7 days before harvest. Do not graze treated fields or feed treated forage within 14 days of application.

SAFFLOWERS—To facilitate harvest and reduce moisture in the fully mature seed heads, apply 1 gallon of DEFOL 6 in sufficient water to give good coverage 7 or more days before normal harvest when safflower heads are fully mature. Apply in 5 to 10 gallons of water per acre by air or in 20 to 30 gallons of water per acre by ground. Do not apply later than 7 days before harvest. Do not graze treated fields or feed treated forage.

RICE—To facilitate harvest and reduce moisture in the fully mature seed heads and to desiccate immature red rice in mature white rice, apply by aircraft 1 gallon of DEFOL 6 per acre in 5 or more gallons of water per acre. Use the lowest rate when crop conditions and weather are right to enhance rapid drying. Apply 7 to 10 days before anticipated harvest. To avoid drift do not apply in windy weather. Thorough coverage of the crop is essential to achieve good results. Do not graze treated fields or feed treated straw or crop residues within 14 days of application.

SOYBEANS—As a soybean harvest aid to dry weeds in early maturing soybeans and to facilitate the harvest, apply 1 gallon of DEFOL 6 per acre in enough water to give thorough coverage. Application may be made by ground equipment or aircraft. Make application 7 to 14 days before anticipated harvesting date when beans are mature and ready for harvest. For air application use a minimum of 5 gallons ready to apply spray solution per acre. For ground equipment apply a minimum of 20 gallons of ready-to-apply spray solution per acre. Do not graze treated fields or feed treated soybean foliage.

GUAR BEANS—To facilitate harvest and reduce bean seed moisture, apply 1 gallon of DEFOL 6 in enough water to provide thorough coverage either by ground or ground equipment. Make application 7 to 14 days before anticipated harvest date. Do not apply under conditions of extreme heat during the middle of the day. For air application, use 5 to 10 gallons of total solution per acre. For ground application, use a minimum of 20 gallons of total solution per acre.

NOTE: DO NOT graze treated fields or feed treated fodder or forage to livestock.

CHILIPEPPERS—Apply 5 to 25 gallons per acre of 1 to 15 gallons of water by air or 10 to 40 gallons of water by ground rig. To be used on peppers for processing only. Apply 10 days before harvest. When grown for processing, do not harvest until processor before applying.

FLAX—For use as a Harvest Aid. To desiccate trashy weeds and desiccate the flax crop, apply 1 gallon DEFOL 6 per acre in enough water to give thorough coverage. Make application on bright, sunny days (preferably when temperature above 60° F) 7-10 days before anticipated harvest date. For air application use minimum of 5 gallons ready to apply material per acre. Activity of the spray will be enhanced by the addition of good nonionic surfactant at the rate of 1 quart per 100 gallons of spray.

NOTE: DO NOT graze treated fields or feed treated flax foliage and fodder for 14 days after application.

CORN—DEFOL 6 may be used as a harvest aid in early maturing corn. Germination of morning glory and other vines may be erratic. Use 2 gallons DEFOL 6 in 5 to 7 gallons of water per acre by air. Apply at least 14 days before anticipated harvest date on clear days when temperature is expected to go above 70° F. Use the 5 gallon rate when weeds are small and the crop canopy is fairly open.

NOTE: DO NOT graze treated fields or feed fodder, forage or residual seeds within 14 days of application.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE INSTRUCTIONS: Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. Do not stack plastic containers (4 X 1 gal or 2 X 2.5 gal) over 2 pallets high. Do not stack metal containers over 4 pallets high. Move containers by handles or cases. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Store in original containers only. Unloading, storage and work areas must be as clean as possible to prevent contamination if spilled. If concentrate or spray solution is allowed to dry on flammable objects, the chlorate component can act as an oxidizer if the contents are leaking or material is spilled follow these steps:

1. Contain spill, absorb with a material such as saw dust, clay granules or dirt.
2. Collect and place in suitable containers for disposal.
3. Wash area with water and soap to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Clothing contaminated with chlorate solution must be removed at once before solution dries. Dried contaminated clothes can be ignited easily. Wash contaminated clothing immediately.
6. Do not allow run off to enter sewer or contaminate water supplies.
7. Dispose of waste as indicated below.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Metal—Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers—Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY—CONDITION OF SALE:

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. In no case shall Drexel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by Drexel Chemical Company and is accepted as such by the Buyer.



Manufactured By

Drexel Chemical Company

P.O. BOX 1000, MEMPHIS, TENN. 38101

BEST AVAILABLE COPY