

PM 21 19713-87

9-8-97

10/2



ACCEPTED
9-8-97
Under the Federal Insecticide, Fungicide, and Rodenticide Act.
as amended for the pesticide
registered 19713-87

Suffa-8

A Flowable 8 Pound Elemental Sulfur

ACTIVE INGREDIENT:
Elemental sulfur* 65.0%

INERT INGREDIENTS: 35.0%

TOTAL: 100.0%

*This product contains 8.0 lbs. of sulfur per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION
Read Entire Label Before Using This Product
SHAKE WELL BEFORE USING

EPA Reg. No. 19713-87
EPA Est. No. 19713-GA-1 # 575F-0297

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
IF SPLASHED IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes and get medical attention if irritation develops.
IF SPILLED ON SKIN: Wash skin with soap and plenty of water. If irritation develops, send for a physician.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION Harmful if swallowed. Avoid breathing spray mist. Avoid contact with eyes, skin and clothing. May cause irritation of nose, throat and skin. Do not get in eyes, on skin or on clothing.
PERSONAL PROTECTIVE EQUIPMENT (PPE)
 Applicators and other handlers must wear:
 Long-sleeved shirt and long pants, Waterproof gloves, Shoes plus socks, and Protective eyewear.
 Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS
 Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS
 For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL HAZARDS
 Keep away from heat, sparks, or flame.

DIRECTIONS FOR USE
 It is a violation of federal law to use this product in a manner inconsistent with its labeling.
DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
 For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.
 Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.
 Personal Protective Equipment required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, Waterproof gloves, Shoes plus socks and Protective eyewear.

IMPORTANT: Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality.
 During periods of high temperatures sulfur may burn foliage and fruit. Do not make sulfur application at excessively high temperatures. Do not use SUFFA 8 with oil or within four (4) weeks of an oil application.
 When SUFFA 8 is used with arsenicals, lime should be added to prevent plant injury.

Use of this material in a manner other than in accordance with this label may cause serious plant, crop or other personal injury, excessive residues, unsatisfactory control or other unintended consequences.
 Consult your State Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your local conditions.

Unless otherwise specified for specific tree fruits, dosage rates are given as pints of SUFFA 8 per 100 gallons of water for use in a full coverage spray.
 Shake well before using.
 Pour recommended amount into partially filled spray tank. Keep agitator running during filling and spraying operation. Do not allow mixture to stand. Failure to maintain agitation will cause SUFFA 8 to settle and may necessitate manual stirring to redispense.

APPLE, PEAR: Scab - Pre-bloom through calyx sprays - Apply 2 1/4 to 5 1/4 cups Cover sprays - Apply 7/8 to 1 1/2 pints. Powdery mildew - Pre-bloom through calyx sprays - Apply 2 1/2 to 5 1/2 cups. Cover sprays - 1 1/2 pints.
PEACH: Brown rot, Powdery mildew, Scab - Pink and bloom sprays - Apply 2 1/2 to 5 1/2 cups. Petal blt, shuck and cover sprays - Apply 1/2 to 1 1/2 pints.
CHERRY, PLUM, PRUNE: Brown rot, Leaf spot - Pink and bloom sprays - Apply 2 1/2 to 5 1/2 cups. Petal, shuck and cover sprays - Apply 1/2 to 1 1/2 pints.
CITRUS (Grapefruit, lemons, limes, oranges, tangelos and tangerines): Rust/mite Apply 1/2 to 3 pints. Do not apply within 4 weeks of an oil application.
PEANUTS: Rust - Apply 1 1/2 to 3 pints per acre in sufficient water for thorough coverage. Begin application approximately 6 weeks after planting. Repeat at 7 to 10 day intervals as long as rust control is required. Do not apply within 4 weeks of an oil application.

Net Contents: 2 1/2 gals.

Manufactured By
Drexel Chemical Company
 P O BOX 13327 MEMPHIS, TN 38113-0327

2012

COTTON: *Atlantic mite* - Apply 1 1/2 pints for early season control and up to 6 pints for mid to late season control.

GRAPE: (California only) *Powdery mildew* - Apply 1/2 to 1 1/2 pints per 100 gallons water, maximum of 6 pints. SUFFA 8 per acre in Spring and Summer. Make first application when shoots are 6 to 8 inches long and third at or about bloom stage. The timing, rate and number of applications will vary with local conditions. Consult State Agricultural Service representative or State Agricultural Experiment Station in your region for specific information. Follow use precautions given on this label.

POTATOES: *Powdery mildew* - Use 2 1/2 to 3 pints in 8 gallons of water per acre by air. Apply at first sign of infection and repeat at 3 week intervals.

STRAWBERRIES: *POWDERY mildew* - Use 1/2 pint. Apply at first sign of infection and repeat at 3 week intervals. For strawberries that will be canned, discontinue application of sulfur well before harvest in accordance with local canner's recommendations.

SUGAR BEETS: *Powdery mildew* - Use 1/2 to 2 1/2 pints in 5 to 8 gallons of water per acre by air. Apply at first sign of infection and repeat at 3 week intervals.

TOMATOES: *Tomato russet mite* - Use 3 to 8 pints per acre depending upon size of plants. Use full coverage spray.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE INSTRUCTIONS: Storage should be under lock and key in a ventilated room and secure from access by unauthorized persons and children. Storage should be in a cool, dry area away from any heat or ignition source. Do not stack plastic containers over 2 pallets high. Move containers by hand or with a pallet fork. Do not move containers from one area to another unless they are securely sealed. Keep containers tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Store in original containers only. If the contents are leaking or material is spilled follow these steps:

1. Contain spill, absorb with a material such as saw dust, clay granules or dirt.
2. Collect and place in suitable container for disposal.
3. Wash area with water and soap to remove remaining pesticide.
4. Follow washing with clean water rinses.
5. Place a leaking container in a plastic tub and transfer contents as soon as possible to any empty original container.
6. Do not allow run off to enter sewer or contaminate water supplies.
7. Dispose of waste as indicated below.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or residue is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY - CONDITION OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall Drexel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by Drexel Chemical Company and is accepted as such by the Buyer.