

CROPS	INSECT	TSPS OF DREXEL CARBARYL PL/GAL	PRE- HARVEST INTERVAL (DAYS)	SPECIFIC DIRECTIONS
All Vegetable Crop in This Section	Grasshoppers	2 to 6	See specific Vegetable Crop	Use 2 to 3 teaspoonfuls/ gallon for nymphs on small plants or sparse vegetation. Use 4 to 6 teaspoonfuls/gallon for adult grasshoppers or application to dense vege- tation.
Asparagus	Asparagus beetle Apache cicada Asparagus beetle	4 to 8 8 to 16	1 Post harvest application only	Treat ferns or brush growth. Do not treat more than once every 3 days.
Beans (including black-eyed peas, cowpeas, crowder or southern peas, dry beans, green beans, lima beans, navy beans and snap beans)	Blister beetles Mexican bean beetle Alfalfa caterpillar, Bean leaf beetle, Cucum- ber beetle, Flea beetles, Green cloverworm, Japanese beetle, Leafhoppers, Three cornered alfalfa hopper, Thrips, Velvetbean cater- pillar, Western bean cutworm	2 to 4 4	0	Observe plant response precautions.
	Armyworm, Corn earworm, Cut- worms, European corn borer, Fall armyworm, Stink bugs, Tarnished plant bug, Webworms	4 to 6		
	Cowpea curculio	8		
	Corn earworm, Lirabean pod borer, Lygus bugs, Stink bugs	8		

CALIFORNIA ONLY