

#### UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460per

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Ms. Luz Piwonka

Drexel Chemical Company
P.O. Box 13327

Memphis, TN 38113-0327

JUL 2 6 2001

Dear Ms. Piwonka:

Subject: Amendment - Delete Beans, Celery, Cucumber, Parsley,
Peas, Peppers, and Potatoes per Diazinon MOA, and
Replacement CSF per Diazinon MOA

Drexel Diazinon 14G [New Name]

Drexel D-264 14G [Old Name]

EPA Registration Number 19713-95

Your submissions dated May 8 and May 17, 2001

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable provided that you:

- Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
- 2. Make the labeling change indicated below before you release the product for shipment bearing the amended labeling:
  - a. In the First Aid section, re-add the **If inhaled** subheading and updated advice appropriate to this product in accordance with PR Notice 2001-1.

3. Submit four (4) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. The Agency's records will be adjusted to reflect the change in product name as follows:

Drexel Diazinon 14G [New Name]
Drexel D-264 14G [Old Name]

The replacement Basic Confidential Statement of Formula (CSF) is acceptable and has been add to this product's registration file.

Sincerely,

George T. LaRocca
Product Manager 13
Insecticide Branch
Registration Division (7505C)

# RESTRICTED USE PESTICIDE DUE TO AVIAN AND AQUATIC TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.



# D-26414G

Diazinon Granular Insecticide

For use outdoors to control certain soil insects including wireworms, cutworms, and other crop pests

50 Pounds Net Weight

Active Ingredient:

Diazinon: O,O-Diethyl O -(2-isopropyl-6-

Total:

100.0%

100.0%

# KEEP OUT OF REACH OF CHILDREN CAUTION

EPA Reg. No. 19713-95 EPA Est. No. 19713-GA-1 ACCEPTED
with COMMENTS
In EPA Letter Dated:
JUL 2 6 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the posticide registered under EPA Reg. No.

change name to:
"Diazinon 146"



#### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Drexe Chemical Company or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Drexe and Seller harmless for any claims relating to such factors.

Drexe . warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or makes no warranty and User assume the risk of any such use. Drexe MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

i

In no event shall Dresco. or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF Dresco. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF DIESE. OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Drese and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of Drese.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Reformulation or repacking of this product is prohibited.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

Unless directions under **Timing/Special Directions** state otherwise, applications to the soil should be worked into the top few inches immediately after application with a rotary hoe, cultivator, disk, harrow, or other suitable means for greatest effectiveness.

#### **WORK SAFETY RULES**

1

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

When handling Diazinon 14G, do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See First Aid section of this label.

#### **VEGETABLE CROPS**

#### SPECIAL PRECAUTIONS TO REDUCE RISKS TO BIRDS

Birds feeding on treated areas may be killed. To reduce risks to birds:

- 1. Thoroughly incorporate this product into the soil during or immediately after application. Refer to the Soil incorporation table that follows.
- 2. Cover or incorporate spilled granules.

diazinon 14G should be applied uniformly as directed for each crop. When a range of rates is given, use the higher rate for heavy infestations.

For application by ground equipment only. Do not apply this product in greenhouses.

Note: Soil Incorporation - Unless directions under Timing/Special Directions state otherwise, for control of pests listed below Biazinon 14G should be incorporated into the soil during or immediately after application.

## Soil Incorporation

ž

| Depth of Incorporation |
|------------------------|
| 2-3 inches             |
| 3-6 inches             |
| 1-2 inches             |
| 3-4 inches             |
| 4-8 inches             |
|                        |

<sup>\*</sup>If cutworms cannot be identified, incorporate 4-6 inches.

| Crop                  | Pest             | Rate         | Timing/Special Directions   |
|-----------------------|------------------|--------------|---|
| Beets, Red<br>(Table) | Cutworms         | 14-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|                       | Mole Crickets    | 7 lbs./A     | One to two days before planting, broadcast and immediately incorporate into soil. |
|                       | Wireworms        | 21-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |
| Broccoli              | Root<br>Maggots  | 14-21 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|                       | Cutworms         | 14-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|                       | Mole<br>Crickets | 7 lbs./A     | One to two days before planting, broadcast and immediately incorporate into soil. |
|                       | Wireworms        | 21-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |

ţ

| Crop                | Pest             | Rate         | Timing/Special Directions   |
|---------------------|------------------|--------------|---|
|                     |                  |              |   |
| Brussels<br>Sprouts | Root<br>Maggots  | 14-21 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|                     | Cutworms         | 14-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|                     | Mole<br>Crickets | 7 lbs./A     | One to two days before planting, broadcast and immediately incorporate into soil. |
|                     | Wireworms        | 21-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |
| Cabbage             | Root<br>Maggots  | 14-21 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|                     | Cutworms         | 14-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|                     | Mole<br>Crickets | 7 lbs./A     | One to two days before planting, broadcast and immediately incorporate into soil. |
|                     | Wireworms        | 21-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |

| Crop        | Pest             | Rate         | Timing/Special Directions   |
|-------------|------------------|--------------|---|
| Carrots     | Root<br>Maggots  | 14-21 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|             | Cutworms         | 14-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|             | Mole<br>Crickets | 7 lbs./A     | One to two days before planting, broadcast and immediately incorporate into soil. |
|             | Wireworms        | 21-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |
| Cauliflower | Root<br>Maggots  | 14-21 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|             | Cutworms         | 14-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|             | Mole<br>Crickets | 7 lbs./A     | One to two days before planting, broadcast and immediately incorporate into soil. |
|             | Wireworms        | 21-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |

| Crop                 | Pest  | Rate         | Timing/Special Directions   |
|----------------------|---|--------------|---|
| Collards             | Cutworms  | 14-28 lbs./A | Broadcast and incorporate into soil just before planting.   |
|                      | Mole<br>Crickets  | 7 lbs./A     | One to two days before planting, broadcast and immediately incorporate into soil.   |
|                      | Wireworms   | 21-28 lbs./A | Broadcast and incorporate into soil just before planting.   |
| Endive<br>(Escarole) | Cutworms  | 14-28 lbs./A | Broadcast and incorporate into soil just before planting.   |
|                      | Mole<br>Crickets  | 7 lbs./A     | One to two days before planting, broadcast and immediately incorporate into soil.   |
|                      | Wireworms   | 21-28 lbs./A | Broadcast and incorporate into soil just before planting.   |
| Ginseng              | Cutworms,<br>Climbing<br>cutworms,<br>Wireworms,<br>White grubs | 14-28 lbs./A | In new beds, broadcast just before planting and incorporate into the top 4-8 inches of soil by tilling with hand-held or powered ground equipment.                    |
|                      |   | 21 lbs./A    | When soil insect damage appears in established beds, broadcast in the Spring or late Summer through early Fall. Incorporate mechanically with 1/2 inch of irrigation. |

Note: Do not apply more than once per growing season. Do not apply within 12 months of harvest.

| Crop   | Pest             | Rate         | Timing/Special Directions   |
|--|------------------|--------------|---|
| Kale   | Cutworms         | 14-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|  | Mole<br>Crickets | 7 lbs./A     | One to two days before planting, broadcast and immediately incorporate into soil. |
|  | Wireworms        | 21-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |
| Lettuce  | Cutworms         | 14-28 lbs./A | Broadcast and incorporate into soil just before planting.                         |
|  | Mole<br>Crickets | 7 lbs./A     | One to two days before planting, broadcast and immediately incorporate into soil. |
|  | Wireworms        | 21-28 lbs./A | Broadcast and incorporate into soil just before planting                          |
| Melons<br>(Cantaloupes,<br>Casabas,<br>Crenshaws,  | Cutworms         | 14-28 lbs./A | Broadcast and incorporate into soil just before planting                          |
| Honeydew,<br>Musk-<br>melons,<br>Persians,<br>and Hybrids<br>of these)<br>and Water-<br>melons | Wireworms        | 21-28 lbs./A | Broadcast and incorporate into soil just before planting                          |

| Crop                          | Pest             | Rate                   | Timing/Special Directions   |
|-------------------------------|------------------|------------------------|---|
| Mustard                       | Mole<br>Crickets | 7 lbs./A               | One to two days before planting, broadcast and immediately incorporate into soil.   |
| Onions<br>(Bulb and<br>Green) | Onion<br>Maggots | 14-28 lbs./A           | Broadcast just before planting and incorporate into the top 3-4 inches of soil.  Note: Diazinon will not control organophosphate resistant onion maggots. |
|                               | Wireworms        | 21-28 lbs./A           | Broadcast and incorporate into soil just before planting.   |
| Radishes                      | Mole<br>Crickets | 7 lbs./A               | One to two days before planting, broadcast and immediately incorporate into soil.   |
|                               | Root<br>Maggots  | 14-21 lbs./A           | Broadcast and incorporate into soil just before planting.   |
|                               | Wireworms        | 21-28 lbs./A           | Broadcast and incorporate into soil just before planting.   |
| Rutabagas                     | Mole<br>Crickets | 7 lbs./A               | One to two days before planting, broadcast and immediately incorporate into soil.   |
|                               | Wireworms        | 21-28 lbs./A           | Broadcast and incorporate into soil just before planting.   |
| inach                         | Cutworms         | 14-28 lbs.<br>per acre | Broadcast and incorporate into soil just before planting.   |
|                               | Wireworms        | 21-28 lbs.<br>per acre | Broadcast and incorporate into soil just before planting.   |

| Crop       | Pest                | Rate         | Timing/Special Directions   |
|------------|---------------------|--------------|---|
| Sweet Corn | Cutworms            | 14-28 lbs./A | Broadcast and incorporate into soil just before planting.                     |
|            | Seed Com<br>Maggots | 14-28 lbs./A | Broadcast just before planting and incorporate into the top 2 inches of soil. |
|            | Wireworms           | 21-28 lbs./A | Broadcast and incorporate into soil just before planting.                     |

#### FIELD CROPS

#### SPECIAL PRECAUTIONS TO REDUCE RISKS TO BIRDS

Birds feeding on treated areas may be killed. To reduce risks to birds:

- 1. Thoroughly incorporate this product into the soil during or immediately after application.
- 2. Collect or incorporate granules that are spilled during loading or are visible on soil surface in turn areas.

Biazinon 14G should be applied uniformly as directed for each crop.

| Crop        | Pest                          | Rate   | Timing/Special Directions   |
|-------------|-------------------------------|--|---|
| Sugar Beets | Sugar Beet<br>Root<br>Maggots | 7-14 lbs./A<br>or 4.7-<br>9.4 oz./<br>1,000 linear<br>ft. of row | Apply in a 5 to 7-inch band (based on 22" row spacing) incorporated 2-3 inches at planting. |
|             | Wireworms                     | 21-28 lbs./A   | Broadcast and incorporate into soil 4-8 inches just before planting.                        |

Note: Do not apply more than 28 lbs. of \_\_\_ Biazinon 14G per acre to the soil

per season.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Store in a cool, dry, well-ventilated area away from food or feed.

Pesticide: Pesticide or equipment rinsate that cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility.

Container: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles into application equipment. Then dispose of empty bag in a sanitary landfill or incinerate, or burn if allowed by state and local authorities. Stay out of smoke during burning. Do not reuse container.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800 130, day or night.

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Do not store near food or feed products.

| FIRST AID                 |   |  |  |
|---------------------------|---|--|--|
| This product coi          | ntains diazinon, an organophosphate insecticide that terase.  |  |  |
| if swallowed              | <ul> <li>Call a polson control center or doctor immediately for treatment advice.</li> <li>Have a person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> </ul> |  |  |
|                           | Do not give anything by mouth to an unconscious or convulsing person.   |  |  |
| If in eyes                | <ul> <li>Hold eye open and rinse slowly and gently<br/>with water for 15-20 minutes. Remove contact<br/>lenses, if present, after the first 5 minutes, then<br/>continue rinsing eye.</li> </ul>  |  |  |
|                           | ı before eating, drinking, chewing gum using  |  |  |
| If on skin<br>or clothing | <ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>ontrol Statements</li> </ul>  |  |  |
| Call a poison contr       | rol center or doctor for treatment advice. Have the product   |  |  |

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

#### **NOTE TO PHYSICIAN**

This product contains diazinon, an organophosphate that inhibits cholinesterase. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal, but should be administered only in conjunction with atropine.

#### **Personal Protective Equipment**

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### **Engineering Control Statements**

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### **User Safety Recommendations**

#### Users should:

 Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.