

PM 25

19713-97

1 of 21



Drexel

LINURON 4L
WEED KILLER

ACCEPTED
MAR 15 1985
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 19713-97

ACTIVE INGREDIENT:

Linuron [3-(3,4-dichlorophenyl)-1-methoxy-1-methylurea] . . . 41.0%

INERT INGREDIENTS 59.0%

Contains 4 Lbs. Linuron Per Gallon

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Statement of Practical Treatment

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF SPLASHED IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if irritation develops.

IF SPILLED ON SKIN: Immediately remove contaminated clothing, INCLUDING SHOES, and wash skin with soap and plenty of water. If irritation develops, send for a physician.

See Side Panel for Additional Precautionary Statements

EPA Reg # 19713-97
EPA Est # 19713-MS-1

Net Contents:

Gal. _____

Liters. _____



Drexel Chemical Company
P.O. BOX 8006, MEMPHIS, TENN 38108

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION! Harmful if Swallowed. May irritate eyes, nose, throat and skin. Avoid breathing spray mist. Avoid contact with skin, eyes, and clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

Required clothing and equipment for mixing/loading and applying linuron:

One-piece overalls which have long sleeves and long pants constructed of finely woven fabric as specified in the USDA/EPA Guide for Commercial Applicators.

Wide-brimmed hat.

Mid-forearm to elbow-length Chemical-resistant gloves.

Any article worn while applying linuron must be cleaned before reusing. Clothing which has been drenched or has otherwise absorbed concentrated pesticide must be disposed in a sanitary landfill, or incinerated, or if allowed by state and local authorities, by burning.

Instead of clothing and equipment specified above, the applicator can use an enclosed tractor cab which provides a filtered air supply (as described by Taschenberg and Bourke, 1975).

Non-aquatic

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Rotational Crop Restrictions

Do not rotate crops used for food or feed, which are not registered for use with linuron onto areas previously treated with this chemical.

Reentry Restriction

Do not reenter treated areas for 24 hours following application unless protective clothing is worn.

3 of 21

BEST AVAILABLE COPY

IMPORTANT

Injury to or loss of desirable trees or other plants may result from failure to observe the following:

Do not apply (except as recommended for crop use), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on home plantings or trees, shrubs or herbaceous plants, nor on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water. Keep from contact with fertilizers, insecticides, fungicides, and seeds.

Thoroughly clean all traces of LINURON 4L from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

GENERAL INFORMATION

DO NOT apply with aerial equipment.

DREXEL LINURON 4L Weed Killer is a liquid suspension to be mixed in water and applied as a spray for selective control of weeds in certain crops and for non-selective weed control on non-cropland areas. It is non-corrosive to equipment, non-flammable and non-volatile.

LINURON 4L may be applied to soil prior to emergence of weeds to control susceptible weed seedlings for an extended period of time; the degree of control and duration of effect will vary with the amount of chemical applied, soil texture, rainfall and other conditions. Soils high in clay or organic matter require higher dosages than soil low in clay or organic matter to obtain equivalent herbicide performance. Moisture is required to activate the chemical; best results occur if rainfall (or irrigation) occurs within 2 weeks of application. In the Columbia River Basin, use LINURON 4L only if crop is sprinkler irrigated.

LINURON 4L may also be used to control emerged weeds. Results vary with rate applied and environmental conditions; best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70° F or higher. Addition of a surfactant such as Drexel Surf Ac 820 to the spray (where recommended) increases contact effects of LINURON 4L.

LINURON 4L is recommended for use in conjunction with certain other herbicides for treatment of field corn, sorghum, soybeans and winter wheat; observe all cautions and limitations on labeling of all products used in mixtures.

Since the effect of LINURON 4L varies with soils, crop varieties, uniformity of application, and environmental conditions, it is suggested that growers limit their first use to small areas.

4 of 21

Drexel Linuron 4L
Weed Killer
Page 3

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

LINURON 4L should be used only in accordance with recommendations on this label. Do not use on any crop in Kern County, California.

Drexel will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Drexel. User assumes all risk associated with such non-recommended use.

SELECTIVE USE IN CROPS

PREEMERGENCE USE (Germinating Weeds): LINURON 4L, at recommended rates, controls weeds such as:

Broadleaves		Grasses
Florida beggarweed	Mustard	Barnyardgrass
Carpetweed	Nettleleaf goosefoot	(watergrass)
Chickweed	Pigweed	Canarygrass
Common dayflower	Purslane	Crabgrass
Florida purslane	Wild radish	Foxtail
(Florida pursley)	Common ragweed	(including giant)
Galinsoga	Smartweed	Goosegrass
Lambsquarters		Fall panicum

Partial Control

- Cocklebur
- Annual morningglory
- Prickly sida (teaweed)
- Sicklepod
- Velvetleaf (buttonweed)

Note: Consult "Amiben"¹, atrazine, "Dual"² 8E, "Lasso"³, "Ortho"⁴ Paraquat CL, propazine, "Prowl"⁵, "Ramrod"³, "Roundup"³, "Surflan"⁶ 75W, "Surflan"⁶ AS, "Tolban"², and "Treflan"⁶, labels for additional weeds controlled when LINURON 4L is used in conjunction with these herbicides.

LINURON 4L will not control established perennials such as bermudagrass, Canada thistle, field bindweed, johnsongrass, and purple nutsedge. The lower dosage rates are effective on coarser soils and the higher rates on finer soils and on the more resistant seedling weeds. Sufficient moisture (1/2" to 1" on moist soils; 1" to 2" on dry soils) in the form of rainfall or sprinkler irrigation is necessary after treatment to carry the chemical into the root zone of germinating weeds; best results are obtained when this occurs within two weeks after application. If heavy rainfall occurs soon after application, injury to crop may result.

5 of 21

A good seed bed must be prepared before application of LINURON 4L as crop injury may result if application is made to ground which is cloddy or compacted resulting in improperly planted seed. Plant seed to depth specified. Surface of the soil should not be cultivated or disturbed after application of LINURON 4L and before emergence of the crop as weed control may be reduced and crop injury may result. However, if moisture is insufficient to activate the herbicide, a shallow cultivation (rotary hoe preferred) should be made after emergence of row crops while weeds are small enough to be controlled by mechanical means.

POSTEMERGENCE USE (Emerged Seedling Weeds): LINURON 4L, at recommended rates, controls weeds such as:

<u>Broadleaves</u>		<u>Grasses</u>
Florida beggarweed	Annual morningglory	Barnyardgrass
Wild buckwheat	Nettleleaf goosefoot	(watergrass)
Carpetweed	Pigweed	Canarygrass
Cocklebur	Prickly sida	Crabgrass
Common dayflower	(teaweed)	Foxtail
Dog fennel	Purslane	(including giant)
Fiddleneck	Common ragweed	Goosegrass
(amsinckia)	Sesbania	Fall panicum
Florida purslane	Sicklepod	Texas panicum
(Florida pursley)	Smartweed	Annual ryegrass
Groundsel	Velvetleaf	Rattail fescue
Knawel	(buttonweed)	
Lambsquarters		
Mustard		

Note: Consult bromoxynil, "Butoxone"⁷, "Butyrac"¹ 000 and "Dyanmyte 3"⁸ labels for additional weeds controlled when LINURON 4L is used in conjunction with these herbicides.

Best results are obtained under conditions of high humidity and temperatures over 70° F. Control of emerged weeds under drought stress is usually impractical.

EQUIPMENT - SPRAY VOLUMES AND PRESSURES: Use a tractor-mounted fixed-boom sprayer properly calibrated to a constant speed and rate of delivery. Openings in screens should be equal to or larger than 50 mesh. Continuous agitation in the spray tank is required to keep the material in suspension. Agitate by mechanical or hydraulic means; if bypass or return line is used, it should terminate at bottom of tank to minimize foaming. Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping, or injury to the crop may result.

BEST AVAILABLE COPY

For preemergence application, use 25 to 40 gals. per acre and spray pressure of 30 to 40 psi. (for mixtures containing Paraquat CL or "Roundup", use 20 to 60 gals. per acre). For postemergence application, use sufficient volume (min. 25 gals. per acre) for thorough coverage of weed foliage; use spray pressure of 20 to 25 psi to keep spray drift to a minimum.

Aerial: LINURON 4L may be applied by aircraft to asparagus, carrots, potatoes, and wheat in 5 to 10 gals. of water per acre and to soybeans (preemergence only) in 2 to 10 gals. per acre. Avoid overlapping of spray swath and avoid application under conditions where excessive drift may occur. Where aerial application is made on bedded plantings, apply parallel to rows.

SPRAY PREPARATION: LINURON 4L and Tank Mixtures - Mix proper amount of LINURON 4L alone or with wettable powder herbicide, in spray tank with agitator running. For tank mixture with a liquid herbicide, dilute liquid product with 2 parts of water and add slowly as last ingredient to spray tank; for tank mix combinations with Paraquat CL or "Roundup", follow mixing instructions under Minimum or No-Tillage Soybeans. Where use of a surfactant is recommended, dilute with 10 parts of water and add as last ingredient to nearly full tank.

Fertilizer Spray Mixtures - For preemergence application, nonpressure nitrogen or fertilizer solution may be used in the spray mixture unless otherwise directed. Small quantities should be tested for compatibility by the following procedure before full scale mixing:

1. Put 1 pint of fertilizer solution in a quart jar.
2. Mix 2 teaspoons LINURON 4L with 2 tablespoons of water; mix thoroughly and add to fertilizer solution.
3. Close jar and shake well.
4. If other herbicides are to be used in the mixture, premix 2 teaspoons of wettable powders or 1 teaspoon of liquid with 2 tablespoons of water; add to LINURON 4L/fertilizer mixture.
5. Close jar and shake well.
6. Watch mixture for several seconds; check again in 30 minutes.
7. If mixture does not separate, foam, gel, or become lumpy, it may be used.
8. Mixing ability may be improved by adding compatibility agents such as Kalo Laboratories "Complex" or Witco Chemicals "Sponto" 168D. Follow directions on container.

7 of 21

If the mixture is compatible, prepare spray by adding the fertilizer solution to the spray tank first; then dilute LINURON 4L with 2 parts of water, thoroughly mix, and add slowly to spray tank with agitator running. For tank mixtures with other herbicides, follow directions above.

USE RATES: All rates of LINURON 4L (and tank mixtures) are expressed as broadcast rates; for band treatment, use proportionately less. For example, use 1/3 of the broadcast rate when treating a 14" band where row spacing is 42". Where a range of dosages is given, use the lower rate on coarser soils (low in clay or organic matter) and the higher rate on finer soils (high in clay or organic matter); for postemergence application, use the lower rate on smaller weeds and the higher rate on larger weeds.

SOIL LIMITATIONS: Unless otherwise directed, do not use on sand, loamy sand, gravelly soils or exposed subsoils nor on soils containing less than 1% organic matter as injury to the treatment or subsequent crops may result.

REPLANTING: If initial seeding fails to produce a stand, the same crop may be replanted in soil treated preemergence with LINURON 4L (or with recommended tank mixtures). Thoroughly rework soil before replanting; do not re-treat field with a second application as injury to the crop may result. West of Rocky Mountains - carrots or celery may be planted 4 months after last application; unless otherwise directed, do not plant any other crop until 1 year after the last application.

ASPARAGUS - California (Includes control of dudain melon and nightshade):

Direct Seeded or Newly Planted Crowns - Do not exceed 4 pints total per acre per season; do not use surfactant or fertilizer solution in spray mixture.

Preemergence Application: Make a single application of 2 to 4 pints per acre after planting seed 1½" deep in coarse soil and 1" deep in fine soils. During planting operation, spray activate charcoal ("Aqua Nu-Char"⁹ or "Gro-Safe"¹⁰) as a 1" band on soil surface directly over seed rows at the rate of 300 lbs. per acre (equivalent to 15 lbs. per acre of crop with 20" row spacing).

Postemergence Application: Make 1 or 2 applications of 1 to 2 pints per acre when ferns are in 6" to 18" stage and weeds are not over 4" tall.

BEST AVAILABLE COPY

Established Beds - Do not apply within 1 day of harvest; do not exceed 8 pints total per acre per season; do not use surfactant or fertilizer solution in spray mixture.

Preemergence Application: Make a single application of 2 to 4 pints per acre.

Postemergence Application: Make 1 to 4 applications of 1 to 2 pints per acre before weeds exceed 4" in height. Apply before cutting season or immediately after cutting.

Directed Postemergence Application (Fern Stage): Make a single application of 4 to 8 pints per acre as a directed spray to base of plants for control of dudain melon.

Note: If more than 4 pints LINURON 4L per acre is applied per season, do not plant any other crop until 1 year after last application.

CARROTS

Preemergence Application - Michigan and Wisconsin - Make a single application of 1 to 3 pints per acre after planting but before carrots emerge; plant seed at least $\frac{1}{2}$ " deep. Use the lower rate on lighter soils (low in clay or organic matter) and the higher rate on heavier soils (high in clay or organic matter). Subsequent postemergence application may be made provided the total does not exceed 4 pints LINURON 4L per acre per season.

Postemergence Application - U.S: Apply $1\frac{1}{2}$ to 3 pints per acre as a non-directed spray after carrots are at least 3" tall. Apply before annual grasses exceed 2" in height and before broadleaf weeds exceed 6" in height. Repeat application may be made but do not exceed 4 pints LINURON 4L per acre (West of Rocky Mountains, do not exceed 3 pints LINURON 4L per acre). Do not exceed 40 psi spray nozzle as crop injury may result.

LINURON 4L may be applied following an application of Stoddard solvent provided treatments are at least one day apart. Stoddard solvent may be applied following an application of LINURON 4L provided treatments are at least 2 weeks apart. Shorter time intervals between applications may result in injury to the crop. Do not apply LINURON 4L as a tank mixture with Stoddard solvent, surfactants, nitrogen or fertilizer solution, other pesticides, nor when temperature exceeds 85° F. as crop injury may result.

Note: Because carrot varieties vary in their resistance, determine tolerance to LINURON 4L prior to adoption as a field practice to prevent possible crop injury; do not treat susceptible varieties which show an initial burning of foliage following postemergence treatment with LINURON 4L.

BEST AVAILABLE COPY

9 of 21

Drexel Linuron 4L
Weed Killer
Page 8
CELERY

East of Rocky Mountains - Post-transplant Application: Make a single application of 1½ to 3 pints per acre. Apply as a non-directed spray after celery is transplanted and established, but before celery is 8" tall. Apply before annual grasses exceed 2" in height and before broadleaf weeds exceed 6" in height. In the Northeast, use only on celery grown on muck soils.

Do not exceed 40 psi spray nozzle pressure, and do not apply when temperature exceeds 85° F. nor as a tank mixture with surfactants, nitrogen or fertilizer solution, or other pesticides as injury to the crop may result. Do not replant to crops other than celery or carrots within 4 months after application as injury to subsequent crops may result.

CORN (FIELD)

East of Rocky Mountains - Preemergence Application: Select one of the following herbicide treatments for application as a tank mixture. Make a single application after planting but before crop emerges. Plant seed at least 1 3/4 deep on flat or raised seedbeds only or injury to the crop may result. Do not spray over top of emerged corn.

LINURON 4L + "Lasso" (Tank Mixture)

Soil Texture	Product Per Acre	
	1 to 3% Organic Matter LINURON 4L (Pts.) + "Lasso" (Qts.)	3 to 6% Organic Matter LINURON 4L (Pts.) + "Lasso" (Qts.)
Sandy loam	2/3 to 1½ + 3/4 to 1	1½ to 2 + 1 to 1½
Silt loam	1 to 1-2/3 + 1 to 1½	1-2/3 to 2½ + 1½ to 2
Clay loam	1½ to 2 + 1½ to 2	2 to 3 + 2 to 2½

Replanting: Corn or soybeans may be replanted within 4 months; after 4 months, any crop may be replanted.

LINURON 4L + "Ramrod" (Tank Mixture)

Soil Texture	Product Per Acre	
	1 to 3% Organic Matter LINURON 4L (Pts.) + "Ramrod" 65 (Lts.)	3 to 6% Organic Matter LINURON 4L (Pts.) + "Ramrod" 65 (Lbs.)
Sandy loam	2/3 to 1-1/3 + 1 to 2	1-1/3 to 2 + 2 to 3
Silt loam	1 to 1-2/3 + 1½ to 2½	1-2/3 to 2½ + 2½ to 4
Clay loam	1½ to 2 + 2 to 3	2 to 3 + 3 to 4-3/4

Replanting: Corn may be replanted within 4 months; after 4 months, any crop may be planted.

AGRI-TECHNICAL 1971

10 of 21

LINURON 4L + atrazine (Tank Mixture)

Soil Texture	Product Per Acre	
	1 to 2% Organic Matter "LINURON 4L (Pts.) + atrazine 80% (Lbs.)	2 to 5% Organic Matter "LINURON 4L (Pts.) + atrazine 80% (Lbs.)
Sandy loam	2/3 to 1 + 1/2 to 2/3	1 to 2 + 2/3 to 1 1/2
Silt loam	1 to 1 1/2 + 2/3 to 1	1 1/2 to 2 1/2 + 1 to 1 1/2
Clay loam	1-1/3 to 1-2/3 + 3/4 to 1	1-2/3 to 3 + 1 to 2

Replanting: Corn may be replanted within 6 months. After 6 months, any crop may be planted except do not follow treated corn with sugar beets, tobacco, or vegetables in rotation.

CORN (FIELD AND SWEET)

Directed Postemergence Application: Make a single application as a directed spray after corn is at least 15" high (measured to the highest leaf surface on free standing plants). Do not spray over top of corn. Apply only when there is sufficient differential between height of corn and weeds so that the directed spray thoroughly covers all weed foliage without contact of upper leaves or whorl of corn by spray or drift, as such contact may cause crop injury. Early cultivation (rotary hoe or other suitable equipment) will aid in achieving proper differential between height of corn and weeds.

Use 1 1/2 to 3 pints per acre; add 1 pint Surf-Ac 820 for each 25 gals. spray mixture. For field corn, non-pressure nitrogen solution may be substituted for all or part of the water. Use the lower rate on lighter soils (low in clay or organic matter) and when weeds do not exceed 2" in height; use the higher rate on heavier soils (high in clay or organic matter) for weeds up to 5" in height.

COTTON

East of Rocky Mountains - Directed Postemergence Application: Apply as a directed spray; adjust nozzles to minimize contact to cotton leaves with spray or drift or crop injury may result. Do not spray over top of cotton.

Apply 1 to 1 1/2 pints per acre when cotton is at least 8" tall and emerged weeds do not exceed 2" in height; add 1 pint Surf Ac 820 for each 25 gals. spray mixture. If needed, a second application at same rate may be made one week or later after initial treatment. Alternatively, after cotton is 20" tall, make a single application of 2 to 3 pints per acre following last cultivation; if emerged weeds are present, add Surf Ac 820 as directed above. Note: Do not use on Pima varieties of cotton. Do not graze treated fields or feed forage from treated areas to livestock. Do not feed gin trash to livestock.

Preemergence Application: Make a single application of 1½ to 3 pints per acre. Apply after planting but before crop emerges. Plant seed at least ½" deep.

POTATOES

Preemergence Application: Make a single application as a broadcast spray after planting but before crop emerges. Plant seed at least 2" deep. Do not spray over top of emerged potatoes. If beds are to be "dragged" and/or "hilled", apply after the final "dragging or "hilling" operation. Apply before grasses are 2" tall and before broadleaf weeds are 6" tall, preferably just before or when weed seedlings emerge. If emerged weeds are present, add 1 pint Surf Ac 820 for each 25 gals. spray mixture. In irrigated areas, best results are obtained when application is made to moist soil, followed within 2 weeks by 1" to 2" of sprinkler irrigation (or rainfall). On powder dry soils, irrigate prior to herbicide application and follow with sprinkler irrigation to activate the herbicide.

East of Rocky Mountains: Apply 1½ to 2½ pints per acre on the coarser soils (sandy loams, silt loams; 1 to 2% organic matter) and 2½ to 4 pints per acre on finer soils (silts, clay loams; 2 to 5% organic matter); on soils over 5% organic matter, use 4 pints per acre and apply to emerged weeds (before potatoes emerge).

Wisconsin - Central Sands Area: Apply 1 pint per acre on sands and 2 pints per acre on loamy sands.

SORGHUM

Preemergence Application: Select one of the following herbicide treatments for application as a tank mixture. Make a single application after planting but before crop emerges. Plant seed at least 1" deep on flat or raised seedbeds only or injury to the crop may result. Do not spray over top of emerged sorghum.

LINURON 4L + propazine - Southwest (Tank Mixture)

	Product Per Acre	
	1 to 2% Organic Matter LINURON 4L (Pts.) + propazine 80% (Lbs.)	2 to 4% Organic Matter LINURON 4L (Pts.) + propazine 80% (Lbs.)
Soil Texture		
Sandy loam	5/8 to 1 + 3/8 to 5/8	1 to 1½ + 5/8 to 1
Silt loam, clay loam	1 to 1½ + 5/8 to 1	1 to 2 + 5/8 to 1

BEST AVAILABLE COPY

Replanting: Do not follow treated sorghum with any fall crop, nor with sugar beets, tobacco, vegetables or potatoes in rotation. Prior to replanting, thorough seedbed preparation including fall or spring plowing is recommended.

LINURON 4L + "Ramrod" - Great Plains (Tank Mixture)

Soil Texture	Product Per Acre	
	1 to 3% Organic Matter LINURON 4L (Pts.) + "Ramrod" 65 (Lbs.)	3 to 6% Organic Matter LINURON 4L (Pts.) + "Ramrod" 65 (Lbs.)
Sandy loam	2/3 to 1-1/3 + 1 to 2	1-1/3 to 2 + 2 to 3
Silt loam	1 to 1-2/3 + 1 1/2 to 2 1/2	1-2/3 to 2 1/2 + 2 1/2 to 4
Clay loam	1 1/2 to 2 + 2 to 3	2 to 3 + 3 to 4-3/4

Note: Do not graze or feed sorghum forage or silage from treated field to dairy animals. Replanting: Sorghum or field corn may be replanted within 4 months; after 4 months, any crop may be planted.

Directed Postemergence Application: Make a single application of LINURON 4L as a directed spray; add 1 pint Surf Ac 820 for each 25 gals. spray mixture. If sprayer is equipped with skids, shoes or shield, apply 1 pint per acre when sorghum is 12" tall (free standing plants) and weeds are up to 2" in height; use 1 to 2 pints per acre when sorghum is 15" tall and weeds are 2" to 4" in height. If boom drops are used, apply 1 to 2 pints per acre when sorghum is at least 18" tall and weeds are 2" to 4" in height. Apply only when there is sufficient differential between height of sorghum and weeds so that the directed spray thoroughly covers all weed foliage without contact of upper leaves or whorl or sorghum by spray or drift as such contact may cause crop injury. Note: Do not graze or feed plants to livestock within 3 months after postemergence application.

SOYBEANS

Preemergence Application: Select one of the following herbicide treatments and make a single application after planting but before crop emergence. LINURON 4L, alone or as a tank mixture with "Lasso", may be applied by aircraft (2 to 10 gals. water per acre). Plant seed at least 1 3/4" deep on flat or raised seedbeds only or injury to the crop may result. Injury to soybeans may result if application is made to fields with standing water or fields too wet to cultivate. Do not spray over top of emerged soybeans.

LINURON 4L Alone - if weeds have emerged, add 1 pint Surf Ac 820 for each 25 gals. spray mixture.

13 of 21

Soil Texture	Pints LINURON 4L Per Acre	
	1/2 to 2% Organic Matter	2 to 5% Organic Matter
Sandy loam	1 to 1-2/3	1-2/3 to 3
Silt loam	1 1/4 to 2-1/3	2-1/3 to 4
Clay loam	1-1/3 to 2-2/3	2-2/3 to 5 (Over 5% organic matter, use 6 pints)

Replanting: Soybeans or field corn may be replanted within 4 months; after 4 months, any crop may be planted.

LINURON 4L Following "Treflan" or "Tolban" Preplant - Where "Treflan" or "Tolban" has been used as a preplant incorporated treatment (according to directions on product label), apply LINURON 4L as a separate operation using rates recommended above for LINURON 4L alone. For rotation crops, follow instructions on "Treflan" or "Tolban" label.

LINURON 4L + "Amiben" (Tank Mixture)

Soil Texture	Product Per Acre	
	1/2 to 3% Organic Matter LINURON 4L (Pts.) + "Amiben" (Qts.)	3 to 6% Organic Matter LINURON 4L (Pts.) + "Amiben" (Qts.)
Sandy loam	2/3 to 1 1/4 + 3	1 1/4 to 2 + 3
Silt loam	1 to 1-2/3 + 3	1-2/3 to 2 1/4 + 4
Clay loam	1 1/4 to 2 + 4	2 to 3 + 5

Replanting: Soybeans or field corn may be replanted within 4 months; after 4 months, any crop may be planted.

LINURON 4L + "Dual" 8E (Tank Mixture)

Soil Texture	Product Per Acre	
	1/2 to 3% Organic Matter LINURON 4L (Pts.) + "Dual" 8E (Pts.)	3 to 6% Organic Matter LINURON 4L (Pts.) + "Dual" 8E (Pts.)
Sandy loam	2/3 to 1 1/4 + 1 1/4	1 1/4 to 2 + 1 1/4
Silt loam	1 to 1-2/3 + 1 1/4	1-2/3 to 2 1/4 + 2
Clay loam	1 1/4 to 2 + 2	2 to 3 + 2 to 2 1/4

Note: Do not graze or feed forage from treated areas to livestock.
Replanting: Soybeans or field corn may be replanted within 4 months; for rotation crops, follow instructions on "Dual" 8E label.

BEST AVAILABLE COPY

14821

LINURON 4L + "Lasso" (Tank Mixture)

Soil Texture	Product Per Acre	
	1/2 to 3% Organic Matter "LINURON 4L (Pts.) + "Lasso" (Qts.)	3 to 6% Organic Matter "LINURON 4L (Pts.) + "Lasso" (Qts.)
Sandy loam	2/3 to 1 1/2 + 3/4 to 1	1 1/2 to 2 + 1 to 1 1/2
Silt loam	1 to 1-2/3 + 1 to 1 1/2	1-2/3 to 2 1/2 + 1 1/2 to 2
Clay loam	1 1/2 to 2 + 1 1/2 to 2	2 to 3 + 2 to 2 1/2

Replanting: Soybeans or field corn may be replanted within 4 months; after 4 months, any crop may be planted.

LINURON 4L + "Prowl" (Tank Mixture) or LINURON 4L Following "Prowl"
Preplant - Apply tank mixture as preemergence treatment only; for pre-plant treatment, incorporate "Prowl" and then follow with LINURON 4L as a separate operation.

Soil Texture	Product Per Acre	
	1 to 3% Organic Matter LINURON 4L (Pts.) + "Prowl" (Pts.)	3 to 6% Organic Matter LINURON 4L (Pts.) + "Prowl" (Pts.)
Sandy loam	2/3 to 1 1/2 + 1 to 1 1/2	1 1/2 to 2 + 1 1/2
Silt loam	1 to 1-2/3 + 1 1/2 to 2	1-2/3 to 2 1/2 + 1 1/2 to 2
Clay loam	1 1/2 to 2 + 1 1/2 to 2	2 to 3 + 2 to 2 1/2

Replanting: Soybeans or field corn may be replanted within 4 months; other crops may be planted the following year.

LINURON 4L + "Ramrod" - Seed Crop Only (Tank Mixture)

Soil Texture	Product Per Acre	
	1 to 3% Organic Matter LINURON 4L (Pts.) + "Ramrod" 65 (Lbs.)	3 to 6% Organic Matter LINURON 4L (Pts.) + "Ramrod" 65 (Lbs.)
Sandy loam	2/3 to 1 1/2 + 1 to 2	1 1/2 to 2 + 2 to 3
Silt loam	1 to 1-2/3 + 1 1/2 to 2 1/2	1-2/3 to 2 1/2 + 2 1/2 to 4
Clay loam	1 1/2 to 2 + 2 to 3	2 to 3 + 3 to 4-3/4

Note: Do not graze or feed forage from treated areas to livestock; do not use seed for food, feed, or oil purposes.

Replanting: Soybeans, sorghum, or field corn may be replanted within 4 months; after 4 months, any crop may be planted.

LINURON 4L + "Surflan" 75W or "Surflan" AS (Tank Mixture)

Soil Texture 1/2 to 3% Organic Matter	Product Per Acre		
	LINURON 4L (Pints)	+	"Surflan" 75W (Lbs.) or "Surflan" AS (Quarts)
Sandy loam	2/3 to 1 1/4	+	2/3 to 1 or 1/2 to 3/4
Silt loam	1 to 1-2/3	+	1 to 1-1/2 or 3/4 to 1
Clay loam	1 1/4 to 2	+	1-1/3 to 1-2/3 or 1 to 1 1/4

Note: Do not use treated vines for feed or forage. Replant: Soybeans may be replanted within 4 months; after 4 months, any crop may be planted except do not plant root crops within 12 months.


SOYBEANS - Minimum or No-Tillage: LINURON 4L may be used with "Ortho" Paraquat CL or "Roundup" in tank mix combinations for preemergence and postemergence control of many broadleaf weeds and grasses and for top kill or suppression of some perennials where soybeans will be planted directly into a preformed bed (stale seed bed), cover crop, or in previous crop residues such as corn or small grain stubble. Addition of "Surflan" 75W, "Surflan" AS, "Dual" 8E or "Lasso" to the tank mixture improves control of grasses and volunteer small grains.

Apply with ground equipment immediately before, during, or after planting but before crop emerges. Use 20 to 60 gals. of water per acre. Use the higher gallonage for dense stubble or vegetation. Maintain constant agitation.

LINURON 4L + Paraquat CL - Select the rate from the table for pre-emergence use of LINURON 4L alone and thoroughly mix in spray tank first; then add Paraquat CL as directed under "Paraquat Combinations" below.

LINURON + "Surflan" 75W (or AS) + Paraquat CL - Select the rate from the table from pre-emergence use of LINURON 4L + "Surflan" 75W or "Surflan" AS tank mixture and thoroughly mix in spray tank first; then add Paraquat CL as directed under Paraquat Combinations below. Do not use treated vines for feed or forage.

LINURON 4L + "Dual" 8E + Paraquat CL or "Roundup" - Thoroughly mix LINURON 4L and "Dual" 8E in spray tank first according to directions; then add Paraquat CL or "Roundup" as directed under Paraquat Combination below. Do not graze or feed forage from treated areas to livestock.



16 of 21

Soil Texture	Product Per Acre	
	$\frac{1}{2}$ to 3% Organic Matter LINURON 4L (Pts.) + "Dual" 8E (Pts.)	3 to 6% Organic Matter LINURON 8L (Pts.) + "Dual" 8E (Pts.)
Sandy loam	$\frac{2}{3}$ to $1\frac{1}{2}$ + $1\frac{1}{2}$	$1\frac{1}{2}$ to 2 + $1\frac{1}{2}$
Silt loam	1 to $1\frac{2}{3}$ + 2	$1\frac{2}{3}$ to $2\frac{1}{2}$ + 2
Clay loam	$1\frac{1}{2}$ to 2 + 2 to $2\frac{1}{2}$	2 to 3 + 2 to $2\frac{1}{2}$

LINURON 4L + "Lasso" + Paraquat CL or "Roundup" - Thoroughly mix LINURON 4L and "Lasso" in spray tank first according to directions; then add Paraquat CL or "Roundup" as directed under Paraquat Combinations or "Roundup" Combinations below.

Soil Texture	Product Per Acre	
	$\frac{1}{2}$ to 3% Organic Matter LINURON 4L (Pts.) + "Lasso" (Qts.)	3 to 6% Organic Matter LINURON 4L (Pts.) + "Lasso" (Qts.)
Sandy loam	$\frac{2}{3}$ to $1\frac{1}{2}$ + 2	$1\frac{1}{2}$ to 2 + $2\frac{1}{2}$
Silt loam	1 to $1\frac{2}{3}$ + $2\frac{1}{2}$	$1\frac{2}{3}$ to $2\frac{1}{2}$ + 3
Clay loam	$1\frac{1}{2}$ to 2 + $2\frac{1}{2}$	2 to 3 + 3

Paraquat Combinations - Select and tank mix one of the above treatments; then add 1 to 2 pints Paraquat CL per acre. Use the higher rate for weeds 4" to 6" tall. As the last ingredient, add $\frac{1}{2}$ pint Surf Ac 820 per 100 gals. of spray mixture. Maintain constant agitation.

"Roundup" Combinations - Select and tank mix one of the above treatments. As last ingredient, add $1\frac{1}{2}$ quarts of "Roundup" per acre for control of emerged annualweeds or 2 to 4 quarts per acre for control of emerged perennial and annual weeds.

SOYBEANS - Directed Postemergence Application: Apply LINURON 4L alone or as a tank mixture with, 2,4-DB or "Dynamyte 3, as a directed spray to cover weed foliage with minimum contact of the soybean plant. Do not spray higher than 3" on the soybean stem or crop injury may result. Do not spray over top of soybean plants. For broadcast application use a single flood-type spray nozzle. ("K" series or equivalent) per middle mounted on an oiling shoe or gauge wheel. For band treatment, use two nozzles per row mounted on oiling shoes or gauge wheels, one on each side of row. To avoid spray drift, which may cause crop injury, do not exceed nozzle pressure of 25 psi nor use nozzle tips smaller than 8002 T-Jet (or equivalent) and do not spray under windy conditions. For best results, use a preemergence treatment (such as LINURON 4L) or cultivation to control early weed growth and to increase the differential between height of soybeans and weeds. NOTE: Do not use on soils with less than $\frac{1}{2}$ % organic matter. Do not apply within 60 days of harvest.

17 of 21

Do not feed soybean forage or hay to livestock from fields treated postemergence. Harvested soybeans may be used for food, feed, or oil purposes.

LINURON 4L Alone: For soybeans at least 8" high and when weeds do not exceed 2" in height, apply $\frac{1}{2}$ to 1 pint per acre. Make a second application at same rate if new flush of weeds occurs.

For soybeans at least 12" high and when weeds do not exceed 4" in height, make a single application of 1 to 2 pints per acre. Alternatively, make a split application of 1 pint per acre followed by a second application at the same rate after 1 week or later.

Add 1 pint Surf Ac 820 for each 25 gals. spray mixture. Do not apply more than 2 pints LINURON 4L per acre per season for postemergence treatment.

LINURON 4L + 2,4-DB (Tank Mixture): For soybeans at least 8" high and when weeds do not exceed 4" in height, apply 1 pint LINURON 4L plus $\frac{1}{5}$ lb. 2,4-DB (13 fl. oz. "Butyrac" 200 or 1 pt. "Butoxone") per acre. A second application may be made if needed, but do not make more than 2 applications per season.

LINURON 4L + "Dynamyte" 3 (Tank Mixture): For soybeans at least 8" high and when weeds do not exceed 3" in height apply 1 pint LINURON 4L plus 2 quarts of "Dynamyte" 3 per acre. Add 1 pint Surf Ac 820 for each 25 gals. spray mixture. A Second application may be made if needed, but do not make more than 2 applications per season. Do not apply after soybeans begin to bloom.

WHEAT (WINTER)

(Drill Planted) - Idaho, Oregon, Washington: Plant seed at least 1" deep; when seed is planted during abnormally dry weather, treat after soil has been settled by rainfall or irrigation. Apply as a broadcast spray prior to emergence of wheat or to semi-dormant wheat plants. Application to actively growing plants may result in temporary yellowing (chlorosis) of wheat. Winter sensitive varieties such as McDermid and Hyslop may be less tolerant of "LINURON 4L than winter hardy varieties such as Gaines and Nugaines.

Crop injury may result where severe winter stress, disease or insect damage follows application, and also from failure to observe the following: Do not use on sand or loamy sand soils, nor on gravelly or sandy loams low in organic matter (less than 1%), nor on thinly covered or exposed sub-soil areas (clay knobs); do not treat wheat planted less than 1" deep; do not treat wheat where winter climatic conditions have caused "heaving" of plants; do not treat wheat plants lacking in vigor due to poor emergence, insect damage, disease, high alkalinity or other causes; do not apply after wheat has reached the "boot" stage of maturity nor when maximum daily temperature exceeds 60° F;



do not use LINURON 4L with surfactants nor nitrogen fertilizer or solution after wheat has emerged.

Do not graze or feed immature plants to livestock. Do not replant treated areas to any rotation crop within 6 months after last application as injury to subsequent crops may result.

West of Cascade Range: Make a single application of 2 to 3½ pints per acre as soon as possible after planting. If wheat and weeds have emerged, apply before weeds are 3" to 4" tall.

East of Cascade Range: Make a single application of LINURON 4L alone or, where recommended below, as a tank mixture with bromoxynil. If fall-planted wheat fails to grow due to winter kill or adverse growing conditions after fall treatment, allow 4 months before planting spring wheat. Do not re-treat field with a second application during the same crop year as injury to the crop may result.

Where Average Annual Rainfall Exceeds 16 Inches:

Fall Treatment: For early fall-planted wheat (seeded before September 10) apply 1 to 1½ pints LINURON 4L per acre either before or after wheat has emerged but before weeds are 2" tall. Treatment after October 1 generally gives best results. Do not apply after soil freezes in the fall.

Spring Treatment: Apply 1 to 1½ pints LINURON 4L per acre as soon as wheat starts to grow in the spring. Application after weeds have reached a height of 2" may give poor results.

Where Average Annual Rainfall is 10 to 16 Inches:

Fall or Winter Treatment: After wheat is planted in the fall, apply 1 to 1½ pints LINURON 4L per acre when sufficient moisture is available to germinate wheat seed. Apply either before or after wheat has emerged, but before weeds are 2" tall and before the soil freezes. Application later than March 1 may give poor results.

Where Average Annual Rainfall is 10 to 20 Inches:

Fall or Spring treatment: Apply ½ pint LINURON 4L plus ¼ lb. bromoxynil per acre as a tank mixture, either in the fall after wheat has emerged but before soil freezes or in the spring as soon as soil thaws; apply before weeds are 2" tall or across.

19-8-21

NON-CROP WEED CONTROL

For short-term control of annual weeds on non-cropland areas such as roadsides and fence rows - apply 2 to 6 pints LINURON 4L per acre in 40 to 100 gals. of water. For best results, apply shortly before weed growth begins or at early seedlings stage of growth. For control of established annual weeds, add Surf Ac 820 at rate of 2 qts. per 100 gals. of spray mixture and apply as a thorough coverage spray during periods when daily temperatures exceed 70° F, and before weed growth exceeds 8" in height.

- 1Registered trademark of Union Carbide Agricultural Products Company, Inc.
- 2Registered trademark of Ciba-Geigy Corporation
- 3Registered trademark of Monsanto Company
- 4Registered trademark of Chevron Chemical Company
- 5Registered trademark of American Cyanamid Company
- 6Registered trademark of Elanco Products Company
- 7Registered trademark of Rhone-Poulenc Inc.
- 8Registered trademark of Drexel Chemical Company
- 9Registered trademark of Westvaco Corporation
- 10Registered trademark of ICI United States, Inc.

20721

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal.

STORAGE INSTRUCTIONS

Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. Avoid storage at high temperatures. Do not stack over 2 pallets high. Move containers by handles or cases. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Avoid contamination with acids or alkalis. Store in original containers only. If the contents are leaking or material is spilled follow these steps:

1. Contain spill, absorb with a material such as saw dust, clay granules or dirt.
2. Collect and place in suitable containers for disposal.
3. Wash area with water and soap to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Place a leaking container in a plastic tub and transfer contents as soon as possible to an empty original container.
6. Do not allow run off to enter sewer or contaminate water supplies.
7. Dispose of waste as indicated below:

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

210821

WARRANTY - CONDITION OF SALE:

DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of Drexel Chemical Company or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result.

ALL SUCH RISKS ARE ASSUMED BY THE BUYER.

Drexel warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. Drexel makes no other warranties, express or implied, including FITNESS OR MERCHANTABILITY. In no case shall Drexel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by Drexel Chemical Company and is accepted as such by the Buyer.