

ATTACHEMENT #1

PM 10 19713-123

PM 72

BEST AVAILABLE COPY

NOT
LAW
FEDERAL NOTICE
DATE MAY 4 1992

DAMOIL

DORMANT & SUMMER SPRAY OIL

ACTIVE INGREDIENT:

Petroleum Oil 98.0%

INERT INGREDIENTS: 2.0%

TOTAL: 100.0%

Petroleum Oil Min. Unsulfonated Residue—92%

Petroleum Oil Classification—Unclassified

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel For Additional Precautionary Statements

EPA Reg. No. 19713-123

EPA Est. No. 19713-TN-1

11234-0001

Manufactured By:

Drexel Chemical Company

P.O. BOX 9208, MEMPHIS, TN 38108-0208

Net Contents ■ 55 Gallons

■ 30 Gallons

■ 5 gallons

ACCEPTED

FEB 14 1992

Under the Federal Insecticide, Fungicide, and
Rodenticide Act, as amended, for the
purpose of registration
EPA Reg. No. 19713-123

69

19713-123-10R-1

WARNING TO HUMANS AND DOMESTIC ANIMALS
CAUTION: HARMFUL IF SWALLOWED. Avoid breathing of dust. Do not take internally. Avoid Contact with Skin and Eyes. Remove clothing off instantly. Wear proper hot-weather work clothing. Change to clean clothing daily. Wash thoroughly after working.
Do not apply this product in such manner as to directly or through dirt come in contact with other persons. The area being treated must be vacated by untreated persons.

IF SWALLOWED: Do not induce vomiting because of aspiration hazard. Call a physician or Poison Control Center.

IF IN EYES: Immediately flush eyes with plenty of water. Get medical attention if irritation persists.

IF-ON SKIN immediately remove contaminated clothing, **SHOULDER SHOULDER**, and wash skin with soap and plenty of water. Get medical attention if irritation persists.

See Side Panel For Additional Preliminary Statements.

Do not spray directly on water. Do not spray when weather conditions favor drift from driven sprayer. This product is toxic to fish. Do not overgraze water by closing of equipment or disuse of water.

Do not use, pour, spill or store near heat or open flame.

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must inform workers of areas or fields that may not be entered without specific protective clothing and sprays have dried, and appropriate actions to take in case of accidental exposure, as described under Precautionary Statements on this label. When oral warnings are given, warnings shall be given in a language customarily understood by the workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. CAUTION: Area treated with Dimecron (state of application). Do not enter without appropriate protective clothing and sprays have dried. In case of accidental exposure, flush eyes or skin with plenty of water. Call physician if irritation persists. Remove and wash contaminated clothing before reuse.

DANOR is an especially refined Phenyonic Oil Contact Insecticide. It is for use on plants in foliage of the domestic stage. When using this product in combination with other pesticides as directed on this label, observe appropriate cautions, instructions for the products involved.

Do not combine Sulfur sprays with DREXEL DAMOL on foliage and do not follow a previous application of Sulfur or apply Sulfur following DREXEL DAMOL until sufficient time has elapsed for safety (at least 30 days). If in doubt, spray a portion of one tree previously treated with Sulfur, using 1 quart to 25 gallons water. If sufficient time has not elapsed, buds will show in several days. Do not use DREXEL DAMOL when temperatures are above 90° F. When temperatures drop below 32° F (usually night is a low), do not spray tree and vine crops until daytime temperatures have warmed to a point above 32° F. and are free from moisture due to melting frost or heavy dew. Do not use on foliage closer than 60 days after or 90 days before using Captan or Politen products.

Notice: This restriction does not apply to use during the dormant period. Do not use in combination with certain oil crops other than cotton. Application to trees weakened by disease, drought, drying winds, or high nitrogen applications may result in oil injury.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF SPRIGATION SYSTEM.
Amounts Given Are For 100 Gallons Of Diluent Spray.

ALFALFA (Seed Crop Only): Use 1 quart per acre in 20 to 30 gallons of water by ground application or 10 to 15 gallons by air when used in combination and at the recommended label rates with either a Malathion 5 Emulsion for control of the pests listed on the product label, or Spiner-Mite. Use 1 quart per acre at the dilutions listed above with Spiner-Mite at the recommended label rates.

ALMONDS, APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES (Dormant or Delayed Dormant): *Brown Apricot Scab*, *Black Scab*, *San Jose Scab*, *Olive Spot* (*Parasitic Scab*), *Cherry Scab*, *Fruited Scab*, *Golden Pear Scab*, *European Fruit Locusts*, *Purple Scab*, *Parasitic*, *Soft Brown Scab*, *Corky* (*Other Scab*), *Stem of Ashes*, *Red Spider Mites*—1½ to 2 gals. **Summer Use**—*Red Spider Mites*—3 to 4 qts.; **Summer Stages of Brown Apricot Scab**, *San Jose Scab*—1 gal.

APPL. 23: Post Mts—Delayed Dormant—1½ to 2 gals. Summer Use—1 to 1½ gals.

APPLER, PEARS: Dormant or Polysed Dormant—San Jose Scale, Indian Pear Scale, Eggs of Aphids, Red Spider Mites—1½ to 2 gal. Summer Use: Red Spider Mites—3 to 4 gal. Concentrated Sprays—Pears (Calif. only)—4 to 6 gal. in 90 to 120 gal. of water per acre. Summer Forms of San Jose and Indian Pear Scale—1 to 1½ gal.

Every education—like 2 yrs. COLLEGE, plus 10 yrs. of 3 Posters & Periodicals.

~~2245726—Mudbug—Seawater—Use 4 to 6 gals. per 2 to 2½ cu. of~~
~~freshwater 4 bushels is 200 to 250 gals. of water per acre. Do not exceed 2½ cu.~~
~~of freshwater 4 bushels per acre.~~

LIMACONS, CRAP-SUCKERS: And Snails. Yellow Snails, Purple Snails, Black Snails, And Snails Alone—1 to 1 1/2 gal. CABBAGE, 26oz 1 lb. of 1/2 dozen 25 vichins. For best results, spray during winter and spring months when the snails are more easily captured and 2000 generally best relative to all sprays. Be not too within 6 weeks of interest on 1/2 acre.

CALVER'S Petroleum Soots, Sulfur Lining, Chloroform Soots, Soot Soots—Use 10 gals. DABCO, or combine 1 gal. DABCO with 1½ pints of a Flammable 4 pound Emulsive per 100 gals. water. Apply in the air-blown or foamed-blown as in footnote.

MAVER. ORANGE: Red Scale, Black Scale, Yellow Scale, Purple Scale, Red Spider Mite.—3 gals. DANCER, plus 1 lb. of a Malicious 25 WORMS. Do not use on Floral Orange after Sept. 1.

YALENCIA GRAPES: Red Skin, Black Skin, Yellow Skin, Purple Skin, Any Combination—1 gal. applied not later than Sept. 7; or 6 lbs. DANCER, plus 1 pound of a DANCER 25 Waikiki, applied not later than Sept. 15.

FEARS: Delayed Dormants For Pests—2 yds. Summer Use—1 to 1½ yds.
Winter Use—(Delayed Dormant and Fall Spray)—2 yds.

ORNAMENTALS AND SHADE TREES (Consult (See Note), Holly, Pyracantha, Sycamorea, Hawthorne, Flowering Peach, Flowering Cherry, Apple, Magnolia, Maple, Dogwood, Oak, Privet, Birch; See Spray Suck, Cydral Shell Scale, Brown Acanth Scale, Black Scale, San Jose Scale, Red Spider Mites and Aphs.—1 gal. to 100 gals. water. Dormant Spray on Deciduous Varieties—1 gal. per 100 gals. water. Dormant Spr'y on Citrus Varieties—1 gal. per 100 gals. water.

Note: May cause discoloration of blue spruce. Do not use as curative or fungicidal on oil spruce.

WALNUTS: Late Spring to Mid-Summer Only; Summer Stages of Froese Scale, European Fruit Scale, San Jose and other Soft-Shelled Scale, Red Spider Mites—2 to 4 quarts. The addition of one of the following phytotoxic insecticides aids in scale control: a Triphor 25 Wettable, a Phosidin 4 Pound Emulsion, an Ecton 28 wettable, or a Parathion 28 Wettable. See label for recommended dosages. Do not apply after black split.

MIXING DIRECTIONS: With agitator running start filling spray tank with water. When combining with Warrantable Powder Insecticide or Fungicide products, add these first when tank is 1/2 filled. Add DAMOL when tank is about 3/4 full and while there is sufficient agitation of the spray to insure complete emulsification.

Do not contain water, food, or food by storage, disposal or cleaning of equipment.

Storage should be under lock and key in a ventilated room and secure from access by unauthorized persons and children. Storage should be in a cool, dry area away from any heat or ignition source. Do not stack over 2 pallets high. Move containers by hand or with a crane. Do not move containers from one area to another unless they are securely sealed. Keep containers tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Store in original containers only. If the contents are leaking or material is spilled follow these steps:

1. Certain spill, absorb with a material such as sawdust, clay granules or dirt.
2. Collect and place in suitable containers for disposal.
3. Wash area with water and soda to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Place a leaking container in a plastic bag and transfer contents as soon as possible to an empty original container.
6. Do not allow runoff to enter sewer or contaminate water supplies.
7. Dispose of waste as indicated below:

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Take items for evaluation. Then either for recycling or reconditioning, or purchase and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, are other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of loss, storage and handling of this material not in exact accordance with directions given hereon.

In no case shall Draxal or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by Draxal Chemical Company and is accepted as such by the Buyer.

BEST AVAILABLE COPY