



MUNICIPALITY OF MEMPHIS
110015 50 00 0000 0000 0000

Plant & Box Seal Treatment
Municipal Plus Molybdenum

6 BUSHES SIZE

ACCEPTED
1982

ACTIVE INGREDIENT:

Captan (N(trichloroethyl) thio-4-cyclohexene-1,2-dicarboximide)	18.86%
INERT INGREDIENTS*	81.14%
	TOTAL: 100.00%

*ADDED Ingredients: Molybdenum (derived from sodium molybdate) - 3.78%

Shake Well Before Each Use

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Immediately flush eyes with plenty of water and get medical attention.

IF ON SKIN: Immediately remove contaminated clothing, INCLUDING SHOES, and wash skin with soap and plenty of water. If irritation develops, send for a physician.

See Side Panel For Additional Precautionary Statements.

EPA Reg. #19713-
EPA Est. #19713-MS-1

Manufactured by: Drexel Chemical Co.
P.O. Box 9306
Memphis, Tenn. 38109

Net Contents: 30 FL. OZ.

Precautionary Statements

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield when handling. Be careful if swallowed. In case of contact, immediately flush eyes with plenty of water and get medical attention promptly. Avoid contamination of food and feed. DO NOT USE TREATED SEED FOR FEED OR OIL PURPOSES. DO NOT USE CONTENTS OF THIS CONTAINER ON LESS THAN 6 BUSHELS OF SEED.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, ponds, and streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed by burial away from bodies of water. Apply this product only as specified on the label.

GENERAL INFORMATION

Molybdenum Plus Captan Flowable is an easy to use liquid soybean seed treatment formulated for planter box application. It is specially formulated to help seed flow easily and evenly from the planter box. Molybdenum Plus Captan Flowable furnishes 1.0 ounce of Captan per bushel to protect against seed and soil-borne diseases, such as seedling blights, damping off and seed decay organisms. This formulation also provides 0.2 ounce of molybdenum per acre for plant nutrition and as an aid in the valuable nitrogen fixation process of the soybean plant.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE EACH USE. Use Molybdenum Plus Captan Flowable at the rate of 5.0 fl. oz. per bushel of soybean seed or the contents of this container on 6 bushels of soybean seed. Plant soybeans at the rate of one (1) bushel per acre for proper application of molybdenum. Fill the planter box (or mixing vessel) with seed, pour the required amount of Molybdenum Plus Captan Flowable evenly on top of the seed. Stir with a stick or paddle so that all seeds are covered. When used in combination with an inoculant, add this product first, stir, add the inoculant, stir. Plant as soon as possible. Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE:

Avoid exposure to excessive heat.

PESTICIDE DISPOSAL:

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

CONTAINER DISPOSAL:

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

WARRANTY - CONDITION OF SALE:

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall Drexel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by Drexel Chemical Company and is accepted as such by the Buyer.