



19713-139

DICOFOL 42 EC

ACTIVE INGREDIENTS:

| | |
|--|----------------------|
| 1,1-Bis (chlorophenyl)-2,2,2-trichloroethanol* | 42.00% |
| Xylene range aromatic solvent. | 45.59% |
| INERT INGREDIENTS: | 12.41% |
| | TOTAL 100.00% |

*Equivalent to 4 lbs. active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
IF IN EYES: Immediately flush eyes with plenty of water and get medical attention.
IF ON SKIN: Immediately remove contaminated clothing, INCLUDING SHOES, and wash skin with soap and plenty of water. If irritation develops, send for a physician.

See Side Panel For Additional Precautionary Statements.

EPA Est. #19713-

Manufactured by: Drexel Chemical Company
P.O. Box 9306
Memphis, Tennessee 38109

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wear regular long-sleeve work clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

GENERAL INFORMATION:

Drexel Dicofol 42 EC is an emulsifiable formulation that gives high initiative kill against most species of agricultural mites. Dicofol 42 EC is a specific miticide and will not kill bees and beneficial insect predators, when used as recommended. Drexel Dicofol 42 EC is compatible with most other commonly used insecticides and fungicides. It should not be combined with lime. Dicofol 42 EC may be applied in dilute or in concentrate ground sprays. For aerial sprays use 3 to 10 gallons of water per acre to provide thorough coverage of all plant parts. Coverage on the underside of leaves is especially important. For further details see specific recommendations on this label, then consult current State Agricultural Experiment Stations spray schedule or State Extension Service Specialists for additional details on compatibility, dosage, timing and uses.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

VEGETABLE AND FIELD CROPS

ALFALFA AND CLOVER (BOTH GROWN FOR SEED ONLY-USE RESTRICTED TO CALIFORNIA, IDAHO, NEVADA, OREGON AND WASHINGTON)-For carmine, Pacific, strawberry (Atlantic), and two-spotted mites. Use 1 to 1½ quarts of Dicofol 42 EC per acre in sufficient water for thorough coverage. Apply when mites become active and repeat if necessary. Treated fields are not to be grazed, and crop residues and cuttings are not to be fed to livestock or baled for such use. The fields must be cleared by burning crop residues or by discing into the soil.

BEANS (dry, green, lima) - Pacific, tropical and two-spotted mites. Use 3/4 to 1 pint Dicofol 42 EC per acre: Apply at first signs of mite buildup and repeat as necessary. Do not apply within 7 days of harvest. Do not feed treated vines (or crop residues) to meat or dairy animals.

BEANS(dry, green, lima) California only--Pacific, tomato russet, tropical and twospotted mites. Use 3 pints Dicofol 42 EC per acre. Apply at first signs of mite buildup and repeat as necessary. Do not apply within 21 days of harvest. Do not feed treated vines (or crop residues) to meat or dairy animals.

COTTON--Desert, Pacific and two-spotted mites. Use 1 to 1½ quarts Dicofol 42 EC per acre, depending on size of plant and density of foliage. Apply in sufficient water to provide thorough coverage. Begin application when mites are first noticed and repeat if necessary but do not apply within 14 days of harvest. Do not feed cotton stalks or trash to meat or dairy animals.

HOPS--Pacific, tropical and two-spotted mites. Use 2 to 2 1/3 pints Dicofol 42 EC per acre. Apply at first signs of mite buildup and repeat as necessary. Do not apply within 7 days of harvest. Do not feed treated vines (or crop residues) to meat or dairy animals.

MINT--Peppermint and spearmint for commercial oil production. For control of two-spotted mites use Dicofol 42 EC at 1 3/4 to 2½ pints per acre in sufficient water for thorough coverage. Make application when mites become active. Do not apply within 30 days of harvest and do not make more than one application per crop per season. Do not feed treated fresh or spent hay to livestock.

FRUITS AND NUTS

APPLES, PEARS (Western States only)--Apple rust, clover, European red, McDaniel, Pacific, Schoene, two-spotted, Willamette and yellow (carpini) mites. Use 3/4 to 1 pint Dicofol 42 EC per 100 gallons dilute spray (1 gallon maximum/acre). Apply in full coverage spray whenever mite population appears and repeat at 10 to 14 day intervals as necessary. Do not apply within 7 days of harvest. To avoid injury do not use glyodin fungicide with Dicofol 42 EC in late cover sprays.

APRICOTS, CHERRIES, FIGS, NECTARINES, PEACHES, PLUMS, PRUNES--Clover (brown almond), European red, McDaniel, Pacific, peach silver, plum nursery, Schoene, two-spotted, Willamette and yellow mites. Use 3/4 to 1 pint Dicofol 42 EC per 100 gallons dilute spray (6.4 pints maximum/acre for Apricots, Cherries, Nectarines and Peaches and 4.0 pints maximum/acre for Figs, Plums and Prunes). Apply in full coverage spray at petal fall to first cover or whenever mite population appears. Dicofol 42 EC may cause fruit spotting on Figs. When fruit is present use a Dicofol 35 wettable formulation. For repeat application on Cherries, Figs, Plums and Prunes wait 30 days to avoid excessive residues, but do not apply within 7 days of harvest. On Apricots, Nectarines and Peaches do not apply within 14 days of harvest.

CITRUS (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangelos, Tangerines) Citrus flat, citrus red (purple), citrus rust, six-spotted, Texas citrus and Yuma mites. Apply Dicofol 42 EC in full coverage sprays using 1 pint per 100 gallons of dilute spray (2 gallons maximum/acre). If concentrate sprays are used adjust dosage accordingly. Apply when mites appear in

threatening numbers and repeat as required for satisfactory control. Do not apply within 7 days of harvest.

Do not use Dicofol in any formulation on avocados or eggplant.

GRAPES--For clover (brown almond), European red, McDaniel, Pacific, two-spotted, Willamette and yellow mites. Use 1 1/3 to 2 1/3 pints Dicofol 42 EC per acre. Apply just after bloom or whenever mites appear and repeat if necessary. Do not apply within 7 days of harvest.

STRAWBERRIES--Six-spotted, strawberry (Atlantic) and two-spotted mites. Use 1 quart Dicofol 42 EC per acre. Apply when mites appear and repeat as necessary but do not apply within 2 days of harvest, or if temperature exceeds 90°F.

WALNUTS --For clover (brown almond), European red, McDaniel, Pacific, two-spotted, Willamette and yellow mites. Use a full coverage spray of 3/4 to 1 pint Dicofol 42 EC per 100 gallons dilute spray (1 gallon maximum/acre). Apply just after bloom or whenever mites appear and repeat if necessary. Do not apply within 14 days of harvest.

NOTE: Dicofol 42 EC may be used without any spreader-sticker. When necessary to improve the spray coverage on foliage difficult to wet and to smooth out spray deposits on maturing fruit Triton B-1956 is recommended. Other suitable pesticide spreader-stickers may also be used except in situations where Triton B-1956 is recommended as the only acceptable spreader-sticker.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

WARRANTY - CONDITION OF SALE:

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall Draxel or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by Draxel Chemical Company and is accepted as such by the Buyer.

| | | |
|--|---------------------------|------------------|
| <p align="center"> U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) WASHINGTON, D.C. 20460 </p> <p> NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i> </p> | EPA REGISTRATION NO. | DATE OF ISSUANCE |
| | TERM OF ISSUANCE | |
| | NAME OF PESTICIDE PRODUCT | |
| NAME AND ADDRESS OF REGISTRANT (Include ZIP code) | | |
| <p> NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number. </p> <p> On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act. </p> <p> A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith. </p> <p> Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others. </p> | | |
| <input type="checkbox"/> ATTACHMENT IS APPLICABLE | | |
| SIGNATURE OF APPROVING OFFICIAL | | DATE |